

SIEVE WEIGHTS AND THEIR SMOOTHINGS

ANDREW GRANVILLE, DIMITRIS KOUKOULOPOULOS, AND JAMES MAYNARD

ABSTRACT. We obtain asymptotic formulas for the $2k$ th moments of partially smoothed divisor sums of the Möbius function. When $2k$ is small compared with A , the level of smoothing, then the main contribution to the moments come from integers with only large prime factors, as one would hope for in sieve weights. However if $2k$ is any larger, compared with A , then the main contribution to the moments come from integers with quite a few prime factors, which is not the intention when designing sieve weights. The threshold for “small” occurs when $A = \frac{1}{2k} \binom{2k}{k} - 1$.

One can ask analogous questions for polynomials over finite fields and for permutations, and in these cases the moments behave rather differently, with even less cancellation in the divisor sums. We give, we hope, a plausible explanation for this phenomenon, by studying the analogous sums for Dirichlet characters, and obtaining each type of behaviour depending on whether or not the character is “exceptional”.

1. INTRODUCTION

Sieve methods begin with the identity

$$(1.1) \quad \sum_{d|(a,m)} \mu(d) = \begin{cases} 1 & \text{if } (a, m) = 1, \\ 0 & \text{otherwise,} \end{cases}$$

since then one has

$$\#\{a \in \mathcal{A} : (a, m) = 1\} = \sum_{a \in \mathcal{A}} \sum_{d|(a,m)} \mu(d) = \sum_{d|m} \mu(d) \sum_{\substack{a \in \mathcal{A} \\ d|a}} 1$$

for any finite set of integers $\mathcal{A}$. Typically, we take $m = \prod_{p \leq z} p$ for some parameter z , though m can be chosen differently.

This simple approach rarely leads to asymptotics (unless m is small), since the combined error terms when estimating the final sum on the right-hand side usually add up to more than the main term. To circumvent this difficulty one restricts the sum to only smallish d (that is we *truncate* the sum in (1.1)), but at what cost? The truncation of (1.1) must be accomplished in a way that simultaneously achieves two goals: the new sum is supported on divisors $d \leq R$, for some suitable parameter R , and the new sum still correlates sufficiently with the characteristic function of integers that are coprime to m . There are many

Date: September 13, 2016.

2010 *Mathematics Subject Classification.* Primary: 11N35, 11N37; Secondary: 11T06, 20B30, 05A05.

Our thanks go to Maksym Radziwiłł for supplying several key references, and to him, Ben Green and Henryk Iwaniec for helpful discussions. AG is funded by the National Science and Engineering Research Council of Canada and by the European Research Council. DK is funded by the National Science and Engineering Research Council of Canada and by the Fonds de recherche du Québec – Nature et technologies. JM is funded by a Clay research fellowship and a fellowship by examination of Magdalen College, Oxford.

ways of doing so. Brun, the pioneer of modern sieve methods, used the inclusion-exclusion inequalities

$$\sum_{\substack{d|(a,m) \\ \omega(d) \leq 2r-1}} \mu(d) \leq \sum_{d|(a,m)} \mu(d) \leq \sum_{\substack{d|(a,m) \\ \omega(d) \leq 2r}} \mu(d),$$

for any $r \in \mathbb{N}$, which when summed over m , often work with a judicious choice of r . In the more general combinatorial sieve one chooses sets $\mathcal{D}^-, \mathcal{D}^+ \subset \{d|m : d \leq R\}$, tailored to the problem at hand, for which

$$\sum_{\substack{d|(a,m) \\ d \in \mathcal{D}^-}} \mu(d) \leq \sum_{d|(a,m)} \mu(d) \leq \sum_{\substack{d|(a,m) \\ d \in \mathcal{D}^+}} \mu(d).$$

See [8, ch. 6] for details on how this can be done.

In [19], Selberg introduced a new approach to sieving. His starting point was the inequality

$$\sum_{d|(a,m)} \mu(d) \leq \left(\sum_{d|(a,m)} \lambda_d \right)^2,$$

which is valid for any real numbers λ_d with $\lambda_1 = 1$. The best upper bound stemming from this identity comes from minimizing the quadratic form

$$\sum_{a \in \mathcal{A}} \left(\sum_{d|(a,m)} \lambda_d \right)^2 = \sum_{d_1, d_2 | m} \lambda_{d_1} \lambda_{d_2} \cdot \#\{a \in \mathcal{A} : [d_1, d_2] | a\}.$$

We will assume that the parameters λ_d are only supported on square-free integers $d \leq R$, so that we only require information about the distribution of $\mathcal{A}$ in arithmetic progressions with step size $\leq R^2$. We make the additional simplifying assumption that $m = \prod_{p \leq z} p$ and that $R \geq z$, so that the condition $d|m$ is guaranteed for square-free integers $d \leq R$. For typical sets $\mathcal{A}$ that arise in arithmetic problems, we find that the optimal choice for the λ_d takes the form

$$\lambda_d \approx c \cdot \mu(d) \cdot \left(\frac{\log(R/d)}{\log R} \right)^A \quad (d \leq R),$$

where c is a suitable normalization constant possibly depending on R and A a sufficiently large exponent. We note that the weights λ_d decay to 0, and the larger the value of A , the higher the level of smoothness at the truncation point R . In the optimal choice, the exponent A is taken to be κ , the *dimension* of the sieve problem. However, for a given dimension κ , it is known [20, pg. 154] that any exponent $A > \kappa - 1/2$ yields weights λ_d whose dominant contribution comes from numbers almost coprime to m , whereas this fails to be true for smaller A . Go to [8, ch. 10] for further discussion.

More generally, one can consider the smoothed sieve weight

$$M_f(a; R) := \sum_{d|a} \mu(d) f\left(\frac{\log d}{\log R}\right),$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function supported on $(-\infty, 1]$. In Selberg sieve arguments one typically takes f to be a polynomial in $[0, 1]$, perhaps of high degree. Such an example is offered by the GPY sieve [11, 24]. However in more recent developments on gaps between primes [16, 21] one works with arbitrarily smooth functions f .

The main motivation of this paper is to understand the exact role of the smoothing in the structure of the Selberg sieve weights. To this end, we consider their moments

$$\sum_{n \leq x} M_f(a; R)^k$$

as a tool of gaining additional insight on the distribution of the values of $M_f(a; R)$. On the practical side, higher moments naturally appear when applying Hölder's inequality, so it would be useful to know their behaviour. Expanding the k -th power, we find that

$$\sum_{n \leq x} M_f(n; R)^k = x \cdot \mathcal{M}_{f,k}(R) + O((\|f\|_\infty R)^k)$$

for any $f : \mathbb{R} \rightarrow \mathbb{R}$ supported on $(-\infty, 1]$, where

$$(1.2) \quad \mathcal{M}_{f,k}(R) := \sum_{d_1, \dots, d_k \geq 1} \frac{\prod_{j=1}^k \mu(d_j) f(\log d_j / \log R)}{[d_1, \dots, d_k]} = \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{p|n \Rightarrow p \leq R} \frac{M_f(n; R)^k}{n}.$$

How smooth should f be so that $M_f(a; R)^{2k}$ behaves like a sieve weight when k varies, that is to say the main contribution to the $2k$ -th moment¹ of $M_f(a; R)$ comes from integers a that have very few prime factors $\leq R$? What happens in the extreme case where f is the discontinuous function $\mathbf{1}_{(-\infty, 1]}$? These are the types of questions that we will study in this paper.

1.1. Some smoothing is necessary. In order to gain a first understanding of the importance of smoothing, let us consider the sharp cut-off function

$$f_0 := \mathbf{1}_{(-\infty, 1]}.$$

If $n = 2m$ with m odd, then we have that

$$(1.3) \quad M_{f_0}(n; R) = \sum_{\substack{d|n \\ d \leq R}} \mu(d) = \sum_{\substack{d|m \\ d \leq R}} \mu(d) + \sum_{\substack{d|m \\ 2d \leq R}} \mu(2d) = \sum_{\substack{d|m \\ R/2 < d \leq R}} \mu(d) = M_{\tilde{f}_0}(m; R),$$

where, with a slight abuse of notation, we have put

$$(1.4) \quad \tilde{f}_0 := \mathbf{1}_{(1 - \log 2 / \log R, 1]}.$$

In particular, if m is square-free and has exactly one divisor $d \in (R/2, R]$, then $M_{f_0}(n; R) = \pm 1$. An easy generalization of a deep result of Ford [7, Theorem 4] implies that the proportion of such $m \leq x/2$ is $\gg (\log R)^{-\delta} (\log \log R)^{-3/2}$ with

$$\delta = 1 - \frac{1 + \log \log 2}{\log 2} = 0.086071332 \dots,$$

whence we conclude that

$$\#\{n \leq x : M_{f_0}(n; R) \neq 0\} \gg \frac{x}{(\log R)^\delta (\log \log R)^{3/2}} \quad (x \geq R^{1+\epsilon}).$$

In particular, we find that $M_{f_0}(n; R)$ is non-zero too often to behave like a sieve weight. This indicates that part of the importance of smoothing is to reduce the contribution of isolated divisors of n to $M_f(n; R)$.

¹We are typically interested in how large sieve weights get. If we took odd powers there might be an irrelevant cancelation, so we focus on even moments.

We will prove in Section 5 that the

$$(1.5) \quad \#\{n \leq x : M_{f_0}(n; R) \neq 0\} \asymp \frac{x}{(\log R)^\delta (\log \log R)^{3/2}} \quad (x \geq R^4).$$

This sharpens a result by Hall and Tenenbaum [10], who used a very similar argument and the best results about divisors of integers available at that time.

1.2. A heuristic argument. Going back to the study of $M_f(n; R)$ for a smooth function f , it is reasonable to believe that the smoother f is, the larger the k are for which $M_f(n; R)^{2k}$ behaves like a sieve weight. One way to explain this phenomenon is by noticing that various integral transformations have faster decay for smooth weights. (See, for example, Section 6.) Nevertheless, we prefer to give a number theoretic explanation in terms of the underlying sieve questions rather than an analytic one focused more on the technical issues: assume that $n = p_1^{\alpha_1} \cdots p_r^{\alpha_r} m$, where $p_1 < \cdots < p_r$ and all of the prime divisors of m are $> p_r$. Then

$$(1.6) \quad \begin{aligned} M_f(n; R) &= \sum_{d|p_2 \cdots p_r m} \mu(d) f\left(\frac{\log d}{\log R}\right) + \sum_{d|p_2 \cdots p_r m} \mu(p_1 d) f\left(\frac{\log(p_1 d)}{\log R}\right) \\ &= \sum_{d|p_2 \cdots p_r m} \mu(d) \left\{ f\left(\frac{\log d}{\log R}\right) - f\left(\frac{\log p_1}{\log R} + \frac{\log d}{\log R}\right) \right\}. \end{aligned}$$

Continuing as above, we find that

$$(1.7) \quad M_f(n; R) = (-1)^r \sum_{d|m} \mu(d) \Delta^{(r)} f\left(\frac{\log d}{\log R}; \frac{\log p_1}{\log R}, \dots, \frac{\log p_r}{\log R}\right),$$

where $\Delta^{(r)} f(x; h_1, \dots, h_r)$ denotes the multi-difference operator defined by

$$\Delta^{(1)} f(x; h) = f(x + h) - f(x)$$

and

$$\Delta^{(r)} f(x; h_1, \dots, h_r) = \Delta^{(r-1)}(x + h_r; h_1, \dots, h_{r-1}) - \Delta^{(r-1)}(x; h_1, \dots, h_{r-1}).$$

In particular, if $f \in C^r(\mathbb{R})$, then

$$(1.8) \quad \Delta^{(r)} f(x; h_1, \dots, h_r) = \int_0^{h_r} \int_0^{h_{r-1}} \cdots \int_0^{h_1} f^{(r)}(x + t_1 + t_2 + \dots + t_r) dt_1 \dots dt_r,$$

Returning to (1.7), we see that if $f \in C^A(\mathbb{R})$ and $n = p_1^{\alpha_1} \cdots p_r^{\alpha_r} m$, $r \leq A$, is as above, then $M_f(n; R)$ should heuristically be $\lesssim M_{f^{(r)}}(m; R) \prod_{j=1}^r (\log p_j / \log R)$. Loosely, this indicates each additional degree of smoothness of the weight function f cuts the average size of $M_f(n; R)$ by about a factor of $1/\log R$.

The above discussion leads us to conjecture that if $f \in C^A(\mathbb{R})$ with $f(0) \neq 0$, then

$$(1.9) \quad \sum_{n \leq x} M_f(n; R)^{2k} \ll \max \left\{ \frac{x}{\log R}, \frac{1}{(\log R)^{2kA}} \sum_{n \leq x} M_{f_0}(n; R)^{2k} \right\}.$$

Notice that the factor $x/\log R$ is necessary because $M_f(n; R) = f(0)$ for all integers n that are free of prime factors $\leq R$.

Naturally, for this relation to be useful, we need to understand the asymptotics of $\sum_{n \leq x} M_{f_0}(n; R)^{2k}$. Recall the relation (1.2). When $k = 1$, Dress, Iwaniec and Tenenbaum [4] showed

$$(1.10) \quad \mathcal{M}_{f_0,2}(R) \sim c_1 \quad (R \rightarrow \infty)$$

for some constant $c_1 > 0$, and when $k = 2$, Motohashi [17] showed that

$$(1.11) \quad \mathcal{M}_{f_0,4}(R) \sim c_2(\log R)^2 \quad (R \rightarrow \infty)$$

for some constant $c_2 > 0$. In general, Balazard, Naimi, and Pétermann [1] proved that

$$\mathcal{M}_{f_0,2k}(R) = P_k(\log R) + O(e^{-c(\log R)^{3/5}(\log \log R)^{-1/5}}),$$

for some polynomial P_k and some constant $c = c(k) > 0$. This built on work of de la Bréteche [2], who showed how a wide class of related sums can be evaluated asymptotically. However, when applying his technique to this question, one would need some strong understanding of the growth of $\zeta(s)$ near to $s = 1$ to recover the result of [1] (which, for example, follows from the Riemann Hypothesis).

Notice that if $\mathcal{E}_k = \deg(P_k)$, so that $\mathcal{E}_1 = 0$ and $\mathcal{E}_2 = 2$, then (1.9) becomes

$$(1.12) \quad \sum_{n \leq x} M_f(n; R)^{2k} \ll \max \left\{ \frac{x}{\log R}, x(\log R)^{\mathcal{E}_k - 2kA} \right\}.$$

This suggests that $M_f(n; R)^{2k}$ acts like a sieve weight as long as $A > \mathcal{E}_k/2k$. The big issue with the result of Balazard, Naimi and Pétermann is that the degree $\mathcal{E}_k$ is not determined for general k , and that is essential if one wishes to gain a better understanding of how the Selberg sieve weights work. Our attention thus turns to calculating $\mathcal{E}_k$.

But first, we study seemingly analogous questions (in different settings), that one might guess would be easier and indicate what kind of estimate we should be looking for

1.3. Analogous settings. It is well-known that many of the analytic properties of integers are shared by both polynomials of finite fields (c.f. [18]), and by permutations (c.f. [9]). Moreover, polynomials and permutations are usually easier objects to understand, so in order to gain an understanding of the exponent $\mathcal{E}_k$, it would be natural to consider what happens in these analogous settings first.

Permutations. The easiest analogy to analyze concerns permutations. Every $\sigma \in S_N$ (the permutations on N letters) can be decomposed in a unique way into a product of cycles. Those cycles cannot be decomposed any further and play the role of irreducibles. Divisors of σ are precisely the set of possible products of cycles. If those cycles act on the subset T of $[N]$, then σ fixes T . Moreover, if σ fixes T , then σ is a product of cycles, a subproduct of which fixes T . Hence “divisors” correspond to sets $T \subset [N]$ for which $\sigma(T) = T$.

To “calibrate” our understandings of the properties of integers and permutations, we note that for a typical integer n , its j -th largest prime factor is about e^{e^j} , whereas for a typical permutation $\sigma \in S_N$, its j -th largest cycle has length about e^j . Thus, the inequality $R/2 < d \leq R$ for a divisor d of n corresponds to having a set T that is fixed by σ of size $|T| = \log R + O(1)$. Hence we will study

$$(1.13) \quad \text{Perm}(N, m; k) := \frac{1}{N!} \sum_{\sigma \in S_N} \left| \sum_{\substack{T \subset [N], |T|=m, \\ \sigma(T)=T}} \mu(\sigma|_T) \right|^{2k},$$

where

$$[N] := \{1, \dots, N\}$$

and if $\sigma|_T = C_1 C_2 \dots C_\ell$ is the product of ℓ cycles, then we have set $\mu(\sigma|_T) = (-1)^\ell$. We claim that $\text{Perm}(N, m; k)$ is more natural than it appears at first sight. A usual function

of permutations is the signature $\epsilon(\sigma)$ which counts the number of transpositions (i.e. the number of interchanges of two elements) needed to create σ . For a cycle C , one knows that $\epsilon(C) = (-1)^{|C|-1}$ and hence $\epsilon(\sigma|_T) = \epsilon(C_1)\epsilon(C_2)\dots\epsilon(C_\ell) = (-1)^{|T|-\ell} = (-1)^m \mu(\sigma|_T)$, since $|T| = m$ here. Therefore

$$\sum_{\substack{\sigma \in S_N, T \subset [N], \\ |T|=m, \sigma(T)=T}} \mu(\sigma|_T) = (-1)^m \sum_{\substack{\sigma \in S_N, T \subset [N], \\ |T|=m, \sigma(T)=T}} \epsilon(\sigma|_T),$$

whence

$$\text{Perm}(N, m; k) = \frac{1}{N!} \sum_{\sigma \in S_N} \left| \sum_{\substack{T \subset [N], |T|=m, \\ \sigma(T)=T}} \epsilon(\sigma|_T) \right|^{2k}.$$

Arguing as in [5], the analogy for permutations of Ford's results [7] for integers, it is possible to show that the summands on the right hand of (1.13) (and, hence, of the above formula) are non-zero for a proportion $\asymp 1/m^\delta (\log m)^{3/2}$ of the permutations in S_N . The following theorem provides a formula and an asymptotic estimate for $\text{Perm}(N, m; k)$.

Theorem 1.1. *For each integer $k \geq 1$ and each integer $m \geq 1$, if $N \geq 2mk$ then*

$$\text{Perm}(N, m; k) = c(m, k),$$

where $c(m, k)$ is the number of $(2^{2k} - 1)$ -tuples $(r_I)_{\emptyset \neq I \subset \{1, \dots, 2k\}}$ of non-negative integers such that

- $r_I \in \{0, 1\}$ for $|I|$ odd;
- $\sum_{I: i \in I} r_I = m$, for each $i \in \{1, \dots, 2k\}$.

Moreover, for fixed $k \in \mathbb{N}$, the function $c(m, k)$ is increasing in m and satisfies the estimate

$$c(m, k) \asymp_k m^{2^{2k-1} - 2k - 1} + 1.$$

Proof of the formula for $\text{Perm}(N, m; k)$. If $T_1, \dots, T_{2k}$ are fixed sets of a permutation $\sigma \in S_N$, then the sets

$$R_I := \left(\bigcap_{i \in I} T_i \right) \setminus \left(\bigcup_{i \in [2k] \setminus I} T_i \right) \quad (I \subset [2k])$$

are all fixed by σ and they provide a partition of $[N]$, that is to say $[N]$ equals $\sqcup_I R_I$, the disjoint union of the sets R_I . We thus find

$$\frac{1}{N!} \sum_{\sigma \in S_N} \left| \sum_{\substack{T \subset [N], |T|=m, \\ \sigma(T)=T}} \epsilon(\sigma|_T) \right|^{2k} = \sum_{\substack{r_I \geq 0 \ \forall I \\ \sum_{I: i \in I} r_I = m}} \sum_{\substack{[N] = \sqcup_I R_I \\ |R_I| = r_I \ \forall I}} \frac{1}{N!} \prod_{I \subset [2k]} \sum_{\rho_I \in S_{R_I}} \epsilon(\rho_I)^{|I|}.$$

The inner sums are each $r_I!$ unless $|I|$ is odd and $r_I > 1$, in which case we get 0. Additionally, we get that the number of choices of sets of the given sizes is $N! / \prod_I r_I!$, and hence the above equals $c(m, k)$.

The upper bound for $c(m, k)$ follows from a straightforward lattice point count. The lower bound will be proven in Section 3. $\square$

Evidently, the above results suggest that $\mathcal{E}_k = \max\{0, 2^{2k-1} - 2k - 1\}$. Relation (1.10) implies that $\mathcal{E}_1 = 0$, but relation (1.11) implies that $\mathcal{E}_2 = 2 \neq 2^3 - 5$. This suggests that there is a discrepancy between the integer and the permutation setting, a very rare exception.

Polynomials over finite fields. Positive integers are uniquely identifiable by their factorization into primes (the Fundamental Theorem of Arithmetic). Note though that every non-zero integer equals a unit (that is 1 or -1) times one of those positive integers. We will work with polynomials in $\mathbb{F}_q[t]$. Monic polynomials in $\mathbb{F}_q[t]$ are uniquely identifiable by their factorization into monic irreducible polynomials of degree ≥ 1 . Again, note that every non-zero polynomial in $\mathbb{F}_q[t]$ equals a unit (that is, any element $a \in \mathbb{F}_q \setminus \{0\}$) times a monic polynomial. We will work only with monic polynomials, for example when considering divisors of a given polynomial (rather like we only consider positive integer divisors of a given integer). The Möbius function of a given polynomial is a multiplicative function, where $\mu(P) = -1$, and $\mu(P^k) = 0$ if $k \geq 2$, whenever P is irreducible.

To “calibrate” our understandings of the arithmetic properties of integers and polynomials, we note that $\sim 1/\log x$ of integers around x are prime, whereas $\sim 1/m$ of monic polynomials of degree m are irreducible in $\mathbb{F}_q[t]$. Here the “ $\sim$ ” symbol means as $q \rightarrow \infty$ running through prime powers. Thus, wherever we see $\log x$ in an estimate about the integers, we try to replace it with m in an estimate about degree m polynomials. Similarly a divisor d of n that is close to R is analogous to a polynomial divisor of $F(t)$ of degree m , where m replaces $\log R$ in estimates. Hence we will study

$$\text{Poly}_q(n, m; k) := \frac{1}{q^n} \sum_{\substack{N \in \mathbb{F}_q[t] \\ \deg N = n}} \left| \sum_{\substack{M|N \\ \deg M = m}} \mu(M) \right|^{2k},$$

where M and N are monic, and we have divided by q^n because this is how many monic polynomials N of degree n are contained in $\mathbb{F}_q[t]$, which is the analogue of

$$\frac{1}{x} \sum_{n \leq x} \left(\sum_{\substack{d|n \\ R/2 < d \leq R}} \mu(d) \right)^{2k},$$

a quantity directly related to $\frac{1}{x} \sum_{n \leq x} M_{f_0}(n; R)^{2k}$ via (1.3). We will prove below the following estimate:

Theorem 1.2. *For integers $k, m \geq 1$ and $n \geq 2mk$, we have that*

$$\text{Poly}_q(n, m; k) = c(m, k)(1 + O_k(1/q)) \asymp_k 1 + m^{2^{2k-1}-2k-1}.$$

We thus see that polynomials behave similarly to permutations (and thus differently than integers).

1.4. Two worlds apart and a bridge between them. Our discussion of the permutation and polynomial analogues, rather than shedding more light on the value of the exponent $\mathcal{E}_k$, gave rise to even more questions. It turns out that the integer setting is substantially more complicated than the permutation and polynomial settings. We now state our main results about integers. First, given $A \in \mathbb{Z}_{\geq 1}$, we let

$$f_A(t) := \begin{cases} (1-t)^A & \text{for } t \leq 1; \\ 0 & \text{otherwise,} \end{cases}$$

an extension of the definition of f_0 . Note that $f_A \in C^{A-1}(\mathbb{R}) \setminus C^A(\mathbb{R})$ for all $A \geq 1$. We then have the following result that determines the value of $\mathcal{E}_k$:

Theorem 1.3. *For fixed integers $k \geq 1$ and $A \geq 0$, there is a constant $c_{k,A} > 0$ such that*

$$(1.14) \quad \mathcal{M}_{f_A, 2k}(R) = c_{k,A}(\log R)^{\mathcal{E}_{k,A}} + O((\log R)^{\mathcal{E}_{k,A}-1}),$$

where

$$\mathcal{E}_{k,A} := \max \left\{ \binom{2k}{k} - 2k(A+1), -1 \right\}.$$

In particular, $\mathcal{E}_k = \mathcal{E}_{k,0} = \binom{2k}{k} - 2k$. Additionally, we find that there is a constant $c'_k > 0$ such that for $R^{2k} \leq x$ we have

$$(1.15) \quad \frac{1}{x} \sum_{n \leq x} \left(\sum_{\substack{d|n \\ R/2 < d \leq R}} \mu(d) \right)^{2k} = c'_k (\log R)^{\binom{2k}{k} - 2k} + O\left((\log R)^{\binom{2k}{k} - 2k - 1}\right).$$

All implied constants depend at most on k and A .

Remark 1.1. We have no nice formula for the constants $c_{k,A}$ and c'_k appearing in Theorem 1.3; we only know how to write them as an enormous rational linear combination of complicated integrals, and leave it as a challenge to come up with an easy explicit description.

Remark 1.2. If the moments of a distribution grow slowly, then the distribution can be determined via its Laplace transform. However, in our case the moments are of rapidly increasing magnitude, indeed with different powers of $\log R$, so one cannot immediately deduce from them the distribution of the weights $M_{f_A}(n; R)$ as n varies over the integers.

For general functions f , we prove that $M_f(n; R)^{2k}$ behaves like a sieve weight as long as $f \in C^A(\mathbb{R})$ with $A > \binom{2k}{k}/2k = \mathcal{E}_{2k}/2k + 1$. Notice that this confirms a weak version of the heuristic estimate (1.12).

Theorem 1.4. *Let $k \in \mathbb{N}$ and $f : \mathbb{R} \rightarrow \mathbb{R}$ be supported in $(-\infty, 1]$. Assume further that $f \in C^A(\mathbb{R})$ and that all functions $f, f', \dots, f^{(A)}$ are uniformly bounded for some integer $A \geq 2$, and fix some $\epsilon \in (0, 1)$.*

(a) *Assume that $A > \frac{1}{2k} \binom{2k}{k}$. For $x \geq R \geq 2$ and $\log 2 / \log R \leq \eta \leq 1 \geq 1$, we have that*

$$\sum_{\substack{n \leq x \\ \exists p|n, p \leq R^\eta}} M_f(n; R)^{2k} \ll \frac{\eta x}{\log R}.$$

If, in addition, $f(0) \neq 0$, then there is a constant $c_{k,f} > 0$ such that for $x \geq R^{2k} \log^2 R$ we have that

$$\frac{1}{x} \sum_{n < x} M_f(n; R)^{2k} = \frac{c_{k,f}}{\log R} + O\left(\frac{1}{(\log R)^{2-\epsilon}}\right).$$

(b) *If $A \leq \frac{1}{2k} \binom{2k}{k}$, then for $x \geq R \geq 2$ we have that*

$$\sum_{n \leq x} M_f(n; R)^{2k} \ll x (\log R)^{\binom{2k}{k} - 2kA}.$$

All implied constants depend at most on f , k and ϵ .

The value of $\mathcal{E}_k = \binom{2k}{k} - 2k$ given by Theorem 1.3 is significantly smaller than the exponent $2^{2k-1} - 2k - 1$ in the polynomial/permutation setting. So we see the usual analogy breaking down in quite a severe way, something surprising. We devote Section 2 to the analysis of this discrepancy. In particular, we will see that the underlying reason is the relation

$$(1.16) \quad \sum_{\substack{\sigma \in S_N, \ T \subset [N], \\ |T|=m, \ \sigma(T)=T}} \mu(\sigma|_T) = (-1)^m \sum_{\substack{\sigma \in S_N, \ T \subset [N], \\ |T|=m, \ \sigma(T)=T}} \epsilon(\sigma|_T)$$

that we saw before. Notice here that while $\mu(\rho) = -1$ for all cycles ρ , we have that $\epsilon(\rho)$ takes the values ± 1 with equal probability as ρ ranges over cycles of all possible lengths. The simplest example of a multiplicative function over $\mathbb{Z}$ demonstrating this kind of behaviour is that of a Dirichlet character. To this end, we consider

$$\mathcal{X}_{2k}(R) = \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{1}{n} \left(\sum_{\substack{d|n \\ R/2 < d \leq R}} \chi(d) \right)^{2k},$$

which, as in (1.2), is the main term of

$$\frac{1}{x} \sum_{n < x} \left| \sum_{\substack{d|n \\ R/2 < d \leq R}} \chi(d) \right|^{2k}.$$

We then have the following theorem, which shows that it is possible to bridge the gap between the two worlds of integers and of permutations/polynomials. All implied constants below depend at most on k , and we have set

$$\mathcal{S}^+(2k) := \{I \subset \{1, 2, \dots, 2k\} : |I| \text{ even}\} \setminus \{\emptyset\}.$$

Theorem 1.5. *Let $\chi \pmod{q}$ be a real non-principal character and $k \in \mathbb{N}$.*

(a) *If $k = 1$, then*

$$\mathcal{X}_2(R) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{P(t, \chi) |L(1 + it, \chi)|^2 \sin^2(t(\log 2)/2)}{t^2} dt + O\left(\frac{1}{(\log R)^{100}}\right),$$

where $P(\cdot, \chi)$ is a real-valued Euler product whose factors are $1 + O(1/p^2)$. In particular, $P(t, \chi) \asymp 1$ for all t , uniformly in χ .

(b) *Assume that $k \geq 2$, let $V_k(m)$ be the Lebesgue volume of the polytope $\{(x_I)_{I \in \mathcal{S}^+(2k)} : x_I \geq 0, \ m - \log 2 \leq \sum_{I \ni i} x_I \leq m\}$ in $\mathbb{R}^{2^{2k-1}-1}$, and let $\mathfrak{S}_k(\chi)$ be the singular series*

$$\mathfrak{S}_k(\chi) = \prod_p \left(1 - \frac{1}{p}\right)^{2^{2k-1}} f_p,$$

where $f_p = \sum_{j \geq 0} (j+1)^{2k}/p^j$ if $\chi(p) = 1$, $f_p = (1 - 1/p^2)^{-1}$ if $\chi(p) = -1$, and $f_p = (1 - 1/p)^{-1}$ if $p|q$. Then $V_k(m) \asymp_k m^{2^{2k-1}-2k-1}$, $\mathfrak{S}_k(\chi) \asymp_k L(1, \chi)^{2^{2k-1}}$, and

$$\mathcal{X}_{2k}(R) = \mathfrak{S}_k(\chi) \cdot V_k(\log R) + O\left((\log R)^{2^{2k-1}-2k-2} (\log(q \log R))^{O(1)}\right).$$

(c) Assume that $k \geq 2$ and that $L(\beta, \chi) = 0$ for some $\beta > 1 - 1/(100 \log q)$. If $Q = e^{1/(1-\beta)}$ and $e^{(\log q)^C} \leq R \leq Q$ for some large enough $C = C(k)$, then there is a constant $c_k(\chi) = (\log q)^{O(1)}$ such that

$$\mathcal{X}_{2k}(R) = c_k(\chi)(\log R)^{\binom{2k}{k} - 2k} \left(1 + O \left(\frac{(\log(q \log R))^{O(1)}}{\log R} \right) \right).$$

In the case of our polynomial and permutation models, we have an exponent of $2^{2k-1} - 2k - 1$ for the $2k^{\text{th}}$ moment with $k \geq 2$, whilst over the integers we have an exponent $\binom{2k}{k} - 2k$. We see that our Dirichlet character model interpolates between these two settings. If the Dirichlet L -function associated with the character has a zero very close to 1, then $\chi(p) = -1$ for many small primes p , and so by multiplicativity χ behaves similarly to μ (at least in appropriate ranges). This is represented by our exponent of $\binom{2k}{k} - 2k$ in this case. On the other hand, χ is a periodic character, and if the L function does not have a zero very close to 1, we see that we have an exponent $2^{2k-1} - 2k - 1$, matching the exponent of our polynomial and permutation models. Notice that if $L(s, \chi)$ does have an exceptional zero, then the asymptotic of case (c) for $\mathcal{X}_{2k}(R)$ holds for small R , and transitions into the asymptotic of case (b) as R grows.

Remark 1.3. Relation (1.16) has a polynomial analogue whose consequences are worth exploring further. Given $I \subset \{F \in \mathbb{F}_q[t] : \deg(F) = n\}$, we consider the sum

$$\sum_{F \in I} \mu(F).$$

For example, we could take $I = \{F \in \mathbb{F}_q[t] : \deg(F) = n\}$, or $I = \{F \in \mathbb{F}_q[t] : \deg(F - F_0) \leq h\}$ for some $F_0 \in \mathbb{F}_q[t]$ of degree n and for some integer $h \in [1, n-1]$, which can be seen as the polynomial analogue of a short interval. Then

$$\sum_{F \in I} \mu(F) = (-1)^n \sum_{F \in I} \chi(F),$$

where $\chi(F) = (-1)^{\deg(F)} \mu(F)$, which is also a multiplicative function. However, we note that, even though $\mu(P) = -1$ for all irreducibles, we have that $\chi(P) = 1$ for about half of the irreducibles P , and $\chi(P) = -1$ for the other half, that is to say χ behaves on average much more like a real Dirichlet character rather than the Möbius function.

This phenomenon is striking and sharply different than what happens over $\mathbb{Z}$, where there is a dichotomy between multiplicative functions that look like the Möbius functions and other ones whose average prime value is 0, as is exemplified by Theorem 1.5 (see, also, [15]).

1.5. Further analysis of truncated Möbius divisor sums. As we saw in Theorem 1.4, if $f \in C^A(\mathbb{R})$ with $A > \frac{1}{2k} \binom{2k}{k} = \mathcal{E}_k/2k + 1$, then $M_f(n; R)^{2k}$ behaves like a sieve weight. When $f = f_A$, we can be more precise:

If $A > \frac{1}{2k} \binom{2k}{k} - 1 = \mathcal{E}_k/2k$, then $\mathcal{E}_{k,A} = -1$ in Theorem 1.3, and so $\mathcal{M}_{f_A, 2k}(R) \asymp (\log R)^{-1}$. Since integers $n \leq x$ with no prime factors less than R contribute a total $\gg x(\log R)^{-1}$ to $\sum_{n \leq x} M_{f_A}(n; R)^{2k}$, we see that $M_{f_A}(n; R)^{2k}$ is behaving like a sieve weight in this case.

If $A \leq \frac{1}{2k} \binom{2k}{k} - 1 = \mathcal{E}_k/2k$, then $\mathcal{E}_{k,A} \geq 0$, and so $\mathcal{M}_{f_A, 2k}(R) \gg 1$. In particular, $M_{f_A}(n; R)^{2k}$ no longer behaves like a sieve weight, and the main contribution is from numbers with several prime factors in $[1, R]$.

The following theorem illustrates further this distinction.

Theorem 1.6. *Let $x \geq R \geq 2$, $k \in \mathbb{N}$ and $A \in \mathbb{Z}_{\geq 0}$. Moreover, let $\Omega(n; R)$ denote the number of prime factors of n in $[1, R]$, counted with multiplicity.*

(a) *If $A > \frac{1}{2k} \binom{2k}{k} - 1$, then*

$$\sum_{\substack{n < x \\ \Omega(n; R) \geq C}} M_{f_A}(n; R)^{2k} \ll_{k,A} \frac{x}{C \log R}.$$

(b) *If $A \leq \frac{1}{2k} \binom{2k}{k} - 1$ and $\epsilon > 0$ is fixed, then there is a $\delta = \delta(\epsilon, k) > 0$ such that*

$$\sum_{\substack{n < x \\ |\Omega(n; R) / \log \log R - \binom{2k}{k}| \geq \epsilon}} M_{f_A}(n; R)^{2k} \ll_{k,A} x (\log R)^{\binom{2k}{k} - 2k(A+1) - \delta}.$$

In other words, if $A > \frac{1}{2k} \binom{2k}{k} - 1$, then the main contribution to the sum defining $\mathcal{M}_{f_A, 2k}(R)$ comes from integers with a bounded number of prime factors $\leq R$; whereas if $A \leq \frac{1}{2k} \binom{2k}{k} - 1$, then the main contribution to the sum comes from integers with $(\binom{2k}{k} + o(1)) \log \log R$ prime factors $\leq R$.

Analogous results hold with $\Omega(n; R)$ replaced by the function $\#\{p|n : p \leq R\}$. We note that typically one requires $x > R^{2k}$, as in Theorem 1.3, to estimate a $2k^{\text{th}}$ moment of a sum of divisors of size at most R , but the estimates of Theorem 1.6 hold in the much wider range $x \geq R$. We can show similar (but slightly weaker) results for general weights f :

Theorem 1.7. *Let $k \in \mathbb{N}$ and $f : \mathbb{R} \rightarrow \mathbb{R}$ be supported in $(-\infty, 1]$. Assume further that $f \in C^A(\mathbb{R})$ and that all functions $f, f', \dots, f^{(A)}$ are uniformly bounded for some integer $A \geq 2$, and fix some $\epsilon \in (0, 1)$.*

(a) *Assume that $A > \frac{1}{2k} \binom{2k}{k}$. For $x \geq R \geq 2$ and $C \geq 1$, we have that*

$$\sum_{\substack{n < x \\ \Omega(n; R) \geq C}} M_f(n; R)^{2k} \ll_{k,f} \frac{x}{C \log R}.$$

(b) *If $A \leq \frac{1}{2k} \binom{2k}{k}$ and $\epsilon > 0$ is fixed, then there is a $\delta = \delta(\epsilon, k) > 0$ such that*

$$\sum_{\substack{n < x \\ |\Omega(n; R) / \log \log R - \binom{2k}{k}| \geq \epsilon}} M_f(n; R)^{2k} \ll_{k,f,\epsilon} x (\log R)^{\binom{2k}{k} - 2kA - \delta} \quad (x \geq R \geq 2).$$

1.6. Outline of the paper. We start the paper in Section 2 with a discussion of the discrepancy between the exponent of $\log R$ in (1.15) and the exponent of m in Theorem 1.1, which is surprising at first sight.

Sections 3 and 4 study the analogies for permutations and polynomials over finite fields, respectively. These analogies are considerably easier to analyze than the integer case.

The main body of the paper is then dedicated to the study of moments of $M_{f_A}(n; R)$ over Sections 5 – 10. Specifically, in Section 5 we establish relation (1.5) for the size of the support of $M_{f_0}(n; R)$, and in Section 6 we study inversion formulas for our divisor sums $M_f(n; R)$ that will be essential when dealing with their moments. The proof of Theorem 1.3 is separated over three sections: in Section 7, we establish certain combinatorial inequalities that will be instrumental in understanding the leading term in the asymptotics for $\mathcal{M}_{f_A, 2k}(R)$. Then, in Section 8 we establish Theorem 1.3 by a multidimensional contour shifting argument, except for showing the positivity of the constants $c_{k,A}$ and c'_k . The latter will be accomplished with

a different argument in Section 9. Section 10 contains an analysis of the anatomy of the integers that give the main contribution to moments of $M_f(n; R)$. Specifically, we prove Theorems 1.4, 1.6 and 1.7 there.

Finally, in Sections 11 and 12 we study the moments of the sum weighted by Dirichlet characters, and establish Theorem 1.5, first for non-exceptional Dirichlet characters (where the proof is similar to Theorem 1.2), and then for exceptional Dirichlet characters (where the proof is similar to Theorem 1.3).

1.7. Notation. Given an integer $N \geq 1$, we set throughout the paper

$$\begin{aligned} [N] &:= \{1, 2, \dots, N\}, \\ \mathcal{S}^+(N) &:= \{\emptyset \neq I \subset [N] : |I| \text{ even}\}, \quad \mathcal{S}^-(N) := \{I \subset [N] : |I| \text{ odd}\}, \\ \mathcal{S}(N) &:= \{I \subset [N]\} \quad \text{and} \quad \mathcal{S}^*(N) := \mathcal{S}^+(N) \cup \mathcal{S}^-(N). \end{aligned}$$

Also, we recall that, given a integer $n \geq 1$, we write $P^+(n)$ and $P^-(n)$ for its largest and smallest prime divisors, respectively, with the convention that $P^+(1) = 1$ and $P^-(1) = \infty$.

Finally, given $2k$ variables $x_1, \dots, x_{2k}$ and $I \subset [2k]$, we will use the notation $x_I = \sum_{i \in I} x_i$.

2. THE DISCREPANCY BETWEEN INTEGERS AND POLYNOMIALS

The goal of this section is to analyze in detail why we have such a different behaviour when considering integers vs. polynomials or permutations

2.1. Integer setting. Assume that $k \geq 2$. We mimic the proof of Theorem 1.1. Recall the definition of $\tilde{f}_0$ in (1.4). Given square-free integers $d_1, \dots, d_{2k}$ and $I \subset \mathcal{S}^*(2k)$, we let D_I be the product of those primes p that divide each of the d_i 's with $i \in I$ but do not divide $\prod_{i \in [2k] \setminus I} d_i$. Then the integers D_I for $I \in \mathcal{S}^*(2k)$ are pairwise coprime and $d_i = \prod_{I: i \in I} D_I$ for each i , so that

$$\begin{aligned} \mathcal{M}_{\tilde{f}_0, 2k}(R) &= \sum_{R/2 < d_1, \dots, d_{2k} < R} \frac{\mu(d_1) \dots \mu(d_{2k})}{[d_1, \dots, d_{2k}]} \\ &= \sum_{\substack{D_I \text{ (I} \in \mathcal{S}^*(2k)) \\ R/2 < \prod_{I \ni i} D_I \leq R \text{ (1} \leq i \leq 2k)}}^b \left(\prod_{I \in \mathcal{S}^-(2k)} \frac{\mu(D_I)}{D_I} \right) \left(\prod_{I \in \mathcal{S}^+(2k)} \frac{1}{D_I} \right), \end{aligned}$$

where the notation $\sum^b$ means that the summation is running over squarefree and pairwise coprime integers D_I . Set $L = e^{(\log \log R)^3}$. The contribution of those tuples with $D_I > L$ for some I odd to $\mathcal{M}_{\tilde{f}_0, 2k}(R)$ can be seen to be $\ll 1/e^{c(\log \log R)^{3/2}}$ for some $c = c(k) > 0$, by the Prime Number Theorem. So assume that $D_I \leq L$ for all I odd. Then it is natural to write

$$\mathcal{M}_{\tilde{f}_0, 2k}(R) \sim \sum_{\substack{D_I \leq L \text{ (I} \in \mathcal{S}^*(2k)) \\ D = \prod_{I \in \mathcal{S}^-(2k)} D_I}}^b \frac{\mu(D)}{D} \cdot T_{2k}(R_1, \dots, R_{2k}; D),$$

where $R_i = R / \prod_{I \ni i, I \in \mathcal{S}^+(2k)} D_I$ and

$$T_{2k}(\mathbf{R}; a) = \sum_{\substack{(D_I, a) = 1 \text{ (I} \in \mathcal{S}^+(2k)) \\ R_i/2 < \prod_{I \in \mathcal{S}^+(2k): i \in I} D_I \leq R_i \text{ (1} \leq i \leq 2k)}}^b \prod_{I \in \mathcal{S}^+(2k)} \frac{1}{D_I}.$$

When $\log R_i = \log R + O(\log L) = \log R + O((\log \log R)^{3/2})$, as it is here, we should be expecting that $T_{2k}(\mathbf{R}; a)$ has an asymptotic formula of the form

$$T_{2k}(\mathbf{R}; a) = g(a)(\log R)^{2^{2k-1}-2k-1} + O\left((\log R)^{2^{2k-1}-2k-2}(\log \log R)^{O(1)}\right),$$

since we have $2^{2k-1} - 1$ variables on a logarithmic scale and $2k$ multiplicative constraints in dyadic intervals, where $g(a)$ is a multiplicative function with $g(p) = 1 + O(1/p)$. Since $\sum_{n=1}^{\infty} \mu(n)/n = 0$, we then find that the total contribution of the main terms to $\mathcal{M}_{\tilde{f}_0, 2k}(R)$ is

$$\sum_{\substack{D_I \leq L \\ D = \prod_{I \in \mathcal{S}^-(2k)} D_I}}^b \frac{\mu(D)f(D)}{D} \ll e^{-c'(\log \log R)^{3/2}}.$$

for some $c' = c'(k) > 0$, which is negligible. Consequently,

$$\mathcal{M}_{\tilde{f}_0, 2k}(R) \ll (\log \log R)^{O(1)} (\log R)^{2^{2k-1}-2k-2},$$

whereas the power of m in Theorem 1.1 is $2^{2k-1} - 2k - 1$. So our heuristics indicate we get more cancellation in the integer setting than we will obtain in the analogous polynomial question.

2.2. Polynomial analogy. The reader might be sceptical of the argument presented above, because a direct analogue exists for polynomials over finite fields too. Specifically, expanding the $2k$ -th power in $\text{Poly}_q(n, m; k)$, we find that

$$\text{Poly}_q(n, m; k) = \sum_{G_1, \dots, G_{2k}} \frac{\mu(G_1) \cdots \mu(G_{2k})}{q^{\deg([G_1, \dots, G_{2k}])}}$$

for $n \geq 2mk$. Given square-free, monic polynomials $G_1, \dots, G_{2k}$ over $\mathbb{F}_q[t]$ and $I \subset \mathcal{S}^*(2k)$, we let G_I be the product of those monic irreducibles P that divide each of the G_i 's with $i \in I$ but do not divide $\prod_{i \in [2k] \setminus I} G_i$. Then the polynomials G_I for $I \in \mathcal{S}^*(2k)$ are pairwise coprime and $G_i = \prod_{I: i \in I} G_I$ for each i , so that

$$\text{Poly}_q(n, m; k) = \sum_{\substack{G_I \text{ } (I \in \mathcal{S}^*(2k)) \\ \sum_{I \ni i} \deg(G_I) = m \text{ } (1 \leq i \leq 2k)}}^b \left(\prod_{I \in \mathcal{S}^-(2k)} \frac{\mu(G_I)}{q^{\deg(G_I)}} \right) \left(\prod_{I \in \mathcal{S}^+(2k)} \frac{1}{q^{\deg(G_I)}} \right).$$

where the notation $\sum^b$ means that the summation is running over squarefree and pairwise monic polynomials G_I . As in the integer case, the contribution to $\text{Poly}_q(n, m; k)$ of those tuples $(G_I)_{I \in \mathcal{S}^*(2k)}$ such that $\deg(G_I)$ is large for some $I \in \mathcal{S}^-(2k)$ is negligible, by the Prime Number Theorem over $\mathbb{F}_q[t]$. Hence, we may assume that $\deg(G_I) \leq \log m$ for all $I \in \mathcal{S}^-(2k)$. Then it is natural to write

$$\text{Poly}_q(n, m; k) = \sum_{\substack{\deg(G_I) \leq \log m \text{ } (I \in \mathcal{S}^-(2k)) \\ G = \prod_{I \in \mathcal{S}^-(2k)} G_I}}^b \frac{\mu(G)}{q^{\deg(G)}} \cdot \tilde{T}_{q, 2k}(m_1, \dots, m_{2k}; G),$$

where $m_i = m - \sum_{I \ni i, I \in \mathcal{S}^+(2k)} \deg(G_I)$ and

$$\tilde{T}_{q,2k}(\mathbf{m}; A) := \sum_{\substack{(G_I, A)=1 \ (I \in \mathcal{S}^+(2k)) \\ \sum_{I \in \mathcal{S}^+(2k): i \in I} \deg(G_I) = m_i \ (1 \leq i \leq 2k)}} \prod_{I \in \mathcal{S}^+(2k)} \frac{1}{q^{\deg(G_I)}}.$$

As before, when $\ell_i = m + O(\log m)$, as above, we should be expecting that $\tilde{T}_{q,2k}(\ell; A)$ has an asymptotic formula of the form

$$(2.1) \quad \tilde{T}_{q,2k}(\ell; A) = \tilde{g}(A) m^{2^{2k-1}-2k-1} + O\left(m^{2^{2k-1}-2k-2}(\log m)^{O(1)}\right),$$

where $\tilde{g}(A)$ is a multiplicative function with $\tilde{g}(P) = 1 + O(1/q^{\deg(P)})$ for irreducibles P .

The above argument suggests that we should have an asymptotic behaviour of $\text{Poly}_q(n, m; k)$ that is smaller than what Theorem 1.2 states, which is absurd. The problem is that if $\sum_{I \ni i} \deg(G_I) = m$ for all i , then we also have that

$$2km = \sum_{i=1}^{2k} \sum_{I: i \in I} \deg(G_I) = \sum_I |I| \deg(G_I).$$

Reducing this formula mod 2, we find that

$$(2.2) \quad \sum_{I \in \mathcal{S}^-(2k)} \deg(G_I) \equiv 0 \pmod{2},$$

a local constraint that is not present in the integer analogue. In particular, we see that (2.1) is true only when (2.2) is satisfied. We thus find that the main term for $\text{Poly}_q(n, m; k)$ equals

$$\begin{aligned} & m^{2^{2k-1}-2k-1} \cdot \sum_{\substack{\deg(G_I) \leq \log m \ (I \in \mathcal{S}^-(2k)) \\ G = \prod_{I \in \mathcal{S}^-(2k)} G_I, \ 2 \mid \deg(G)}} \frac{\mu(G) \tilde{g}(G)}{q^{\deg(G)}} \\ &= \frac{m^{2^{2k-1}-2k-1}}{2} \cdot \sum_{\substack{\deg(G_I) \leq \log m \ (I \in \mathcal{S}^-(2k)) \\ G = \prod_{I \in \mathcal{S}^-(2k)} G_I}} \frac{\mu(G)(1 + (-1)^{\deg(G)})}{q^{\deg(G)}} \\ &\asymp m^{2^{2k-1}-2k-1}, \end{aligned}$$

because $\sum_F \mu(F)/q^{\deg(F)} = 0$ and $\sum_F \mu(F)(-1/q)^{\deg(F)} = \prod_P (1 - (-1/q)^{\deg(P)}) > 0$. Thus we see the local constraint associated to the discreteness of degrees in the polynomial setting means we have genuinely different asymptotic behavior.

2.3. Further analysis. The above arguments suggest a possible route to proving Theorem 1.3, by working out the full asymptotic expansion of $N_{2k}(\mathbf{x}; a)$. Controlling the coefficients in this expansion is a highly non-trivial problem. Instead, we take another route, using a high-dimensional contour shifting argument. Our starting point is Perron's inversion formula which, ignoring convergence issues, yields

$$T_{2k}(\mathbf{x}; a) \sim \frac{1}{(2\pi i)^{2k}} \int \cdots \int_{\substack{\text{Re}(s_j)=1/\log R \\ 1 \leq j \leq 2k}} \sum_{\substack{(D_I, a)=1 \\ I \in \mathcal{S}^+(2k)}} \prod_{I \in \mathcal{S}^+(2k)} \frac{1}{D_I^{1+s_I}} \prod_{j=1}^{2k} \frac{x_j^{s_j} (1 - 2^{-s_j})}{s_j} ds_1 \cdots ds_{2k},$$

with the notational convention that $s_I = \sum_{j \in I} s_j$. Therefore

$$\mathcal{M}_{\tilde{f}_0, 2k}(R) \sim \frac{1}{(2\pi i)^{2k}} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=1/\log R \\ 1 \leq j \leq 2k}} F(\mathbf{s}) \frac{\prod_{I \in \mathcal{S}^+(2k)} \zeta(1+s_I)}{\prod_{I \in \mathcal{S}^-(2k)} \zeta(1+s_I)} \prod_{j=1}^{2k} \frac{1-2^{-s_j}}{s_j} ds_1 \cdots ds_{2k},$$

where $F(\mathbf{s})$ is analytic and non-zero when $\operatorname{Re}(s_j) > -1/4k$ for all j . As we will see in Section 8, shifting contours, we pick up poles any time $s_I = 0$ for some $I \in \mathcal{S}^+(2k)$. What is the difficulty in proving Theorem 1.3 is that some of these poles can get annihilated by poles of the zeta factors in the denominator, which is an analytic way of saying that the higher order terms in the asymptotic expansions of $T_{2k}(\mathbf{x}; a)$ are cancelled out.

It is clear from the above discussion that the underlying reason why we got a genuinely smaller main term for $\mathcal{M}_{\tilde{f}_0, 2k}(R)$ is the identity $\sum_{n=1}^{\infty} \mu(n)/n = 0$, that is to say the fact that $1/\zeta$ has a zero at 1. This also explains the phenomenon we see in Theorem 1.5. If we replace μ by a real valued multiplicative function f whose Dirichlet series $F(s) = \sum_{n=1}^{\infty} f(n)/n^s$ which is not very small at $s = 1$, then the behaviour of the respective divisor sums should be similar as the permutation analogue, whilst if $F(s)$ is close to 0 at $s = 1$ (which occurs if F has a zero very close to 1) the behaviour is the same as in the original integer setting.

2.4. Further obstructions to the analogy. Is it possible that the local constraints at the prime 2 described above are the only thing separating integers and polynomials? In order to study this question, we consider the variations

$$\operatorname{Poly}_q(n, m, h; k) := \frac{1}{q^n} \sum_{\substack{N \in \mathbb{F}_q[t] \\ \deg N = n}} \left| \sum_{\substack{M|N \\ m-h < \deg M \leq m}} \mu(M) \right|^{2k},$$

where $h \in \mathbb{Z} \cap [1, m+1]$. If $h \geq 2$, then the local problems at the factor should be resolved. However, we will see that this is not sufficient, and that the discrepancy between the integer and the polynomial analogues goes even deeper.

First, let us consider the case $h = m+1$ in order to convince the reader that resolving the constraints at the prime 2 is not sufficient. It is known that a positive proportion of polynomials $N \in \mathbb{F}_q[t]$ has a simple zero over $\mathbb{F}_q$, and the number of these zeroes is, on average, bounded. So it should be

$$\operatorname{Poly}_q(n, m, m+1; k) \asymp \frac{1}{q^n} \sum_{\substack{N \in \mathbb{F}_q[t] \\ \deg N = n}} \sum_{\substack{\alpha \in \mathbb{F}_q \\ N(\alpha)=0 \\ N'(\alpha) \neq 0}} \left| \sum_{\substack{M|N \\ \deg M \leq m}} \mu(M) \right|^{2k}.$$

If N has a simple zero at α , then we can factor $N(x) = (x - \alpha)\tilde{N}(x)$, where $\tilde{N}(\alpha) \neq 0$, that is to say $x - \alpha$ and $\tilde{N}$ are co-prime. Then

$$\sum_{\substack{M|N \\ \deg M \leq m}} \mu(M) = \sum_{\substack{M|\tilde{N} \\ \deg M \leq m}} \mu(M) + \sum_{\substack{M|\tilde{N} \\ 1 + \deg M \leq m}} \mu((x - \alpha) \cdot M) = \sum_{\substack{M|\tilde{N} \\ \deg M = m}} \mu(M),$$

so that

$$\text{Poly}_q(n, m, m+1; k) \asymp \frac{1}{q^n} \sum_{\alpha \in \mathbb{F}_q} \sum_{\substack{\tilde{N} \in \mathbb{F}_q[t] \\ \deg \tilde{N} = n-1 \\ \tilde{N}(\alpha) \neq 0}} \left| \sum_{\substack{M | \tilde{N} \\ \deg M = m}} \mu(M) \right|^{2k} \asymp m^{2^{2k-1}-2k-1} + 1$$

for $n \geq 2mk$, by an easy variation of Theorem 1.2. This argument can be made rigorous; we leave this task to the interested reader.

Let us now study $\text{Poly}_q(n, m, h; k)$ more generally. For any $h \in \mathbb{N}$ and $n \geq 2mk$, we note that

$$\begin{aligned} \text{Poly}_q(n, m, h; k) &= \sum_{\substack{G_1, \dots, G_{2k} \\ m-h < \deg(G_i) \leq m \\ 1 \leq i \leq 2k}} \frac{\mu(G_1) \cdots \mu(G_{2k})}{q^{\deg([G_1, \dots, G_{2k}])}} \\ &= \sum_{\substack{G_I \ (I \in \mathcal{S}^*(2k)) \\ m-h < \sum_{I \ni i} \deg(G_I) \leq m \ (1 \leq i \leq 2k)}}^b \left(\prod_{I \in \mathcal{S}^-(2k)} \frac{\mu(G_I)}{q^{\deg(G_I)}} \right) \left(\prod_{I \in \mathcal{S}^+(2k)} \frac{\mu^2(G_I)}{q^{\deg(G_I)}} \right), \end{aligned}$$

as before. Applying Fourier inversion $2k$ times, we find that, for any $r \in (0, 1)$,

$$\begin{aligned} \text{Poly}_q(n, m, h; k) &= \sum_{\substack{m-h < \ell_j \leq m \\ 1 \leq j \leq 2k}} \sum_{G_I \ (I \in \mathcal{S}^*(2k))}^b \left(\prod_{I \in \mathcal{S}^-(2k)} \frac{\mu(G_I)}{q^{\deg(G_I)}} \right) \left(\prod_{I \in \mathcal{S}^+(2k)} \frac{\mu^2(G_I)}{q^{\deg(G_I)}} \right) \\ &\quad \times \prod_{j=1}^{2k} \int_0^1 (re(\theta_j))^{-\ell_j + \sum_{I \ni j} \deg(G_I)} d\theta_j. \end{aligned}$$

So, if we set

$$\mathcal{Z}_q(w) = \sum_{G \in \mathbb{F}_q[t]} \left(\frac{w}{q} \right)^{\deg(G)} = \prod_P (1 - (w/q)^{\deg(P)})^{-1},$$

the $\mathbb{F}_q[t]$ analogue of the Riemann zeta function, then

$$\begin{aligned} \text{Poly}_q(n, m, h; k) &= \sum_{\substack{m-h < \ell_j \leq m \\ 1 \leq j \leq 2k}} \int_{[0,1]^{2k}} \tilde{F}_q((re(\theta_j))_j) \frac{\prod_{I \in \mathcal{S}^+(2k)} \mathcal{Z}_q(r^{|I|}e(\theta_I))}{\prod_{I \in \mathcal{S}^-(2k)} \mathcal{Z}_q(r^{|I|}e(\theta_I))} \prod_{j=1}^{2k} \frac{e(-\ell_j \theta_j)}{r^{\ell_j}} d\theta \\ &= \int_{[0,1]^{2k}} \tilde{F}_q((re(\theta_j))_j) \frac{\prod_{I \in \mathcal{S}^+(2k)} \mathcal{Z}_q(r^{|I|}e(\theta_I))}{\prod_{I \in \mathcal{S}^-(2k)} \mathcal{Z}_q(r^{|I|}e(\theta_I))} \prod_{j=1}^{2k} \sum_{\ell=m-h+1}^m \frac{e(-\ell \theta_j)}{r^\ell} d\theta, \end{aligned}$$

where $\theta_I = \sum_{j \in I} \theta_j$ and $\tilde{F}_q(\mathbf{w})$ is a certain function that is analytic and non-zero when $|w_j| < \sqrt{q}/2k$ for all j .

We take $r = 1 - 1/m$ and note that the main contribution to $\text{Poly}_q(n, m, h; k)$ should come from those values of θ_I for which there are many $I \in \mathcal{S}^+(2k)$ such that $\theta_I \equiv 0 \pmod{1}$. This is the key difference with the integer case: before, we needed many $I \in \mathcal{S}^+(2k)$ with $s_I = 0$. So we see two different linear algebra problems: one over the group $\mathbb{R}/\mathbb{Z}$, which has

torsion, and one over $\mathbb{R}$, which is a field. The presence of torsion in $\mathbb{R}/\mathbb{Z}$ is a reflection of the discreteness of the polynomial setting (of the degree of the polynomials, more precisely), and the fact that $\mathbb{R}$ is a field reflects the continuous nature of the integer problem (of the logarithms of integers, more precisely).

When $h = 1$, then the integrand is $\asymp 1/m$ when $\theta_j = O(1/m) \bmod 1$ for all j , much like the integer analogue. However, if we take $\theta_j = 1/2 + O(1/m)$ for all j , then we see that $\theta_I = O(1/m) \bmod 1$ for $I \in \mathcal{S}^+(2k)$, whereas $\theta_I = 1/2 + O(1/m) \bmod 1$ for $I \in \mathcal{S}^-(2k)$, so the integrand has size $m^{2^{2k-1}-1}$ for such θ . The volume of this region is $\asymp 1/m^{2k}$, leading to a contribution of size $m^{2^{2k-1}-2k-1}$ to $\text{Poly}_q(n, m, 1; k)$, which is precisely its order of magnitude for $k \geq 2$. Note that the fact the maximum occurs when $\theta_j \sim 1/2$ and not when $\theta_j \sim 0$ is a reflexion of the local constraint at the prime 2 we noticed above.

Similarly to the above case, if $h = 2$ and $\theta_j = 1/2 + O(1/m)$, then the integrand becomes

$$\tilde{F}_q(1/2, \dots, 1/2) \frac{m^{2^{2k-1}-1}(1 + O(1/m))}{\mathcal{Z}_q(1/2)^{4k}} \prod_{j=1}^{2k} (1 + e(\theta_j)).$$

By Taylor expansion, we have that

$$1 + e(\theta_j) = 1 - e(\theta_j - 1/2) = -(\theta_j - 1/2) - \frac{(\theta_j - 1/2)^2}{2} - \dots$$

By symmetry, we should then have that

$$\begin{aligned} & \int \cdots \int_{\substack{|\theta_j - 1/2| \leq 1/m \\ 1 \leq j \leq 2k}} \tilde{F}_q(1/2, \dots, 1/2) \frac{m^{2^{2k-1}-1}(1 + O(1/m))}{\mathcal{Z}_q(1/2)^{4k}} \prod_{j=1}^{2k} (1 + e(\theta_j)) d\theta \\ &= (1 + O(1/m)) \int \cdots \int_{\substack{|\theta_j - 1/2| \leq 1/m \\ 1 \leq j \leq 2k}} \tilde{F}_q(1/2, \dots, 1/2) \frac{m^{2^{2k-1}-1}}{\mathcal{Z}_q(1/2)^{4k}} \prod_{j=1}^{2k} \frac{(\theta_j - 1/2)^2}{2} d\theta, \end{aligned}$$

which leads to a contribution of size $m^{2^{2k-1}-6k-1}$ to $\text{Poly}_q(n, m, 2; k)$. This should be the dominant contribution for large k , even though for small k other regions can dominate. For example, if $k = 2$ and we take $\theta_1, \theta_2 \in [0.33, 0.34]$, $\theta_3, \theta_4 \in [0.66, 0.67]$, and $\theta_1 + \theta_2, \theta_3 + \theta_4, \theta_1 + \theta_4 = O(1/m)$, then the integrand becomes $\asymp m^5$, and we are integrating over a region of volume $\asymp 1/m^3$, so we see that $\text{Poly}_q(n, m, 1; 2) \gg m^2$. In fact, this is the exact order of magnitude of $\text{Poly}(n, m, 1; 2)$.

We conclude our discussion with another peculiar fact: if $h = 3$, then

$$\sum_{\ell=m-h+1}^m e(\ell\theta) = e(m\theta)(1 + e(\theta) + e(2\theta)) = e(m\theta) + O(1/m)$$

when $\theta = 1/2 + O(1/m)$. So $\text{Poly}_q(n, m, 3; k)$ should have the same size as $\text{Poly}_q(n, m, 1; k)$, whereas $\text{Poly}_q(n, m, h; k)$ is a bit smaller, by a factor of size $m^{O(k)}$. In general, no matter how we choose h , we cannot make the sum $\sum_{\ell=m-h+1}^m e(\ell\theta)$ small enough to cancel the contribution of the factors $\mathcal{Z}_q(r^{|I|}e(\theta_I))$ for even I in the region $\theta_j \sim 1/2$, so the quantities $\text{Poly}_q(n, m, h; k)$ do not behave in the same way as $\mathcal{M}_{\tilde{f}_0, 2k}(R)$ for k large.

3. THE ANALOGY FOR PERMUTATIONS

Completion of the proof of Theorem 1.1. It remains to prove the two claims for $c(m, k)$. Given any vector $\{r_I : \emptyset \neq I \subset [2k]\}$ counted by $c(m, k)$, the vector $\{r'_I : \emptyset \neq I \subset [2k]\}$ is counted by $c(m+1, k)$ where $r'_{\{1,2\}} = r_{\{1,2\}} + 1$ and $r'_{\{3,4,\dots,2k\}} = r_{\{3,4,\dots,2k\}} + 1$, and $r'_I = r_I$ otherwise. Hence $c(m, k) \leq c(m+1, k)$ for all $m \geq 0$, as claimed.

We now estimate $c(m, k)$. When $k = 1$, we find immediately that $c(m, 1) = 2$, so there is nothing to prove. Assume now that $k \geq 2$. Since $c(m, k)$ is increasing in m and $c(0, k) = 1$, we may assume that m is even and large enough. We note that there are $\asymp_k 1$ possibilities for the r_I for the odd-sized I . Otherwise we have to satisfy $2k$ equations with $2^{2k-1} - 1$ variables. Hence the number of solutions should be

$$\asymp_k m^{2^{2k-1}-2k-1} + 1,$$

as claimed. Certainly, this argument yields an appropriate upper bound. To prove the lower bound for $k \geq 2$ we will construct this number of solutions. Let

$$\mathcal{I} = \{\{i, j\} : 1 \leq i < j \leq 4\} \cup \{\{1, j\} : 5 \leq j \leq 2k\},$$

so that $|\mathcal{I}| = 2k + 2$. Set $r_I = 0$ if $I \in \mathcal{S}^-(2k)$ and, given $\delta > 0$ to be chosen later, let r_I be any even integer from the range $[0, \delta m/4^k]$ if $I \in \mathcal{S}^+(2k) \setminus (\mathcal{I} \cup \{5, 6, \dots, 2k\})$. Finally, if $k > 2$, let $r_{\{5,6,\dots,2k\}}$ be an even integer from the range $[m - 2\delta m, m - \delta m]$. There are $\asymp_{k,\delta} m^{2^{2k-1}-3-2k}$ such choices of r_I , $I \in \mathcal{S}^+(2k) \setminus \mathcal{I}$. Then select

$$r_{\{1,j\}} := m - \sum_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}, j \in I} r_I \quad (5 \leq j \leq 2k),$$

which is an even integer lying in the interval $[0, \delta m]$, and set

$$m_j = m - \sum_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}, j \in I} r_I \quad (1 \leq j \leq 4),$$

which are even integers lying in the interval $[m - 2k\delta m, m]$. It remains to choose r_I , $I \in \mathcal{I}$, such that $\sum_{I \in \mathcal{I}, j \in I} r_I = m_j$. Then, we select any even integers $r_{\{2,4\}}, r_{\{3,4\}}$ from $[\sqrt{\delta}m - \delta m, \sqrt{\delta}m + \delta m]$, and we set

$$r_{\{1,4\}} = m_4 - r_{\{2,4\}} - r_{\{3,4\}}.$$

Finally, we define $r_{\{1,2\}}, r_{\{1,3\}}, r_{\{2,3\}}$ such that

$$\begin{aligned} r_{\{1,2\}} + r_{\{1,3\}} &= m_1 - r_{\{1,4\}} = m_1 - m_4 + r_{\{2,4\}} + r_{\{3,4\}}; \\ r_{\{1,2\}} + r_{\{2,3\}} &= m_2 - r_{\{2,4\}}; \text{ and} \\ r_{\{1,3\}} + r_{\{2,3\}} &= m_3 - r_{\{3,4\}}. \end{aligned}$$

Note that the right-hand sides are all even so there is no parity problem, and the solutions we obtain are non-negative integers for δ small enough. We have thus constructed $\gg_k m^{2^{2k-1}-2k-1}$ solutions counted by $c(m, k)$. This completes the proof of the lemma. $\square$

Remark 3.1. It should not be too difficult to determine $c(m, k)$ exactly in some special cases. For example, we have that $c(m, 1) = 2$ and $c(m, 2) = \frac{1}{3}(64m^3 - 135m^2 + 182m - 66)$ for all $m \geq 1$.

Finally, we prove a probabilistic interpretation for $c(m, k)$. In its statement, we have set with a slight abuse of notation

$$(3.1) \quad M(\mathbf{c}; m) := \sum_{\substack{0 \leq b_j \leq c_j \\ 1 \leq j \leq m \\ \sum_j j b_j = m}} (-1)^{b_1 + \dots + b_m}$$

for an m -tuple of non-negative integers $\mathbf{c} = (c_1, \dots, c_m)$.

Proposition 3.1. *Let $\mathbf{X} = (X_1, X_2, \dots, X_m)$ be a vector of pairwise independent Poisson random variables, where X_j has parameter $1/j$. For every $k \in \mathbb{N}$, we have that*

$$c(m, k) = \mathbb{E}[M(\mathbf{X}; m)^{2k}].$$

Before we prove Proposition 3.1, we need a lemma.

Lemma 3.2. *Let $n \geq m \geq 1$. The proportion of permutations $\sigma \in S_n$ that have no cycles of length $\leq m$ is*

$$\prod_{j=1}^m e^{-1/j} + O\left(\frac{m^2}{n}\right).$$

Proof. Note that this lemma was proven in [9] for large m , but here we are mainly interested in the case when m is very small compared to n . We apply inclusion-inclusion. If $\mathcal{C}_j$ denotes the j -cycles in S_n , we write $|\pi| = j$ for an element π of $\mathcal{C}_j$, and we let $\mathcal{C}$ be the union of $\mathcal{C}_1, \dots, \mathcal{C}_m$. Then

$$\begin{aligned} & \#\{\sigma \in S_n : \sigma \text{ has no cycles of length } \leq m\} \\ &= n! - \sum_{\pi \in \mathcal{C}} (n - |\pi|)! + \sum_{\substack{\pi_1, \pi_2 \in \mathcal{C} \\ \pi_1, \pi_2 \text{ disjoint}}} (n - |\pi_1| - |\pi_2|)! \mp \dots \\ &= \sum_{\substack{c_1, \dots, c_m \geq 0 \\ c_1 + 2c_2 + \dots + mc_m \leq n}} (-1)^{c_1 + \dots + c_m} (n - c_1 - 2c_2 - \dots - mc_m)! \sum_{\substack{\pi_1, \pi_2, \dots \in \mathcal{C} \text{ disjoint} \\ \#\{i: |\pi_i| = j\} = c_j \ \forall j}} 1. \end{aligned}$$

In order to count the inner quantity, we note that if $r = c_1 + \dots + c_m$ is the total number of disjoint cycles we are choosing, and we have fixed our choice for $\pi_1, \pi_2, \dots, \pi_{r-1}$, then there are

$$\frac{(n - |\pi_1| - \dots - |\pi_{r-1}|)!}{|\pi_r|! (n - |\pi_1| - \dots - |\pi_{r-1}| - |\pi_r|)!}$$

choices for the set π_r will permute, and then $(|\pi_r| - 1)!$ possibilities for a cycle on $|\pi_r|$ given elements. Inductively, we then find that the total number of possibilities for $\pi_1, \dots, \pi_r$ should be

$$\frac{n!}{(n - |\pi_1| - \dots - |\pi_r|)!} \cdot \frac{1}{|\pi_1| \dots |\pi_r|} = \frac{n!}{(n - c_1 - 2c_2 - \dots - mc_m)!} \prod_{j=1}^m \frac{1}{j^{c_j}}.$$

Note though we have overcounted: each possibility of j -cycles occurs $c_j!$ times, depending on the order they are picked, so we must divide the above expression by $c_1! \dots c_m!$. We then

find that

$$\begin{aligned} \frac{\#\{\sigma \in S_n : \sigma \text{ has no cycles of length } \leq m\}}{n!} &= \sum_{\substack{c_1, \dots, c_m \geq 0 \\ c_1 + 2c_2 + \dots + mc_m \leq n}} \prod_{j=1}^m \frac{(-1/j)^{c_j}}{c_j!} \\ &= \prod_{j=1}^m e^{-1/j} + O\left(\frac{m^2}{n}\right), \end{aligned}$$

where the error term is obtained by noting that $\mathbf{1}_{c_1 + 2c_2 + \dots + mc_m \leq n} \leq (c_1 + 2c_2 + \dots + mc_m)/n$. $\square$

Proof of Proposition 3.1. We recall that we have already proved that

$$c(m, k) = \frac{1}{n!} \sum_{\sigma \in S_n} \left(\sum_{\substack{T \subset [n] \\ \sigma(T)=T \\ |T|=m}} \mu(\sigma|_T) \right)^{2k}$$

for any $n \geq 2mk$. We will now rewrite the right hand side for n much larger than m and k . Note that if σ has c_j cycles of length j for each $j \in \{1, \dots, m\}$, then

$$\sum_{\substack{T \subset [n] \\ \sigma(T)=T \\ |T|=m}} \mu(\sigma|_T) = M(\mathbf{c}; m) = \sum_{\substack{0 \leq b_j \leq c_j \\ 1 \leq j \leq m \\ \sum_j j b_j = m}} (-1)^{b_1 + \dots + b_m}$$

where $\mathbf{c} = (c_1, \dots, c_m)$. Moreover, a generalization of Cauchy's formula (see Lemma 2.2 in [5]) implies that if $t := c_1 + 2c_2 + \dots + mc_m \leq n$, then

$$\begin{aligned} &\frac{\#\{\sigma \in S_n : \sigma \text{ has } c_j \text{ } j\text{-cycles of length } j \text{ } (1 \leq j \leq m)\}}{n!} \\ &= \left(\prod_{j=1}^m \frac{1}{j^{c_j} c_j!} \right) \cdot \frac{\#\{\sigma \in S_{n-t} : \sigma \text{ has no cycles of length } \leq m\}}{(n-t)!}. \end{aligned}$$

Applying Lemma 3.2, it is then easy to conclude that

$$\begin{aligned} c(m, k) &= \lim_{n \rightarrow \infty} \frac{1}{n!} \sum_{\sigma \in S_n} \left(\sum_{\substack{T \subset [n] \\ \sigma(T)=T \\ |T|=m}} \mu(\sigma|_T) \right)^{2k} \\ &= \sum_{c_1, \dots, c_m \geq 0} M(\mathbf{c}; m)^{2k} \prod_{j=1}^m \frac{e^{-1/j}}{j^{c_j} c_j!}, \end{aligned}$$

which completes the proof. $\square$

4. THE ANALOGY FOR POLYNOMIALS OVER FINITE FIELDS

Proof of Theorem 1.2. Throughout this proof all polynomials we consider are monic, and P denotes a generic monic irreducible polynomial over $\mathbb{F}_q$. Note that

$$\begin{aligned} \text{Poly}_q(n, m; k) &= \frac{1}{q^n} \sum_{\deg(F)=n} \left(\sum_{\substack{G|F \\ \deg(G)=m}} \mu(G) \right)^{2k} \\ &= \prod_{\deg(P) \leq m} (1 - q^{-\deg(P)}) \sum_{\substack{P|F \Rightarrow \deg(P) \leq m}} \frac{1}{q^{\deg(F)}} \left(\sum_{\substack{G|F \\ \deg(G)=m}} \mu(G) \right)^{2k} \end{aligned}$$

for $n \geq 2km$, as it can be proven by expanding the $2k$ -th power in both sides, and noticing that if $G_j|F$ for each $j \leq 2k$, then we may write $F = [G_1, \dots, G_{2k}]H$ for some monic polynomial H .

Next, note that if $F = P_1^{n_1} \dots P_r^{n_r}$ is the factorisation of F into monic irreducible factors, and we write $c_j = \#\{i : \deg(P_i) = j\}$ for $1 \leq j \leq m$, then

$$\sum_{\substack{G|F \\ \deg(G)=m}} \mu(G) = M(\mathbf{c}; m)$$

with $M(\mathbf{c}; m)$ defined by (3.1). In particular, we see that $\sum_{G|F, \deg(G)=m} \mu(G)$ is a function of the vector $\mathbf{c} = (c_1, \dots, c_m)$, which we call the *factorisation type* of F . Moreover,

$$\begin{aligned} \prod_{\deg(P) \leq m} (1 - q^{-\deg(P)}) \sum_{\substack{P|F \Rightarrow \deg(P) \leq m \\ F \text{ has factorisation type } \mathbf{c}}} \frac{1}{q^{\deg(F)}} &= \prod_{\deg(P) \leq m} (1 - q^{-\deg(P)}) \prod_{j=1}^m \frac{\binom{N_j}{c_j}}{(q^j - 1)^{c_j}} \\ &= \prod_{j=1}^m \frac{\binom{N_j}{c_j} (1 - q^{-j})^{N_j}}{(q^j - 1)^{c_j}}, \end{aligned}$$

where

$$N_j := \#\{P \in \mathbb{F}_q[t] : P \text{ irreducible, } \deg(P) = j\}.$$

(Note that we have $(q^j - 1)^{c_j}$ and not q^{jc_j} in the denominator because we have to sum over powers of P_j too.) Galois theory implies that $q^j = \sum_{j'|j} j' N_{j'}$, whence

$$(4.1) \quad N_j = \frac{1}{j} \sum_{j'|j} \mu(j') q^{j/j'} = \frac{q^j}{j} \left(1 + O\left(\frac{\mathbf{1}_{j \geq 2}}{(q^j/j)^{1/2}}\right) \right)$$

and

$$(4.2) \quad q + j N_j \leq q^j \quad (j \geq 2).$$

Our next task is to control the quantity

$$\prod_{j=1}^m \frac{\binom{N_j}{c_j} (1 - q^{-j})^{N_j}}{(q^j - 1)^{c_j}}$$

and remove the dependence on q . First, note that

$$\prod_{j=1}^m (1 - q^{-j})^{N_j} = (1 + O(1/q)) \prod_{j=1}^m e^{-1/j}.$$

Furthermore,

$$\frac{\binom{N_j}{c_j}}{(q^j - 1)^{c_j}} = \frac{N_j^{c_j} (1 + O(c_j/N_j))^{c_j}}{c_j! (q^j - 1)^{c_j}} = \frac{1}{c_j! j^{c_j}} \left(1 + O\left(\frac{\mathbf{1}_{j \geq 2} \cdot c_j}{(q^j/j)^{1/2}} + \frac{\mathbf{1}_{j=1} c_j}{q^j} \right) \right),$$

provided that $c_1 \leq q$ and that $c_j \leq \sqrt{q^j/j}$ if $j \geq 2$. Therefore, if $c_1 \leq q$ and $\sum_{2 \leq j \leq m} c_j \cdot (j/q^j)^{1/2} \leq 1$, then

$$\prod_{j=1}^m \frac{\binom{N_j}{c_j} (1 - q^{-j})^{N_j}}{(q^j - 1)^{c_j}} = \left(1 + O\left(\frac{c_1 + 1}{q} + \sum_{j=2}^m \frac{c_j j^{1/2}}{q^{j/2}} \right) \right) \prod_{j=1}^m \frac{e^{-1/j}}{c_j! j^{c_j}}.$$

Together with Proposition 3.1, this implies that

$$\frac{1}{q^n} \sum_{\deg(F)=n} \left(\sum_{\substack{G|F \\ \deg(G)=m}} \mu(G) \right)^{2k} = c(m, k) + O(R_1 + R_2 + R_3) \quad (n \geq 2mk),$$

where

$$\begin{aligned} R_1 &= \sum_{c_1, \dots, c_m \geq 0} M(\mathbf{c}; m)^{2k} \left(\frac{c_1 + 1}{q} + \sum_{j=2}^m \frac{c_j j^{1/2}}{q^{j/2}} \right) \prod_{j=1}^m \frac{e^{-1/j}}{c_j! j^{c_j}}, \\ R_2 &= \sum_{\substack{c_1, \dots, c_m \geq 0 \\ c_1 > q \text{ or } \sum_{j>1} c_j j^{1/2}/q^{j/2} > 1}} M(\mathbf{c}; m)^{2k} \prod_{j=1}^m \frac{e^{-1/j}}{c_j! j^{c_j}} \\ &\leq \sum_{c_1, \dots, c_m \geq 0} M(\mathbf{c}; m)^{2k} \left(\frac{c_1}{q} + \sum_{j=2}^m \frac{c_j j^{1/2}}{q^{j/2}} \right) \prod_{j=1}^m \frac{e^{-1/j}}{c_j! j^{c_j}} \leq R_1, \end{aligned}$$

and

$$R_3 = \sum_{\substack{c_1, \dots, c_m \geq 0 \\ c_1 > q \text{ or } \sum_{j>1} c_j j^{1/2}/q^{j/2} > 1}} M(\mathbf{c}; m)^{2k} \prod_{j=1}^m \frac{e^{-1/j} \binom{N_j}{c_j}}{(q^j - 1)^{c_j}}.$$

For R_3 , we note that $c_j \leq N_j$ in its range; otherwise, $\binom{N_j}{c_j} = 0$. In particular, $c_1 \leq N_1 = q$. Moreover, (4.2) implies that

$$\binom{N_j}{c_j} \leq \frac{N_j^{c_j}}{c_j!} \leq \begin{cases} \frac{(q^j - 1)^{c_j}}{c_j! j^{c_j}} & \text{if } j \geq 2, \\ \frac{q^{c_1}}{c_1!} \leq (1 - 1/q)^{-q} \cdot \frac{(q - 1)^{c_1}}{c_1!} & \text{if } j = 1, \end{cases}$$

Therefore

$$R_3 \ll \sum_{c_1, \dots, c_m \geq 0} M(\mathbf{c}; m)^{2k} \left(\sum_{j=2}^m \frac{c_j j^{1/2}}{q^{j/2}} \right) \prod_{j=1}^m \frac{e^{-1/j}}{c_j! j^{c_j}} \leq R_1.$$

We thus see that Theorem 1.2 is reduced to proving that $R_1 \ll_k c(m, k)/q$. It suffices to show that

$$T_i := \sum_{c_1, \dots, c_m \geq 0} c_i M(\mathbf{c}; m)^{2k} \prod_{j=1}^m \frac{e^{-1/j}}{j^{c_j} c_j!} \ll c(m, k) \quad (1 \leq i \leq m).$$

Indeed, we note that the term with $c_i = 0$ does not contribute, and we replace c_i by $c_i + 1$ to find that

$$T_i = \frac{1}{i} \sum_{c_1, \dots, c_m \geq 0} M(\mathbf{e}_i + \mathbf{c}; m)^{2k} \prod_{j=1}^m \frac{e^{-1/j}}{j^{c_j} c_j!},$$

where $\mathbf{e}_i$ denotes the m -th dimensional vector all of whose coordinates are 0 except for the i -th coordinate that equals 1. Note that

$$M(\mathbf{e}_i + \mathbf{c}; m) = \sum_{\substack{0 \leq b_j \leq c_j \quad \forall j \neq i \\ 0 \leq b_i \leq c_i + 1 \\ \sum_j j b_j = m}} (-1)^{b_1 + \dots + b_m} = M(\mathbf{c}; m) + (-1)^{c_i + 1} M(\mathbf{c}_i; m - i(c_i + 1)),$$

where $\mathbf{c}_i = (c_1, \dots, c_{i-1}, 0, c_{i+1}, \dots, c_m)$, so that

$$M(\mathbf{e}_i + \mathbf{c}; m)^{2k} \leq 2^{2k-1} (M(\mathbf{c}; m) + M(\mathbf{c}_i; m - i(c_i + 1)))^{2k},$$

by Hölder's inequality. We thus conclude that

$$\begin{aligned} T_i &\leq \frac{2^{2k-1}}{i} c(m, k) + \frac{2^{2k-1}}{i} \sum_{c_i=0}^{\infty} \frac{e^{-1/i}}{c_i! i^{c_i}} \sum_{(c_j)_{j \leq m, j \neq i}} M(\mathbf{c}_i; m - i(c_i + 1))^{2k} \prod_{j \neq i} \frac{e^{-1/j}}{c_j! j^{c_j}} \\ &\leq \frac{2^{2k-1}}{i} c(m, k) + \frac{2^{2k-1}}{i} \sum_{c_i=0}^{\infty} \frac{e^{-1/i}}{c_i! i^{c_i}} c(m - i(c_i + 1), k), \end{aligned}$$

since the $M(\mathbf{c}_i; m - i(c_i + 1))^{2k}$ is independent of the value of the c_j 's with $j > m - i(c_i + 1)$. Recalling that $c(\ell, k)$ is an increasing function of ℓ by Theorem 1.1, we arrive to the claimed bound $T_i \ll_k c(m, k)$, whence Theorem 1.2 follows. $\square$

5. THE SUPPORT OF $M_{f_0}(n; R)$

We prove here (1.5). The lower bound was proven in the introduction. For the upper bound, let q be the smallest prime dividing n and write $n = q^j m$ with $q \nmid m$. Relation (1.6) implies that

$$M_{f_0}(n; R) = \sum_{\substack{d|m \\ R/q < d \leq R}} \mu(d).$$

Therefore,

$$(5.1) \quad \#\{n \leq x : M_{f_0}(n; R) \neq 0\} \leq \sum_{q^j \leq y} H(x/q^j, q; R/q, R) + O\left(\frac{x}{\log y}\right),$$

for any parameter $y \leq R^{1/3}$ to be chosen later, where

$$H(X, Y; Z, W) := \#\{n \leq X : P^-(n) > Y, \exists d|n \text{ with } Z < d \leq W\}.$$

Combining the methods of [7] and of [13], it is possible to show the *local-to-global* estimate

$$(5.2) \quad H(X, Y; Z, W) \asymp \frac{X}{(\log Z)^2} \sum_{p|a \Rightarrow Y < p \leq Z} \frac{\mu^2(a) L(a; \log(W/Z))}{a}$$

uniformly in $1 \leq Y^2 \leq Z \leq X^{1/3}$ and $Z + Z/(\log Z)^{100} \leq W \leq Z^2$, where

$$L(a; \eta) := \text{meas} \left(\bigcup_{d|a} [\log d - \eta, \log d) \right)$$

with meas denoting the Lebesgue measure on $\mathbb{R}$. Consequently,

$$H(x/q^j, q; R/q, R) \asymp \frac{x/q^j}{(\log R)^2} \sum_{p|a \Rightarrow q < p \leq R} \frac{\mu^2(a) L(a; \log q)}{a}$$

uniformly in $q^j \leq y \leq R^{1/3}$ and $x \geq R^4$. Moreover, note that

$$\begin{aligned} \frac{\log q}{(\log R)^2} \sum_{p|a \Rightarrow q < p \leq R} \frac{L(a; \log q)}{a} &\asymp \frac{1}{(\log R)^2} \sum_{P^+(a') \leq q} \frac{1}{a'} \sum_{p|a \Rightarrow q < p \leq R} \frac{L(a; \log q)}{a} \\ &\leq \frac{1}{(\log R)^2} \sum_{P^+(b) \leq R} \frac{L(b; \log q)}{b} \\ &\asymp \frac{H(x, 1; R/q, R)}{x} \end{aligned}$$

by setting $b = a'a$ and reapplying (5.2). Theorem 1 in [7] then yields the estimate

$$H(x/q^j, q; R/q, R) \ll \frac{x}{q^j \log q} \cdot \left(\frac{\log q}{\log R} \right)^\delta \left(\log \frac{\log R}{\log q} \right)^{-3/2},$$

uniformly in $2 \leq q^j \leq y \leq R^{1/3}$ and $x \geq R^4$. Inserting this bound to (5.1), we deduce that

$$\#\{n \leq x : M_{f_0}(n; R) \neq 0\} \ll \frac{x}{(\log R)^\delta (\log \log R)^{3/2}} + \frac{1}{\log y}.$$

Selecting $y = \exp((\log R)^\delta (\log \log R)^{3/2})$ completes the proof of (1.5).

Remark 5.1. It is possible to construct integers n for which $M_{f_0}(n; R)$ can get quite large. Indeed, let $y \geq 3$ and $k \in \mathbb{N}$ be two parameters such that the interval $(y, 2^{1/k}y)$ contains at least $2k$ primes, and let $q_1 < \dots < q_{2k}$ be the smallest such primes. Then we set $n = 2q_1 \cdots q_{2k}$ and $R = 2y^k$. By (1.6),

$$M_{f_0}(n; R) = \sum_{\substack{d|q_1 \cdots q_{2k} \\ R/2 < d \leq R}} \mu(d).$$

The choice of R implies that the above sum runs over all divisors d of $q_1, \dots, q_{2k}$ with precisely k prime factors, so that

$$M_{f_0}(n; R) = (-1)^k \binom{2k}{k}.$$

Optimizing the choice of k and y , and using the fact that there infinitely many y 's such that $\pi(y + \sqrt{y \log y}) - \pi(y) \gg \sqrt{y/\log y}$ (see, for example, [12, Exercice 5, p. 266]), we find that there integers n such that $|M_f(n; R)| \gg n^{c/\log \log n}$, for any fixed $c < \frac{\log 2}{2}$.

On the other hand, such extreme values of $M_{f_0}(n; R)$ are very rare, as Theorem 1.3 indicates.

6. INVERSION FORMULAS

Given $f : \mathbb{R} \rightarrow \mathbb{R}$, $R \geq 2$ and $s \in \mathbb{C}$, we set

$$\widehat{f}_R(s) = \int_0^\infty f\left(\frac{\log x}{\log R}\right) x^{s-1} dx = (\log R) \int_{-\infty}^\infty f(u) R^{su} du,$$

provided that the above integral converges. If f is Lebesgue measurable, supported in $(-\infty, 1]$ and uniformly bounded, which will always be the case for us, then $\widehat{f}_R$ defines an analytic function for $\operatorname{Re}(s) > 0$. If, in addition, $f \in C^j(\mathbb{R})$ for some $j \geq 1$ and the derivatives $f', f'', \dots, f^{(j)}$ are all uniformly bounded, then integrating by parts j times yields the formula

$$(6.1) \quad \widehat{f}_R(s) = \frac{(-1)^j}{s^j (\log R)^{j-1}} \int_{-\infty}^\infty f^{(j)}(u) R^{su} du.$$

In particular, we see that

$$(6.2) \quad \left| \widehat{f}_R(s) \right| \leq \|f^{(j)}\|_\infty \cdot \frac{R^{\operatorname{Re}(s)}}{\operatorname{Re}(s)(|s| \log R)^j}$$

for $\operatorname{Re}(s) > 0$, where we used our assumption that $\operatorname{supp}(f) \subset (-\infty, 1]$.

Now, for $m \in \mathbb{N}$, Perron's inversion formula implies that

$$(6.3) \quad f\left(\frac{\log m}{\log R}\right) = \frac{1}{2\pi i} \int_{\operatorname{Re}(s)=c} \widehat{f}_R(s) m^{-s} ds.$$

In the proof of Theorem 1.3 with $A \geq 1$ and of Theorem 1.6, our assumption that f is a few times differentiable in $\mathbb{R}$ allows us to apply (6.2) and write $M_f(a; R)$ in terms of an absolutely convergent integral, which can easily be truncated at some appropriate height. However, when $A = 0$ in Theorem 1.3, we have $f_0 = \chi_{(-\infty, 1]}$, so that $(\widehat{f_0})_R(s) = 1/s$. Truncating Perron's formula is still feasible but rather technical. Instead, we perform a technical manoeuvre and smoothen f_0 a bit. We consider a smooth function $h : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\begin{cases} h(x) = 1 & \text{if } x \leq 1 - \eta, \\ 0 \leq h(x) \leq 1 & \text{if } 1 - \eta \leq x \leq 1, \\ h(x) = 0 & \text{if } x \geq 1, \end{cases}$$

where $\eta = 1/(\log R)^C$ is some constant $C = C(k)$ that will be chosen appropriately later. The function h can be chosen so that $h^{(j)}(x) \ll_j \eta^{-j}$, for all $j \in \mathbb{Z}_{\geq 0}$. We claim that, for any fixed $L > 0$ and $k \geq 1$, there is $C = C(k, L)$ such that

$$(6.4) \quad \mathcal{M}_{f_0, 2k}(R) = \mathcal{M}_{h, 2k}(R) + O\left(\frac{1}{(\log R)^L}\right).$$

Indeed, we have that

$$|\mathcal{M}_{f_0, 2k}(R) - \mathcal{M}_{h, 2k}(R)| \leq 2k \sum_{\substack{d_1, \dots, d_{2k-1} \leq R \\ R^{1-\eta} < d_{2k} \leq R}} \frac{\prod_{j=1}^{2k} \mu^2(d_j)}{[d_1, \dots, d_{2k}]} \leq 2k \sum_{m \leq R^{2k}} \frac{\tau(m)^{2k-1}}{m} \sum_{\substack{d|m \\ R^{1-\eta} < d \leq R}} 1,$$

by setting $m = [d_1, \dots, d_{2k}]$ and $d = d_{2k}$. We split the above sum according to whether $\tau(m) \leq (\log R)^B$ or not, where B is some parameter. We then find that

$$\begin{aligned} |\mathcal{M}_{f_0, 2k}(R) - \mathcal{M}_{h, 2k}(R)| &\leq 2k \sum_{\substack{m \leq R^{2k} \\ \tau(m) \leq (\log R)^B}} \frac{(\log R)^{(2k-1)B}}{m} \sum_{\substack{d|m \\ R^{1-\eta} < m \leq R}} 1 \\ &\quad + 2k \sum_{\substack{m \leq R^{2k} \\ \tau(m) > (\log R)^B}} \frac{\tau(m)^{2k+1} (\log R)^{-B}}{m} \\ &\ll_k (\log R)^{(2k-1)B+2-C} + (\log R)^{2k+1-B}. \end{aligned}$$

We choose $B = L + 2^{2k+1}$ and $C \geq (2k-1)2^{2k+1} + 2kL + 2$ to complete the proof of our claim. For the purposes of Theorem 1.3, we may take $L = 1$, so that having $C \geq (2k-1)2^{2k+1} + 2k + 2$ suffices. We also note that

$$\widehat{h}_R(s) = -\frac{1}{s} \int_{1-\eta}^1 h'(u) R^{su} du = -\frac{R^s}{s} \int_0^\eta h'(1-u) R^{-su} du,$$

by (6.1) and the fact that h is constant outside $[1-\eta, 1]$. In particular, this relation implies that $\widehat{h}_R$ has a meromorphic continuation to $\mathbb{C}$ with only a simple pole at $s = 0$ of residue $-\int_0^\eta h'(1-u) du = 1$. We further note that

$$\begin{aligned} (6.5) \quad \frac{d^j}{ds^j} \left(\frac{s \widehat{h}_R(s)}{R^s} \right) &= (-1)^{j-1} \int_0^\eta h'(1-u) (u \log R)^j R^{-su} du \\ &\ll \eta \cdot \eta^{-1} \cdot (\eta \log R)^j = (\eta \log R)^j \quad (s \in \mathbb{C}, |\operatorname{Re}(s)| \leq 1). \end{aligned}$$

Moreover, we have that

$$\begin{aligned} (6.6) \quad \widehat{h}_R(s) &= (\log R) \int_{-\infty}^1 R^{us} du + O(\eta (\log R) R^{\operatorname{Re}(s)}) \\ &= \frac{R^s}{s} + O((\log R)^{-C+1} R^{\operatorname{Re}(s)}) \quad (\operatorname{Re}(s) \geq 0), \end{aligned}$$

a relation that we will use at the very end of the proof of Theorem 1.3.

7. A COMBINATORIAL PROBLEM IN LINEAR ALGEBRA

Recall the notations from Section 1.7. Consider the $2k$ -dimensional vector space of linear forms in the free variables $s_1, \dots, s_{2k}$. Given a subspace V of this space, we define

$$\mathcal{A}(V) = \sum_{\substack{J \in \mathcal{S}^*(2k) \\ s_J \in V}} (-1)^{|J|}.$$

We will prove the following result:

Proposition 7.1. *Let $k \geq 1$ and V be a subspace as above containing the form $s_{[2k]}$.*

- (a) *If $s_j \in V$ for some $j \in \{1, \dots, 2k\}$, then $\mathcal{A}(V) = -1$.*
- (b) *If $\dim(V) = 2k - 1$, then*

$$\mathcal{A}(V) - \dim(V) \leq \binom{2k}{k} - 2k,$$

with equality if, and only, there is a set $J \subset [2k]$ such that $|J| = k$, $1 \in J$, and $V = \text{Span}_{\mathbb{Q}}(\{s_j - s_1\}_{j \in J}, \{s_j + s_1\}_{j \in [2k] \setminus J})$.
(c) If $\dim(V) \leq 2k - 2$, then

$$\mathcal{A}(V) - \dim(V) \leq \binom{2k}{k} - 2k - 2.$$

Proof. (a) If $s_j \in V$, then we immediately see that $\mathcal{A}(V) = -1$ by pairing s_J with $s_{J \cup \{j\}}$ for each $J \subset [2k] \setminus \{j\}$.

(b) We may assume that $s_1, \dots, s_{2k} \notin V$, by part (a). Since $\dim(V) = 2k - 1$ and $s_1 \notin V$, we have that $s_j \equiv r_j s_1 \pmod{V}$, for some $r_j \in \mathbb{Q} \setminus \{0\}$. We may write $r_j = b_j/q$ for some $b_j \in \mathbb{Z} \setminus \{0\}$ and $q \in \mathbb{N}$, so that $s_J \in V$ if, and only if, $b_J = 0$. Therefore

$$\mathcal{A}(V) = -1 + \int_0^1 \prod_{j=1}^{2k} (1 - e(b_j \theta)) d\theta \leq -1 + \int_0^1 \prod_{j=1}^{2k} |1 - e(b_j \theta)| d\theta$$

Hölder's inequality then implies that

$$\mathcal{A}(V) \leq -1 + \prod_{j=1}^{2k} \left(\int_0^1 |1 - e(b_j \theta)|^{2k} d\theta \right)^{\frac{1}{2k}} = -1 + \binom{2k}{k},$$

whence

$$(7.1) \quad \mathcal{A}(V) \leq \binom{2k}{k} - 1.$$

Finally, we claim that (7.1) is an equality if, and only if, the multiset $\{b_1, \dots, b_{2k}\}$ is of the form $\{b, -b, \dots, b, -b\}$ with $b = b_1$ (which must equal q). It is trivial to check the equality if the multiset $\{b_1, \dots, b_{2k}\}$ is of the form $\{b, -b, \dots, b, -b\}$. Conversely, we know that Hölder's inequality above is an equality if, and only if, there exist real numbers $\lambda_1, \dots, \lambda_{2k}$ such that $|1 - e(b_j \theta)| = \lambda_j |1 - e(b_1 \theta)|$ for $t \in [0, 1]$ and $j \in \{1, \dots, 2k\}$. Since $\int_0^1 |1 - e(b \theta)| d\theta = 4/\pi$ for $b \neq 0$, we must have that $\lambda_j = 1$ for all j . Moreover, taking θ close enough to 0, we find that the condition $|1 - e(b_j \theta)| = |1 - e(b_1 \theta)|$ implies that $|b_j| = |b_1|$ for all j . So $\{b_1, \dots, b_{2k}\}$ has ℓ copies of b_1 and $2k - \ell$ copies of $-b_1$, for some $\ell \in \{1, \dots, 2k\}$. Since $b_{[2k]} = 0$ by our assumption that $s_{[2k]} \in V$, we must have that $\ell = k$, which completes the proof of our claim.

(c) Write $\dim(V) = 2k - n$, where $n \geq 2$. By part (a), we may assume that $s_1, \dots, s_{2k} \notin V$.

We first deal with the case $n = 2, k = 2$ by direct computation. In this case, we have $s_1 + s_2 + s_3 + s_4 \in V$ and $s_1, \dots, s_4 \notin V$, by assumption. It is thus easy to see that either $V \cap \{s_I : I \in \mathcal{S}^*(2k)\} = \{s_1 + \dots + s_4\}$ or $V \cap \{s_I : I \in \mathcal{S}^*(2k)\} = \{s_1 + \dots + s_4, s_J\}$, for some J containing two elements. In any case, $\mathcal{A}(V) \leq 2$, as required. This completes the proof of part (c) when $n = 2$ and $k = 2$.

We now assume that either $n > 2$ or $k > 2$. Choose a maximal subset of linear forms $\{s_{j_1}, \dots, s_{j_{n'}}\}$ that are linearly independent when reduced mod V . Clearly, $n' = n$. Moreover, a permutation of the variables $s_1, \dots, s_{2k}$ allows to assume without loss of generality that $j_i = i$ for each i . Then

$$s_j \equiv \sum_{i=1}^n r_{i,j} s_i \pmod{V} \quad (1 \leq j \leq 2k),$$

for certain $r_{i,j} \in \mathbb{Q}$. We write $r_{i,j} = b_{i,j}/q$, where $b_{i,j} \in \mathbb{Z}$ and $q \in \mathbb{N}$, so that $s_J \in V$ if, and only if, $b_{i,J} := \sum_{j \in J} b_{i,j} = 0$ for each $i \in \{1, \dots, n\}$. Thus

$$\mathcal{A}(V) + 1 = \int_{[0,1]^n} \prod_{j=1}^{2k} (1 - e(b_{1,j}\theta_1 + \dots + b_{n,j}\theta_n)) d\theta_1 \cdots d\theta_n.$$

We set

$$J_m = \{1 \leq j \leq 2k : b_{n-m+1,j} = \dots = b_{n,j} = 0\} \quad (0 \leq m \leq n),$$

so that $J_0 = [2k]$, $J_m = \emptyset$. Note here that $b_{i,j} = \mathbf{1}_{i=j} \cdot q$ for $1 \leq i, j \leq n$, by construction. In particular, $|J_m \setminus J_{m+1}| \geq 1$ for $1 \leq m \leq n-1$. Then

$$\begin{aligned} \mathcal{A}(V) + 1 &\leq \int_{[0,1]^{n-1}} \prod_{j \in J_1} |1 - e(b_{1,j}\theta_1 + \dots + b_{n-1,j}\theta_{n-1})| \\ &\quad \times \left(\int_0^1 \prod_{j \in [2k] \setminus J_1} |1 - e(b_{1,j}\theta_1 + \dots + b_{n,j}\theta_n)| d\theta_n \right) d\theta_1 \cdots d\theta_{n-1}. \end{aligned}$$

By Hölder's inequality, the innermost integral is bounded by

$$\prod_{j \in [2k] \setminus J} \left(\int_0^1 |1 - e(b_{1,j}\theta_1 + \dots + b_{n,j}\theta_n)|^{2k-|J_1|} d\theta_n \right)^{\frac{1}{2k-|J_1|}}.$$

Since $b_{n,j} \neq 0$, we make the change of variables $\theta_n \rightarrow b_{1,j}\theta_1 + \dots + b_{n,j}\theta_n$ and use periodicity to find that

$$\begin{aligned} \int_0^1 |1 - e(b_{1,j}\theta_1 + \dots + b_{n,j}\theta_n)|^{2k-|J_1|} d\theta_n &= \frac{1}{|b_{n,j}|} \int_0^{|b_{n,j}|} |1 - e(\theta)|^{2k-|J_1|} d\theta \\ &= \int_0^1 |1 - e(\theta)|^{2k-|J_1|} d\theta. \end{aligned}$$

We set

$$M(\lambda) = \int_0^1 |1 - e(\theta)|^\lambda d\theta = 2^\lambda \int_0^1 |\sin(\pi\theta)|^\lambda d\theta.$$

Thus we find that

$$H_n(B) \leq M(2k - |J_1|) \int_{[0,1]^{n-1}} \prod_{j \in J_1} |1 - e(b_{1,j}\theta_1 + \dots + b_{n-1,j}\theta_{n-1})| d\theta_1 \cdots d\theta_{n-1}.$$

We repeat the same process to obtain

$$\begin{aligned} \mathcal{A}(V) + 1 &\leq M(2k - |J_1|) M(|J_1| - |J_2|) \\ &\quad \times \int_{[0,1]^{n-1}} \prod_{j \in J_2} |1 - e(b_{1,j}\theta_1 + \dots + b_{n-1,j}\theta_{n-1})| d\theta_1 \cdots d\theta_{n-2} \\ &\leq M(2k - |J_1|) M(|J_1| - |J_2|) \cdots M(|J_{n-2}| - |J_{n-1}|) M(|J_{n-1}|). \end{aligned}$$

Thus

$$(7.2) \quad \mathcal{A}(V) + 1 \leq \sup\{M(\lambda_1) \cdots M(\lambda_n) : \lambda_1 + \dots + \lambda_n = 2k, \lambda_j \geq 1 \ (1 \leq j \leq n)\}.$$

By Cauchy-Schwarz, for any positive reals x, y we have

$$2(xy)^{(A+B)/2} = 2(xy)^B (xy)^{(A-B)/2} \leq (xy)^B (x^{A-B} + y^{A-B}) = x^A y^B + x^B y^A.$$

Thus, applying this with $x = |\sin \theta_1|$, $y = |\sin \theta_2|$ we find

$$\begin{aligned} M(\lambda_1)M(\lambda_2) &= \frac{2^{\lambda_1+\lambda_2}}{2} \int_0^1 \int_0^1 \left(|\sin(\pi\theta_1)^{\lambda_1} \sin(\pi\theta_2)^{\lambda_2}| + |\sin(\pi\theta_1)^{\lambda_1} \sin(\pi\theta_2)^{\lambda_2}| \right) d\theta_1 d\theta_2 \\ &\geq 2^{\lambda_1+\lambda_2} \left(\int_0^1 |\sin(\pi\theta)|^{(\lambda_1+\lambda_2)/2} d\theta \right)^2 = M\left(\frac{\lambda_1+\lambda_2}{2}\right)^2. \end{aligned}$$

In particular, $\log M(\lambda)$ is a convex function. It is then easy to see that supremum in (7.2) is attained when $\lambda_j = 1$ for $n-1$ of the indices $j \in \{1, \dots, n\}$, and with the remaining λ_j being equal to $2k-n+1$. So

$$\mathcal{A}(V) \leq M(1)^{n-1}M(2k-n+1) - 1.$$

Thus, it suffices to show that

$$(7.3) \quad M(1)^{n-1}M(2k-n+1) < \binom{2k}{k} - n = M(2k) - n$$

for $2 \leq n \leq 2k-1$ and $k \geq 2$.

Firstly, consider $n=2$ and $k \geq 3$. The function $k \rightarrow M(1)M(2k-1)/M(2k)$ is decreasing in k by the convexity of $\log M(\lambda)$. Thus

$$M(1)M(2k-1) \leq \frac{M(1)M(3)}{M(4)}M(2k) = \frac{64}{9\pi^2} \binom{2k}{k} < \binom{2k}{k} - 2.$$

Here we have used the fact $M(1) = 4/\pi$, $M(3) = 32/3\pi$ and performed a quick computation to verify $64\binom{2k}{k}/(9\pi^2) < \binom{2k}{k} - 2$ for all $k \geq 3$.

Now consider $3 \leq n \leq 2k-1$. The functions $n \rightarrow M(1)^{n-1}M(2k-n+1)$ and $k \rightarrow M(2k-2)/M(2k)$ are decreasing in n by the convexity of $\log M(\lambda)$. Thus we have

$$\begin{aligned} M(1)^{n-1}M(2k-n+1) &\leq M(1)^2M(2k-2) \\ &\leq \frac{M(1)^2M(2)}{M(4)}M(2k) \\ &= \frac{4}{3\pi^2} \binom{2k}{k} \\ &< \binom{2k}{k} - 2k + 1 = M(2k) - 2k + 1. \end{aligned}$$

Here we have performed a short computation to verify the final inequality. This completes the proof of the proposition. $\square$

8. CONTOUR INTEGRATION

In this section we begin our attack on Theorem 1.3. All implied constants might depend on k and on A . We will actually prove a result that is a little weaker than Theorem 1.3: we will show that there exist constants $c_{k,A}$ and c'_k for which

$$(8.1) \quad \mathcal{M}_{f_A, 2k}(R) = c_{k,A}(\log R)^{\mathcal{E}_{k,A}} + O((\log R)^{\mathcal{E}_{k,A}-1})$$

and

$$(8.2) \quad \mathcal{M}_{\tilde{f}_0, 2k}(R) = c'_k(\log R)^{\binom{2k}{k}-2k} + O((\log R)^{\binom{2k}{k}-2k-1}),$$

where $\widetilde{f}_0(x) = f_0(x) - f_0(x + \frac{\log 2}{\log R})$, so that

$$M_{\widetilde{f}_0}(n; R) = \sum_{\substack{d|n \\ R/2 < d \leq R}} \mu(d).$$

Notice that we do not claim here that $c_{k,A} \neq 0$ and $c_k \neq 0$, as is required in order to prove Theorem 1.3. We do obtain a very complicated expression for these constants, but we are unable to prove they are non-zero (or evaluate them at all). Showing that $c_{k,A} > 0$ and $c'_k > 0$ is the objective of Section 9.

8.1. Initial preparations. We will first prove relation (8.1). The proof of relation (8.2) is very similar, and we indicate the necessary changes in the end of Section 8.

We note that

$$(8.3) \quad (\widehat{f_A})_R(s) = \frac{A!R^s}{(\log R)^A s^{A+1}}.$$

This decays fast when $A \geq 1$, but we have a problem when $A = 0$. However, recall from the end of Section 1.1, that

$$\mathcal{M}_{f_0,2k}(R) = \mathcal{M}_{h,2k}(R) + O\left(\frac{1}{\log R}\right),$$

where h is a smooth function such that $h(x) = 1$ for $x \leq 1 - 1/(\log R)^C$ and $h(x) = 0$ for $x \geq 1$, for some constant $C \geq (2k-1)2^{2k+1} + 2k + 2$ to be chosen later. Therefore relation (8.1) is reduced to showing that

$$\mathcal{M}_{g,2k}(R) = c_{k,A}(\log R)^{\mathcal{E}_{k,A}} + O((\log R)^{\mathcal{E}_{k,A}-1}),$$

where $g = h$ when $A = 0$, and $g = f_A$ when $A \geq 1$. For any $\lambda > 1$, which will be chosen to be sufficiently large in terms of k , and for

$$T := \exp\{(\log \log R)^2\},$$

we have that

$$\begin{aligned} \mathcal{M}_{g,2k}(R) &= \frac{1}{(2i\pi)^{2k}} \int \cdots \int \sum_{\substack{m_j \in \mathbb{N} \\ 1 \leq j \leq 2k \\ \text{Re}(s_j) = \lambda^j / \log R}} \frac{\prod_{j=1}^{2k} \mu(m_j) m_j^{-s_j}}{[m_1, \dots, m_{2k}]} \left(\prod_{j=1}^{2k} \widehat{g}_R(s_j) \right) ds_{2k} \cdots ds_1 \\ &= I_{g,2k}(R) + O\left(\frac{1}{(\log R)^2}\right), \end{aligned}$$

where

$$I_{g,2k}(R) := \frac{1}{(2i\pi)^{2k}} \int \cdots \int \sum_{\substack{m_j \in \mathbb{N} \\ 1 \leq j \leq 2k \\ \text{Re}(s_j) = \lambda^j / \log R \\ |\text{Im}(s_j)| \leq T}} \frac{\prod_{j=1}^{2k} \mu(m_j) m_j^{-s_j}}{[m_1, \dots, m_{2k}]} \left(\prod_{j=1}^{2k} \widehat{g}_R(s_j) \right) ds_{2k} \cdots ds_1,$$

with the truncation at height T being accomplished by noticing that $\widehat{g}_R(s) \ll (\log R)^{O(1)}/|s|^2$ for $\text{Re}(s) = 1/\log R$, a consequence of (6.2) when $A = 0$.

Recall that $s_I := \sum_{i \in I} s_i$ for $I \in \mathcal{S}(2k)$. We set

$$\begin{aligned} P(\mathbf{s}) &= \sum_{\substack{m_j \in \mathbb{N} \\ 1 \leq j \leq 2k}} \frac{\prod_{j=1}^{2k} \mu(m_j) m_j^{-s_j}}{[m_1, \dots, m_{2k}]} \prod_{I \in \mathcal{S}^*(2k)} \zeta(1 + s_I)^{(-1)^{|I|}} \\ &= \prod_p \left\{ \left(1 - \frac{1}{p} + \frac{1}{p} \prod_{j=1}^{2k} \left(1 - \frac{1}{p^{s_j}} \right) \right) \prod_{I \in \mathcal{S}^*(2k)} \left(1 - \frac{1}{p^{1+s_I}} \right)^{(-1)^{|I|}} \right\}, \end{aligned}$$

which is analytic when $\operatorname{Re}(s_j) > -1/4^k$ for all j , as well as

$$F(\mathbf{s}) := P(\mathbf{s}) \prod_{j=1}^{2k} \frac{(\log R)^A \widehat{g}_R(s_j)}{R^{s_j} \zeta(1 + s_j)^{A+1}}$$

and

$$e_I := \begin{cases} +1 & \text{if } |I| = \text{even}, \\ A & \text{if } |I| = 1, \\ -1 & \text{otherwise,} \end{cases}$$

so that

$$I_{g,2k}(R) = \frac{1}{(2i\pi)^{2k}} \int \cdots \int_{\substack{\operatorname{Re}(s_j) = \lambda^j / \log R \\ |\operatorname{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k}} \frac{F(\mathbf{s}) R^{s_1 + \cdots + s_{2k}}}{(\log R)^{2kA}} \prod_{I \in \mathcal{S}^*(2k)} \zeta(1 + s_I)^{e_I} ds_{2k} \cdots ds_1.$$

We claim that F is analytic and non-zero in the region

$$(8.4) \quad \Omega := \{\mathbf{s} \in \mathbb{C}^{2k} : |\operatorname{Re}(s_j)| \leq 1/(\log T)^{4/3}, \ |\operatorname{Im}(s_j)| \leq T \ (1 \leq j \leq 2k)\}$$

and that its derivatives satisfy the bound

$$(8.5) \quad \frac{\partial^{j_1 + \cdots + j_{2k}} F}{\partial s_1^{j_1} \cdots \partial s_{2k}^{j_{2k}}}(\mathbf{s}) \ll_{j_1, \dots, j_{2k}} \frac{(\log \log R)^{O(j_1 + \cdots + j_{2k})}}{(|s_1| + 1) \cdots (|s_{2k}| + 1)}$$

for such $\mathbf{s}$. Indeed, we know that there is some absolute constant $c_0 > 0$ such that

$$(8.6) \quad \zeta^{(j)}(s), \left(\frac{1}{\zeta} \right)^{(j)}(s) \ll_j \frac{1}{|s-1|^j} + \log^{j+1}(|t|+2) \quad \text{whenever } \sigma \geq 1 - \frac{c_0}{\log(|t|+2)},$$

a consequence of the classical zero-free region for ζ (see, for example, Titchmarsh [23, Theorems 3.8 and 3.11]). Moreover, we have that

$$(8.7) \quad \frac{d^j}{ds^j} \left(\frac{s^{A+1} \widehat{g}_R(s)}{R^s} \right) \ll \frac{1}{(\log R)^A} \quad (s \in \mathbb{C}, \ j \in \mathbb{Z}_{\geq 0}),$$

an estimate that follows from (6.6) when $A = 0$ and from the formula (8.3) for $(\widehat{f_A})_R$ otherwise. Relations (8.6) and (8.5) then prove our claims about G . We write $\mathcal{C}_{2k}$ for the class of functions in $2k$ variables that have this property, and similarly $\mathcal{C}_j$ for such functions in j variables, for any $j \in \mathbb{N}$.

8.2. Contour shifting. We will simplify $I_{g,2k}(R)$ and prove (8.1) by a $2k$ -dimensional contour shifting argument that we will demonstrate in an iterative fashion. The general idea is to move the variables s_j to the left in a certain order that will be specified below and to pick up pole contributions, coming from solutions to linear equations of the form $s_I = 0$, $I \in \mathcal{S}^*(2k)$. We need to set-up some notation to carry out the iterative argument.

Assume that the first variable we move to the left is s_{j_1} and we have picked the contribution of some pole with $s_{I_1} = 0$, where $j_1 \in I_1$. We use this equation to write s_{j_1} in terms of s_j , $j \in \{1, \dots, 2k\} \setminus \{j_1\}$. Then s_I becomes a linear form $L_{1,I}$ in the variables s_j , $j \in \{1, \dots, 2k\} \setminus \{j_1\}$. Trivially, $L_{1,I} = 0$ if, and only if, $I \in \mathcal{I}_1 := \{\emptyset, I_1\}$.

Next, we choose some other variable s_{j_2} in a way to be specified below, and move the s_{j_2} contour to the left to pick the pole contribution from some other linear equation $L_{1,I_2} = 0$. It is implicit here that s_{j_2} has a non-zero coefficient in the linear form L_{1,I_2} , and we use this fact to write s_{j_2} in terms of s_j , $j \in \{1, \dots, 2k\} \setminus \{j_1, j_2\}$. Then s_I becomes a linear form $L_{2,I}$ in the variables s_j , $j \in \{1, \dots, 2k\} \setminus \{j_1, j_2\}$. Some of these linear forms will vanish identically. Call the set of these forms which vanish identically $\mathcal{I}_2$.

We continue as above. At the N -th step, we have moved to the left some variables $s_{j_1}, \dots, s_{j_N}$, and we have picked poles from the equations $s_{I_n} = 0$, $n \leq N$, which are all linearly independent over $\mathbb{Q}$. In order to highlight the connection with Section 7, we let

$$V_N = \text{Span}_{\mathbb{Q}}(s_{I_1}, \dots, s_{I_N}) \quad \text{and} \quad \mathcal{I}_N = \{I \in \mathcal{S}(2k) : s_I \in V_N\},$$

so that $\dim(V_N) = N$. We use these equations to write $s_{j_1}, \dots, s_{j_N}$ in terms of the remaining variables, so that s_I becomes a linear form $L_{N,I}$. Note that $I \in \mathcal{I}_N$ if, and only if, $L_{N,I} = 0$. We will also show that we may permute the variables $s_1, \dots, s_{2k}$ in a way that allows us to assume that $j_n = 2k - n + 1$ for all n .

Given sets $I_1, \dots, I_N$ as above, call a pair $(\mathbf{h}, d)$, where $d \in \mathbb{Z}_{\geq 0}$ and $\mathbf{h} = \{h_{n,I} : 0 \leq n \leq N, I \in \mathcal{S}^*(2k)\}$ is a vector of non-negative integers, admissible if:

- $0 = h_{0,I} \leq h_{1,I} \leq \dots \leq h_{N,I}$ for $I \in \mathcal{S}^*(2k)$;
- If $I \in \mathcal{I}_n \setminus \mathcal{I}_{n-1}$ for some $n \in \{1, \dots, N\}$, then $h_{m,I} = h_{n,I}$ for all $m \geq n$;
- $H_N \geq N + d$, where

$$H_N := \sum_{I \in \mathcal{I}_N \setminus \{\emptyset\}} (-1)^{|I|} - \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_N} h_{N,I} + (A+1) \cdot \#\{1 \leq j \leq 2k : \{j\} \in \mathcal{I}_N\}.$$

We claim that $I_{g,2k}(R)$ after N contour shiftings as described above equals a linear combination of terms of the form

$$\begin{aligned} J_N(I_1, \dots, I_N; \mathbf{h}, d) &:= \frac{(\log R)^{H_N - N - d - 2kA}}{(2i\pi)^{2k-N}} \int \cdots \int_{\substack{\text{Re}(s_j) = \lambda_j / \log R \\ |\text{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k-N}} G(\mathbf{s}) R^{E_N(\mathbf{s})} \\ &\quad \times \prod_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_N} (\zeta^{e_I})^{(h_{N,I})} (1 + L_{N,I}(\mathbf{s})) ds_{2k-N} \cdots ds_1, \end{aligned}$$

up to an error term of size $O(e^{-(1+o(1))(\log \log R)^2})$, with $\lambda_j / \lambda_{j-1} \geq \lambda$, $E_N(\mathbf{s}) := L_{N,[2k]}(\mathbf{s})$ and G a function in the class $\mathcal{C}_{2k-N}$ such that

$$G(\mathbf{s}) = F(L_{N,\{1\}}(\mathbf{s}), \dots, L_{N,\{2k\}}(\mathbf{s})) \quad \text{when} \quad d = 0.$$

Note that $I_{g,2k}(R)$ is the above integral when $N = 0$ and $h_I = 0$ for all I , with the convention that $L_{0,I}(s_1, \dots, s_{2k}) = s_I$. Now, we need to understand what happens to $J_N(\mathbf{I}; \mathbf{h}, d)$ when

we move some appropriate variable to the left. We write

$$E_N(\mathbf{s}) = \gamma_1 s_1 + \cdots + \gamma_{2k-N} s_{2k-N}$$

and denote by j_{N+1} the largest element of $\{1, \dots, 2k - N\}$, if it exists, for which $\gamma_{j_{N+1}} \neq 0$. We note that if λ is big enough, then $\text{Re}(E_N(\mathbf{s}))$ has constant sign throughout the region of integration, unless $E_N = 0$ (i.e. $[2k] \in \mathcal{I}_N$). As a matter of fact, if $E_N \neq 0$, then $\text{Re}(E_N(\mathbf{s}))$ has the same sign as $\gamma_{j_{N+1}}$. We then separate the four following cases:

- $\text{Re}(E_N(\mathbf{s})) > 0$;
- $E(\mathbf{s}) = 0$ and $N = 2k - 1$;
- $\text{Re}(E_N(\mathbf{s})) < 0$;
- $E(\mathbf{s}) = 0$ and $N \leq 2k - 2$.

We treat each of these cases separately in the subsequent sections.

8.3. Case 1: $\text{Re}(E_N(\mathbf{s})) > 0$. Note that we must have that either $N = 0$, either $k \geq 2$ or $A \geq 1$ for this case to occur: when $N = k = 1$ and $A = 0$, it must be that $E_1 = 0$, since after having moved the s_2 variable, the only possible pole we could have picked is at $s_2 = -s_1$.

Since $\text{Re}(E_N(\mathbf{s})) > 0$, we have $\gamma_{j_{N+1}} > 0$. We make the change of variables

$$s'_j = s_j \ (1 \leq j < j_{N+1}), \ s'_j = s_{j+1} \ (j_{N+1} \leq j \leq 2k - N), \ s'_{2k-N} = s_{j_{N+1}},$$

which corresponds to a cyclic permutation of the variables $s_1, \dots, s_{2k-N}$. We impose the same permutation to elements of the sets $I_1, \dots, I_N$ we have already chosen and, respectively, to the coefficients of the linear forms $L_{n,I}$. We shift s'_{2k-N} contour to the line $\text{Re}(s'_{2k-N}) = -1/(\log T)^{3/2}$. The contribution of the vertical integrals is $\ll (\log R)^{O(1)}/T$. Moreover, when $\text{Re}(s'_{2k-N}) = -1/(\log T)^{3/2}$ and $\text{Re}(s'_j) = O(1/\log R)$ for $j < 2k - N$, then we have that

$$\text{Re}(E_N(\mathbf{s}')) = -\frac{\gamma_{j_{N+1}}}{(\log T)^{3/2}} + O\left(\frac{1}{\log R}\right).$$

It thus follows that the contribution of the integral with $\text{Re}(s'_{2k-N}) = -1/(\log T)^{3/2}$ is $\ll e^{-\sqrt{\log R}}$, say, which is of negligible size. So we need only worry about the poles that the contour shifting introduces.

The poles occur when $L_{N,I_{N+1}}(\mathbf{s}') = 0$ for some $I_{N+1} \in \mathcal{S}(2k) \setminus \mathcal{I}_N$ of even cardinality or of cardinality 1 such that the coefficient of s'_{2k-N} in $L_{N,I_{N+1}}$ is non-zero. Using the relation $L_{N,I_{N+1}}(\mathbf{s}') = 0$ to write s'_{2k-N} as a linear combination of $s'_1, \dots, s'_{2k-N-1}$, say $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$, we deduce that

$$L_{N,I}(s'_1, \dots, s'_{2k-N}) = L_{N,I}(s'_1, \dots, s'_{2k-N-1}, C(s'_1, \dots, s'_{2k-N-1})) =: L_{N+1,I}(s'_1, \dots, s'_{2k-N-1})$$

at this pole. We also set $E_{N+1} = L_{N+1,[2k]}$ and let $\mathcal{I}_{N+1}$ be the set of $I \subset [2k]$ such that $L_{N+1,I} = 0$. We need to understand the order of the pole at $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$. We only look at *generic points* $(s'_1, \dots, s'_{2k-N-1})$: it could be the case that for some measure-zero subset of points, we get a different pole order. For example, for fixed $s_1 \in \mathbb{C}$, the function $s_2 \rightarrow s_1/s_2$ has generically a pole of order 1 at $s_2 = 0$, unless $s_1 = 0$, when there is no pole. This reduced pole order however would not affect an integral over s_1 , because it only occurs for a measure-zero set of s_1 values.

With the above discussion in mind, we let $\nu \geq 0$ be the generic order of the zero of the analytic function

$$G(\mathbf{s}) \prod_{\substack{I \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N \\ e_I < 0, h_{N,I} \geq |e_I| + 1}} (\zeta^{e_I})^{(h_{N,I})} (1 + L_{N,I}(\mathbf{s})) = G(\mathbf{s}) \prod_{\substack{I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N) \\ |I| \geq 3, h_{N,I} \geq 2}} \left(\frac{1}{\zeta}\right)^{(h_{N,I})} (1 + L_{N,I}(\mathbf{s}))$$

at $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$. If $d = 0$ (so that $G(\mathbf{s}) = F(L_{N,\{1\}}(\mathbf{s}), \dots, L_{N,\{2k\}}(\mathbf{s}))$) and $h_{N,I} = 0$ for all $I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N)$, then we note that $\nu = 0$, because F does not vanish in Ω , nor does the product of the zeta values (unless $L_{N+1,I}(\mathbf{s}) = 0$, in which case though we are referring to a non-generic zero). Then the generic order of the pole of the integrand in the definition of $J_N(\mathbf{I}; \mathbf{h}, d)$ at $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$ is

$$\begin{aligned} m &= \sum_{\substack{I \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N \\ |I| = \text{even}}} (h_{N,I} + 1) - \sum_{\substack{I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N) \\ h_{N,I} = 0, |I| \geq 3}} 1 + \sum_{\substack{1 \leq j \leq 2k \\ \{j\} \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N}} (h_{N,\{j\}} + A) - \nu \\ (8.8) \quad &= \sum_{I \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N} (h_{N,I} + (-1)^{|I|}) + (A + 1) \cdot \#\{1 \leq j \leq 2k : \{j\} \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N\} \\ &\quad - \nu - \sum_{\substack{I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N) \\ h_{N,I} \geq 2, |I| \geq 3}} (h_{N,I} - 1). \end{aligned}$$

Note that it could be the case that $m \leq 0$, in which case there is no pole contribution to $J_N(\mathbf{I}; \mathbf{h}, d)$ from the pole with $s_{I_{N+1}} = 0$. Assume, now, that $m \geq 1$. Then, $m + H_N \geq 1 + N + d$ by our assumption that $H_N \geq N + d$. Moreover,

$$\begin{aligned} m + H_N &= \sum_{I \in \mathcal{I}_{N+1} \setminus \{\emptyset\}} (-1)^{|I|} - \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{N+1}} h_{N,I} \\ &\quad + (A + 1) \cdot \#\{1 \leq j \leq 2k : \{j\} \in \mathcal{I}_{N+1}\} - \nu - \sum_{\substack{I \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N \\ |I| = \text{odd} \\ h_{N,I} \geq 1, |I| \geq 3}} (h_{N,I} - 1) \\ (8.9) \quad &= H_{N+1} + \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{N+1}} (h_{N+1,I} - h_{N,I}) - \nu - \sum_{\substack{I \in \mathcal{I}_{N+1} \setminus \mathcal{I}_N \\ |I| = \text{odd} \\ h_{N,I} \geq 2, |I| \geq 3}} (h_{N,I} - 1). \end{aligned}$$

In order to continue, we separate two subcases.

Case 1a: $N = 2k - 1$. In this case, we have that $s'_j = L_{2k-1,\{j\}}(s'_1) = a_j s'_1$ for all j , where $a_j \in \mathbb{Q}$. Thus, the only potential pole is at $s'_1 = 0$. We note that such a pole does occur for some choice of $I_1, \dots, I_{2k-1}$, so that Case 1a is feasible. Indeed, if $k = 1$ and $A \geq 1$, so that $N = 1$, then we note that at least one of the zeta factors must have survived in the numerator after shifting the s_2 contour, and there are none in the denominator, so there is a pole at $s'_1 = 0$. On the other hand, if $k \geq 2$, then if $I_1 = \{2k - 1, 2k\}$, $I_2 = \{2k - 1, 2k - 2\}$ and $I_3 = \{2k - 2, 2k\}$, then $a_{2k} = a_{2k-1} = a_{2k-2} = 0$, and $a_1 \neq 0$. Taking $h_{2k-1,I} = 0$ for all

I , we see that

$$\begin{aligned} m &= \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}} (-1)^{|I|} + (A+1) \# \{1 \leq j \leq 2k : a_j \neq 0\} \\ &= (A+1) \# \{1 \leq j \leq 2k : a_j \neq 0\} \geq 1, \end{aligned}$$

where we used Proposition 7.1(a). Thus we see indeed that there is a genuine pole at $s'_1 = 0$.

Now, if $m \geq 1$, so that there is a genuine pole at $s'_1 = 0$, then we obtain an evaluation of $J_{2k-1}(\mathbf{I}; \mathbf{h}, d)$ as a finite linear combination of powers of $\log R$ (up to an error term of size $O((\log R)^{O(1)}/T)$), the highest of which has exponent

$$H_{2k-1} + m - 2k - 2kA - d = \sum_{I \in \mathcal{S}^*(2k)} (-1)^{|I|} - \nu - \sum_{\substack{I \in \mathcal{S}^-(2k) \setminus \mathcal{I}_{2k-1} \\ h_{2k-1,I} \geq 2, |I| \geq 3}} (h_{2k-1,I} - 1) - d \leq -1,$$

since $\mathcal{I}_{2k} = \mathcal{S}(2k)$ in this case. We thus see that this contributes towards the main term of $I_{g,2k}(R)$ when $A > \frac{1}{2k} \binom{2k}{k} - 1$, $d = 0$, $\nu = 0$ and $h_{2k-1,I} \in \{0, 1\}$ for $I \in \mathcal{S}^-(2k) \setminus \mathcal{I}_{2k-1}$, as well as that $G(s) = F(a_1 s, \dots, a_{2k} s)$. The residue is then

$$G(0) = A!^{2k} F(0, \dots, 0) = A!^{2k}.$$

Case 1b: $N \leq 2k - 2$. Then the contribution of the pole $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$ to $J_N(\mathbf{I}, \mathbf{h})$ is a constant multiple of

$$\begin{aligned} & \frac{(\log R)^{H_N - N - d - 2kA}}{(2i\pi)^{2k-N-1}} \int \cdots \int_{\substack{\operatorname{Re}(s'_j) = \lambda'_j / \log R, |\operatorname{Im}(s'_j)| \leq T \\ 1 \leq j \leq 2k-N-1}} \left. \frac{d^{m-1}}{d(s'_{2k-N})^{m-1}} \right|_{s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})} \\ & \left\{ G(\mathbf{s}') R^{E_N(\mathbf{s}')} \prod_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{N+1}} (\zeta^{e_I})^{(h_{N,I})} (1 + L_{N,I}(\mathbf{s}')) \right\} ds'_{2k-N-1} \cdots ds'_1, \end{aligned}$$

where $\lambda'_j = \lambda_j$ for $j < j_{N+1}$ and $\lambda'_j = \lambda_{j+1}$ for $j \geq j_{N+1}$. Applying the generalized product rule and writing s_j in place of s'_j , we find that the above integral can be expressed as a finite sum of terms of the form

$$\begin{aligned} & c \cdot \frac{(\log R)^{H_N + m - h - N - 1 - d - 2kA}}{(2i\pi)^{2k-N-1}} \int \cdots \int_{\substack{\operatorname{Re}(s'_j) = \lambda'_j / \log R, |\operatorname{Im}(s'_j)| \leq T \\ 1 \leq j \leq 2k-N-1}} \tilde{G}(\mathbf{s}) R^{E_{N+1}(\mathbf{s})} \\ & \times \prod_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{N+1}} (\zeta^{e_I})^{(h_{N+1,I})} (1 + L_{N+1,I}(\mathbf{s})) ds_{2k-N-1} \cdots ds_1, \end{aligned}$$

where $c \ll 1$, $h \in \{0, \dots, m-1\}$, $\tilde{G}$ is in the class $\mathcal{C}_{2k-N-1}$, $h_{N+1,I} \geq h_{N,I}$ with equality if $I \in \mathcal{I}_{N+1} \setminus \{0\}$, and $\sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{N+1}} (h_{N+1,I} - h_{N,I}) \leq h$.

Relation (8.9) implies that the exponent of $\log R$ is $H_{N+1} - N - 1 - d' - 2kA$, with

$$d' = d + \nu + \sum_{\substack{I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N) \\ h_{N,I} \geq 2, |I| \geq 3}} (h_{N,I} - 1) + h - \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{N+1}} (h_{N+1,I} - h_{N,I}) \geq 0.$$

Moreover, we have that

$$H_{N+1} - d' = H_N + m - h \geq N + 1 \quad \Rightarrow \quad H_{N+1} \geq d' + N + 1,$$

as needed. Finally, if $d' = 0$, it is easy to check that $G(\mathbf{s}) = F(L_{N+1, \{1\}}(\mathbf{s}), \dots, L_{N+1, \{2k\}}(\mathbf{s}))$.

This completes the inductive step, which we keep iterating until we arrive to one of Cases 2, 3 and 4 below, or to Case 1a above. The main contribution to $I_{g, 2k}(R)$ when $A \leq \frac{1}{2k} \binom{2k}{k} - 1$ will then come when, after $2k - 1$ applications of Case 1b, we arrive to the following case:

8.4. Case 2: $E_N(\mathbf{s}) = 0$ and $N = 2k - 1$. As in Case 1a, we have that $s_j = L_{2k-1, \{j\}}(s_1) = a_j s_1$ for all j , where $a_j \in \mathbb{Q}$. Then the only possible pole of the integrand is when $s_1 = 0$. If there is such a pole, then it means that $\{1\} \notin \mathcal{I}_{2k-1}$. In this case, the order of this potential pole would be given by (8.8) with $N = 2k - 1$ and $\mathcal{I}_{2k} = \mathcal{S}(2k)$, so that (8.9) implies that

$$(8.10) \quad m + H_{2k-1} - (2k - 1) - 2kA = 1 + \sum_{I \in \mathcal{S}^*(2k)} (-1)^{|I|} - \sum_{\substack{I \in \mathcal{S}^-(2k) \setminus \mathcal{I}_{2k-1} \\ |I| \geq 3, h_{2k-1, I} \geq 2}} (h_{2k-1, I} - 1) - \nu \leq 0.$$

If $m \geq 1$ (i.e. there is a genuine pole at $s_1 = 0$), we find that $H_{2k-1} + 1 - 2k(A + 1) \leq -1$. We then move the line of integration of s_1 to the contour $\sigma_1 = 1/(\log(2 + |t_1|))^{3/2}$, $|\operatorname{Im}(s_j)| \leq T$. No poles are encountered and the vertical integrals contribute $\ll (\log R)^{O(1)}/T$. Moreover, the integral converges fast enough now (even when $A = 0$) that we may remove the condition $|\operatorname{Im}(s_j)| \leq T$ at the cost of an error term of size $\ll (\log R)^{O(1)}/T$. We thus conclude that

$$J_{2k-1}(\mathbf{I}; \mathbf{h}, d) = c \cdot (\log R)^{H_{2k-1} - 2k(A+1) + 1 - d} + O\left(e^{-(1+o(1))(\log \log R)^2}\right),$$

where

$$c = \int_{\sigma_1 = 1/(\log(2 + |t_1|))^{3/2}} G(s_1) \prod_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}} (\zeta^{e_I})^{(h_{2k-1, I})} (1 + a_I s_1) ds_1$$

is some constant. This contributes towards the error term if $m \geq 2$, $d \geq 1$ or $A \leq \frac{1}{2k} \binom{2k}{k} - 1$, and towards the main term if $m = 1$, $d = 0$, $A > \frac{1}{2k} \binom{2k}{k} - 1$ and $h_{2k-1, I} \in \{0, 1\}$ for each $I \in \mathcal{S}^-(2k)$, in which case $H_{2k-1} - 2k(A + 1) + 1 = -1$ by (8.10).

Finally, assume that $m \leq 0$, so that there is no pole at $s_1 = 0$. We move s_1 to the line $\operatorname{Re}(s_1) = 0$. The vertical lines contribute $\ll (\log R)^{O(1)}/T$. Finally, we note that we may extend the range of integration to all $s_1 \in \mathbb{C}$ with $\operatorname{Re}(s_1) = 0$ at the cost of an error term of size $\ll (\log R)^{O(1)}/T$. Consequently,

$$J_N(\mathbf{I}; \mathbf{h}, d) = c \cdot (\log R)^{H_{2k-1} + 1 - 2k(A+1) - d} + O((\log R)^{O(1)}/T),$$

where

$$c = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(it) \prod_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}} (\zeta^{e_I})^{(h_{2k-1, I})} (1 + ia_I t) dt$$

with $a_I := \sum_{j \in I} a_j$. The power of $\log R$ is

$$\begin{aligned} H_{2k-1} + 1 - 2k(A+1) - d &= 1 + \sum_{I \in \mathcal{S}_{2k-1} \setminus \{0\}} (-1)^{|I|} - \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}} h_{2k-1,I} \\ &\quad - (A+1) \cdot \#\{1 \leq j \leq 2k : a_j \neq 0\} - d \\ &= 1 + \mathcal{A}(V_{2k-1}) - \sum_{I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}} h_{2k-1,I} \\ &\quad - (A+1) \cdot \#\{1 \leq j \leq 2k : a_j \neq 0\} - d \end{aligned}$$

in the notation of Proposition 7.1. Clearly, this is maximized when $h_{2k-1,I} = 0$ for all $I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}$ and $d = 0$, in which case

$$(8.11) \quad G(s) = F(a_1 s, \dots, a_{2k} s) = \frac{P(a_1 s, \dots, a_{2k} s)}{\prod_{j=1}^{2k} (a_j s \zeta(1 + a_j s))^{A+1}} + O\left(\frac{(\log(2 + |s|))^{O(1)}}{(1 + |s|)^{(2k-1)(A+1)}}\right)$$

by (6.6). Moreover, if $a_j = 0$ for some $j \in \{1, \dots, 2k\}$, then $\mathcal{A}(V_{2k-1}) = -1$, by Proposition 7.1(a), so that the maximal power of $\log R$ becomes $-(A+1) \cdot \#\{1 \leq j \leq 2k : a_j \neq 0\}$. This is $\leq -A-1$: there is at least one j such that $a_j \neq 0$; otherwise, the dimension of V_{2k-1} would be $2k$, as it would contain the independent forms $s_1, \dots, s_{2k}$, which is a contradiction. Therefore,

$$J_{2k-1}(\mathbf{I}; \mathbf{h}, d) \ll (\log R)^{-A-1}$$

in this case, which is of negligible size. Finally, assume that there is no $a_j = 0$. Then Proposition 7.1(b) implies that the maximal power of $\log R$ is

$$\mathcal{A}(V_{2k-1}) - 2k(A+1) \leq \binom{2k}{k} - 2k(A+1),$$

with equality attained when half of the a_j 's equal $+1$ and the other half -1 . We can then give a formula for $J_{2k-1}(\mathbf{I}; \mathbf{h}, d)$. For simplicity, let us assume that $a_j = 1$ for $j \leq k$ and $a_j = -1$ for $j > k$. Then $a_I = |I \cap [1, k]| - |I \cap (k, 2k]|$, which has the same parity as $|I|$. In particular, $\mathcal{I}^-(2k) \cap \mathcal{I}_{2k-1} = \emptyset$, so that $h_{2k-1,I} = 0$ for all $I \in \mathcal{S}^-(2k)$. Moreover, given $\ell \in \mathbb{Z}$, we have that $a_I = \ell$ for exactly $\binom{2k}{k+|\ell|}$ sets $I \subset [2k]$. Therefore

$$\begin{aligned} J_{2k-1}(\mathbf{I}; \mathbf{h}, 0) &= \frac{(\log R)^{\binom{2k}{k} - 2kA}}{2\pi} \int_{-\infty}^{\infty} \frac{P(it, -it, \dots, it, -it)}{t^{2k(A+1)}} \cdot \frac{\prod_{\ell \text{ even}, \geq 2} |\zeta(1 + i\ell t)|^{2\binom{2k}{k+\ell}}}{\prod_{\ell \text{ odd}, \geq 1} |\zeta(1 + i\ell t)|^{2\binom{2k}{k+\ell}}} dt \\ &\quad + O\left(e^{-(1+o(1))(\log \log R)^2}\right) \end{aligned}$$

This completes the study of this second case. Relation 8.1 will then follow if we can show that Cases 3 and 4 below contribute only towards the error term.

8.5. Case 3: $\operatorname{Re}(E_N(\mathbf{s})) < 0$. In this case, we must have $\gamma_{j_{N+1}} < 0$. We then shift the contours of $s_{2k-N}, s_{2k-N-1}, \dots, s_{j_{N+1}}$ in this order to the lines $\operatorname{Re}(s_j) = \lambda_j/(\log T)^{3/2}$, $j_{N+1} \leq j \leq 2k - N$. If λ is large enough, then we do not encounter any poles and the vertical lines contribute $\ll (\log R)^{O(1)}/T$ when we make this shift. Finally, when $\operatorname{Re}(s_j) = \lambda_j/(\log T)^{3/2}$ for $j_{N+1} \leq j \leq 2k - N$, and $\operatorname{Re}(s_j) = \lambda^j/\log R$ for $1 \leq j < j_{N+1}$, then we have that

$$\operatorname{Re}(E(\mathbf{s})) = -\frac{|\gamma_{j_{N+1}}| \lambda_{j_{N+1}} (1 + O(1/\lambda))}{(\log T)^{3/2}},$$

so that our integrand is $\ll e^{-\sqrt{\log R}}$ if λ is large enough. We thus find that in this case

$$J_N(\mathbf{I}; \mathbf{h}) \ll e^{-(1+o(1))(\log \log R)^2}.$$

8.6. Case 4: $E(\mathbf{s}) = 0$ and $N \leq 2k - 2$. We shift the contours of $s_{2k-N}, s_{2k-N-1}, \dots, s_1$ in this order to the lines $\operatorname{Re}(s_j) = \lambda^j / (\log T)^{3/2}$, $1 \leq j \leq 2k - N$. As before, if λ is large enough, then we do not encounter any poles. The vertical lines contribute $\ll (\log R)^{O(1)}/T$. Finally, we note that the integrand on the new lines of integration is $\ll (\log \log R)^{O(1)}/[(1 + |s_1|) \cdots (1 + |s_{2k-N}|)]$, by (8.6) and our assumption that G is in the class $\mathcal{C}_{2k-N}$. We thus find that

$$J_N(\mathbf{I}; \mathbf{h}, d) \ll (\log R)^{H_N - N - 2kA - d} (\log \log R)^{O(1)}$$

in this case. We need to understand the power of $\log R$. We have that

$$H_N - N - 2kA - d \leq 2k - N + \sum_{I \in \mathcal{I}_N \setminus \{\emptyset\}} (-1)^{|I|} - (A+1) \cdot \#\{1 \leq j \leq 2k : \{j\} \notin \mathcal{I}_N\}.$$

If there is $\{j\} \in \mathcal{I}_N$, then $\sum_{I \in \mathcal{I}_N \setminus \{\emptyset\}} (-1)^{|I|} = -1$ by Proposition 7.1(a). Since $\{j\} \in \mathcal{I}_N$ for at most N j 's for dimension reasons, the power of $\log R$ is

$$\leq 2k - N - 1 - (A+1)(2k - N) = -1 - A(2k - N) \leq -1 - 2 \cdot \mathbf{1}_{A \geq 1},$$

and we can see that the contribution of $J_N(\mathbf{I}; \mathbf{h})$ is towards the error term of (8.1). So assume that there is no $\{j\} \in \mathcal{I}_N$. Then we find that

$$H_N - N - 2kA \leq 2k - N + \sum_{I \in \mathcal{I}_N \setminus \{\emptyset\}} (-1)^{|I|} - 2k(A+1) \leq \binom{2k}{k} - 2k(A+1) - 2,$$

by Proposition 7.1(c), which means that $J_N(\mathbf{I}; \mathbf{h})$ contributes towards the error term in (8.1) in this case too.

8.7. Dyadic intervals. We conclude this section with a brief explanation of the proof of (8.2). We have that

$$\mathcal{M}_{\tilde{f}_0, 2k}(R) = \mathcal{M}_{\tilde{h}, 2k}(R) + O\left(\frac{1}{\log R}\right),$$

where $\tilde{h}(x) = h(x) - h(x + \frac{\log 2}{\log R})$, by the argument leading to (6.4). We then note that

$$\widehat{(\tilde{h})}_R(s) = \widehat{h}_R(s)(1 - 2^{-s}),$$

so that Perron's inversion formula and relation (6.2) imply that

$$\begin{aligned} \mathcal{M}_{\tilde{f}_0, 2k}(R) &= \frac{1}{(2i\pi)^{2k}} \int \cdots \int \sum_{\substack{\operatorname{Re}(s_j) = \lambda^j / \log R \\ 1 \leq j \leq 2k}} \sum_{\substack{m_j \in \mathbb{N} \\ 1 \leq j \leq 2k}} \frac{\prod_{j=1}^{2k} \mu(m_j) m_j^{-s_j}}{[m_1, \dots, m_{2k}]} \left(\prod_{j=1}^{2k} \widehat{h}_R(s_j)(1 - 2^{-s_j}) \right) ds_{2k} \cdots ds_1 \\ &\quad + O\left(\frac{1}{\log R}\right), \end{aligned}$$

where λ and T are as before. We thus find that

$$\mathcal{M}_{\tilde{f}_0, 2k}(R) = \frac{1}{(2i\pi)^{2k}} \int \cdots \int_{\substack{\operatorname{Re}(s_j) = \lambda^j / \log R \\ 1 \leq j \leq 2k}} \tilde{F}(\mathbf{s}) R^{s_1 + \cdots + s_{2k}} \prod_{I \in \mathcal{S}^*(2k)} \zeta(1 + s_I)^{(-1)^{|I|}} + O\left(\frac{1}{\log R}\right),$$

where

$$\tilde{F}(\mathbf{s}) := P(\mathbf{s}) \prod_{j=1}^{2k} \frac{\widehat{h}_R(s_j)(1 - 2^{-s_j})}{R^{s_j}}$$

and P is defined as above. The function $\tilde{F}$ is in the class $\mathcal{C}_{2k}$, since the factor $1 - 2^{-s_j}$ annihilates the pole of $\widehat{h}_R(s_j)$ at $s_j = 0$. We thus see that the above integral has the same shape as the integral $I_{g, 2k}(R)$ with $A = 0$, with the difference that $e_I = -1$ when $|I| = 1$. We thus follow the argument leading to (8.1) when $A = 0$ with the obvious modifications and relation (8.2) follows.

9. LOWER BOUNDS

In this section we complete our proof of Theorem 1.3 by showing that, for fixed $k \in \mathbb{N}$, $A \in \mathbb{Z}_{\geq 0}$ and $\epsilon > 0$, there are positive constants $c'_{k,A} > 0$ and $c''_k > 0$ such that

$$(9.1) \quad \mathcal{M}_{f_A, 2k}(R) \geq c'_{k,A} (\log R)^{\mathcal{E}_{k,A} - \epsilon} + O_\epsilon((\log R)^{\mathcal{E}_{k,A} - 1}),$$

and

$$(9.2) \quad \mathcal{M}_{\tilde{f}_0, 2k}(R) \geq c''_k (\log R)^{D_{k,0} - \epsilon} + O_\epsilon((\log R)^{D_{k,0} - 1}),$$

where $\tilde{f}_0(x) = f_0(x) - f_0(x + \frac{\log 2}{\log R})$, as before. Evidently, this completes the proof of Theorem 1.3, since if the constants in the leading terms there were 0, we would obtain a contradiction to the above lower bounds by letting $R \rightarrow \infty$ in (8.1) and (8.2).

As in the previous section, there is some smooth function h such that $h(x) = 1$ for $x \leq 1 - 1/(\log R)^C$ and $h(x) = 0$ for $x \geq 1$, with $C = (2k - 1)2^{2k+1} + 2k + 2$, so that

$$\mathcal{M}_{f_0, 2k}(R) = \mathcal{M}_{h, 2k}(R) + O\left(\frac{1}{\log R}\right) \quad \text{and} \quad \mathcal{M}_{\tilde{f}_0, 2k}(R) = \mathcal{M}_{\tilde{h}, 2k}(R) + O\left(\frac{1}{\log R}\right),$$

where $\tilde{h}(x) = h(x) - h(x + \frac{\log 2}{\log R})$. So, it suffices to prove that

$$(9.3) \quad \mathcal{M}_{g, 2k}(R) \geq c'_{k,A} (\log R)^{\mathcal{E}_{k,A} - \epsilon} + O((\log R)^{\mathcal{E}_{k,A} - 1}),$$

where $g \in \{h, \tilde{h}\}$ when $A = 0$, and $g = f_A$ when $A \geq 1$.

Positivity is a key to this proof: we will consider the sum restricted to those integers with a convenient prime factorization, which clearly provides a lower bound. The fact that these integers have a convenient prime factorization means the corresponding sum is technically easier to analyze.

9.1. First manipulations. We start by observing that

$$\mathcal{M}_{g,2k}(R) = \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{1}{n} \left(\sum_{d|n} \mu(d) g \left(\frac{\log d}{\log R} \right) \right)^{2k},$$

since $\text{supp}(g) \subset (-\infty, 1]$. So (9.1) follows immediately when $A > -1 + \frac{1}{2k} \binom{2k}{k}$ by noticing that $\mathcal{E}_{k,A} = -1$ in this case and that we always have that $\mathcal{M}_{g,2k}(R) \geq \prod_{p \leq R} (1 - 1/p) \gg 1/\log R$.

For the rest of the proof, we assume that $A \leq -1 + \frac{1}{2k} \binom{2k}{k}$, so that

$$\mathcal{E}_{k,A} = \binom{2k}{k} - 2k(A+1) \geq 0.$$

Let $q_1 < q_2 < \dots$ be the sequence of all prime numbers and set

$$Q = \prod_{j=1}^{A+1} q_j.$$

If $A \geq 1$, or if $A = 0$ and $g = h$, then we restrict our attention to integers of the form Qn with $P^-(n) > Q$, so that

$$\mathcal{M}_{g,2k}(R) \geq \frac{\prod_{p \leq R} (1 - 1/p)}{Q} \sum_{p|n \Rightarrow q_{A+1} < p \leq R} \frac{1}{n} \left(\sum_{J \subset [A+1]} (-1)^{|J|} \sum_{d|n} \mu(d) g \left(\frac{\log(q_J d)}{\log R} \right) \right)^{2k},$$

where $q_J := \prod_{j \in J} q_j$. To this end, we define

$$(9.4) \quad w(x) := \sum_{J \subset [A+1]} (-1)^{|J|} g \left(x + \frac{\log q_J}{\log R} \right),$$

so that

$$(9.5) \quad \mathcal{M}_{g,2k}(R) \geq \frac{\prod_{p \leq R} (1 - 1/p)}{Q} \sum_{p|n \Rightarrow q_{A+1} < p \leq R} \left(\sum_{d|n} \mu(d) w \left(\frac{\log d}{\log R} \right) \right)^{2k}.$$

We further note that $w = \tilde{h}$ when $A = 0$, so that the right hand side of (9.5) is a trivial lower bound for $\mathcal{M}_{\tilde{h},2k}(R)$. Therefore, relation (9.3) will follow in all cases if we can show that

$$(9.6) \quad \begin{aligned} W &:= \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{p|n \Rightarrow q_{A+1} < p \leq R} \left(\sum_{d|n} \mu(d) w \left(\frac{\log d}{\log R} \right) \right)^{2k} \\ &\geq c''_{k,A} (\log R)^{\mathcal{E}_{k,A} - \epsilon} + O_\epsilon((\log R)^{\mathcal{E}_{k,A} - 1}) \end{aligned}$$

for some $c''_{k,A} > 0$, where w is defined by (9.4) with $g = h$ if $A = 0$ and $g = f_A$ if $A \geq 1$.

Next, observe that

$$(9.7) \quad \widehat{w}_R(s) = (\log R) \sum_{J \subset [A+1]} (-1)^{|J|} \int_{-\infty}^{\infty} g \left(u + \frac{\log q_J}{\log R} \right) R^{su} du = \widehat{g}_R(s) \prod_{j=1}^{A+1} (1 - q_j^{-s}).$$

In particular, we see that $\widehat{w}_R$ has an analytic continuation to $\mathbb{C}$ and it satisfies the bound

$$(9.8) \quad \widehat{w}_R(s) \ll \frac{R^{\operatorname{Re}(s)}}{(|s|+1)(\log R)^A} \quad (\operatorname{Re}(s) \geq -1),$$

which follows by the definition of f_A when $A \geq 1$ and by (6.5) otherwise. Finally, we note that we also have the bound

$$(9.9) \quad \widehat{w}_R(s) \ll \frac{R^{\operatorname{Re}(s)}(\log R)^{O(1)}}{1+|s|^2} \quad (\operatorname{Re}(s) \geq 1/\log R),$$

which follows from (6.2) when $A = 0$. (This bound can be shown to hold in a larger range, but the above range is good enough for our purposes.)

Before we apply Perron's inversion formula to write the right hand side of (9.5) in terms of $\widehat{w}_R$, we use positivity again to focus on integers n of a certain convenient form. We set

$$y = \exp\{(\log R)^{1-\epsilon'}\} \quad \text{and} \quad Y = \exp\{(\log R)^{1-\epsilon'/2}\},$$

where ϵ' will be taken to be small enough in terms of ϵ , and write

$$\mathcal{N} = \{n \in \mathbb{N} : p|n \Rightarrow q_{A+1} < p \leq y\}.$$

We then focus on integers of the form $n = mp_1 \cdots p_k$, where $m \in \mathcal{N}$ with $m \leq Y$, and $p_1, \dots, p_k$ are distinct primes from the interval $(R^{1/2}, R]$. For such an integer n , if $d|n$, then either $d = d'$ or $d = d'p_a$, for some $d'|m$ and some $\ell \in \{1, \dots, k\}$. So, we conclude that

$$\begin{aligned} W &\geq \frac{\prod_{p \leq R} (1 - 1/p)}{k!} \sum_{\substack{m \in \mathcal{N} \\ m \leq Y}} \sum_{\substack{\sqrt{R} < p_1, \dots, p_k \leq R \\ \text{distinct primes}}} \frac{1}{mp_1 \cdots p_k} \\ &\quad \times \left(\sum_{\ell=1}^k \sum_{d|m} \mu(d) w \left(\frac{\log(p_\ell d)}{\log R} \right) - \sum_{d|m} \mu(d) w \left(\frac{\log d}{\log R} \right) \right)^{2k}. \end{aligned}$$

Note that the condition that $m \leq Y$ implies that $d < R/(2q_{A+1})$ for all divisors d of m , so that

$$\begin{aligned} \sum_{d|m} \mu(d) w \left(\frac{\log d}{\log R} \right) &= \sum_{J \subset [A+1]} (-1)^{|J|} \sum_{d|m} \mu(d) \left(1 - \frac{\log(q_J d)}{\log R} \right)^A \\ &= \sum_{d|Qm} \mu(d) \left(1 - \frac{\log d}{\log R} \right)^A = 0 \end{aligned}$$

by observing that $\Delta^{(A+1)}(1-x)^A = 0$ by (1.8), and by applying (1.7) with $r = A+1$, since $\omega(Qm) \geq \omega(Q) \geq A+1$ here. We thus conclude that

$$\begin{aligned} W &\geq \frac{\prod_{p \leq R} (1 - 1/p)}{k!} \sum_{\substack{m \in \mathcal{N} \\ m \leq Y}} \sum_{\substack{\sqrt{R} < p_1, \dots, p_k \leq R \\ \text{distinct primes}}} \frac{1}{mp_1 \cdots p_k} \left(\sum_{\ell=1}^k \sum_{d|m} \mu(d) w \left(\frac{\log(p_\ell d)}{\log R} \right) \right)^{2k} \\ &\geq \frac{\prod_{p \leq R} (1 - 1/p)}{k!(\log R)^k} \sum_{\substack{m \in \mathcal{N} \\ m \leq Y}} \sum_{\substack{\sqrt{R} < p_1, \dots, p_k \leq R \\ \text{distinct primes}}} \frac{\prod_{j=1}^k \log p_j}{mp_1 \cdots p_k} \left(\sum_{\ell=1}^k \sum_{d|m} \mu(d) w \left(\frac{\log(p_\ell d)}{\log R} \right) \right)^{2k}. \end{aligned}$$

We note that

$$\sum_{n \in \mathcal{N}, n > Y} \frac{\tau_r(n)}{n} \leq Y^{-1/\log y} \sum_{n \in \mathcal{N}} \frac{\tau_r(n)}{n^{1-1/\log y}} \ll \frac{1}{(\log R)^B},$$

for any fixed $B \geq 1$ and $r \in \mathbb{N}$. Therefore, we may drop the conditions that $m \leq Y$ and that the p_ℓ 's are distinct at the cost of an admissible error term, finding that

$$W \geq \frac{\prod_{p \leq R} (1 - 1/p)}{k! (\log R)^k} \sum_{m \in \mathcal{N}} \sum_{\sqrt{R} < p_1, \dots, p_k \leq R} \frac{\prod_{j=1}^k \log p_j}{m p_1 \cdots p_k} \left(\sum_{\ell=1}^k \sum_{d|m} \mu(d) w \left(\frac{\log(p_\ell d)}{\log R} \right) \right)^{2k} - O \left(\frac{1}{(\log R)^{100}} \right).$$

Next, we expand the $2k$ -th power as follows:

$$\left(\sum_{\ell=1}^k \sum_{d|m} \mu(d) w \left(\frac{\log(p_\ell d)}{\log R} \right) \right)^{2k} = \sum_{\substack{J_1 \cup \dots \cup J_k = [2k] \\ J_i \cap J_j = \emptyset}} \prod_{\ell=1}^k \prod_{j \in J_\ell} \sum_{d_j | m} \mu(d_j) w \left(\frac{\log(p_\ell d_j)}{\log R} \right),$$

with the convention that if $J_\ell = \emptyset$, then the corresponding factor equals 1. Clearly, the summands corresponding to those k -tuples $\mathbf{J} = (J_1, \dots, J_k)$ all of whose components have an even number of terms are non-negative. We write $\mathcal{J}$ for the set of k -tuples $\mathbf{J}$ such that $(J_1, \dots, J_k)$ is a partition of $[2k]$ and either $|J_\ell| = 2$ for all ℓ , or there is some ℓ such that $|J_\ell|$ is an odd number. Then

$$\left(\sum_{\ell=1}^k \sum_{d|m} \mu(d) w \left(\frac{\log(p_\ell d)}{\log R} \right) \right)^{2k} \geq \sum_{\mathbf{J} \in \mathcal{J}} \prod_{\ell=1}^k \prod_{j \in J_\ell} \sum_{d_j | m} \mu(d_j) w \left(\frac{\log(p_\ell d_j)}{\log R} \right).$$

Moreover, if $D \in \mathcal{N}$, then

$$\sum_{\substack{m \in \mathcal{N} \\ D|m}} \frac{1}{m} = \frac{1}{D} \prod_{q_{A+1} < p \leq y} \left(1 - \frac{1}{p} \right)^{-1} = \frac{c_1 \prod_{p \leq y} (1 - 1/p)}{D}$$

with $c_1 = \prod_{p \leq q_{A+1}} (1 - 1/p) \asymp 1$. So, we conclude that

$$(9.10) \quad W \geq \frac{c_1 \prod_{y < p \leq R} (1 - 1/p)}{k! (\log R)^k} \sum_{\mathbf{J} \in \mathcal{J}} \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\mu(d_1) \cdots \mu(d_{2k})}{[d_1, \dots, d_{2k}]} \times \prod_{\ell=1}^k \sum_{\sqrt{R} < p_\ell \leq R} \frac{\log p_\ell}{p_\ell} \prod_{j \in J_\ell} w \left(\frac{\log(p_\ell d_j)}{\log R} \right) - O \left(\frac{1}{(\log R)^{100}} \right).$$

For the convenience of notation, set

$$W(\mathbf{J}) = \frac{1}{(\log R)^k} \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\mu(d_1) \cdots \mu(d_{2k})}{[d_1, \dots, d_{2k}]} \prod_{\ell=1}^k \sum_{\sqrt{R} < p_\ell \leq R} \frac{\log p_\ell}{p_\ell} \prod_{j \in J_\ell} w \left(\frac{\log(p_\ell d_j)}{\log R} \right),$$

so that

$$W \geq \frac{c_1 \prod_{y < p \leq R} (1 - 1/p)}{k!} \sum_{\mathbf{J} \in \mathcal{J}} W(\mathbf{J}) - O \left(\frac{1}{(\log R)^{100}} \right).$$

We will show that the dominant contribution comes from the terms $\mathbf{J}$ with $|J_\ell| = 2$ for all ℓ .

9.2. Mellin transformation. Fix a choice of sets $\mathbf{J} = (J_1, \dots, J_k) \in \mathcal{J}$ and let

$$\mathcal{L} = \{1 \leq \ell \leq k : J_\ell \neq \emptyset\}.$$

For $\ell \notin \mathcal{L}$, the sum over p_ℓ is

$$\sum_{\sqrt{R} < p_\ell \leq R} \frac{\log p_\ell}{p_\ell} = \frac{\log R}{2} + O\left(e^{-c\sqrt{\log R}}\right).$$

So

$$W(\mathbf{J}) = \frac{2^{|\mathcal{L}|-k}}{(\log R)^{|\mathcal{L}|}} \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\mu(d_1) \cdots \mu(d_{2k})}{[d_1, \dots, d_{2k}]} \prod_{\ell \in \mathcal{L}} \sum_{p_\ell > \sqrt{R}} \frac{\log p_\ell}{p_\ell} \prod_{j \in J_\ell} w\left(\frac{\log(p_\ell d_j)}{\log R}\right) + O\left(\frac{1}{(\log R)^{100}}\right),$$

where the condition that $p_\ell \leq R$ was dropped because it is implied by the fact that $\text{supp}(w) \subset (-\infty, 1]$.

Next, we use Perron's formula $2k$ times to write each appearance of w as an integral of $\hat{w}_R$. We thus find that

$$W(\mathbf{J}) = \frac{2^{|\mathcal{L}|-k} (\log R)^{-|\mathcal{L}|}}{(2\pi i)^{2k}} \int \cdots \int_{\substack{\text{Re}(s_j) = 1/\log R \\ 1 \leq j \leq 2k}} \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \left(\prod_{\ell \in \mathcal{L}} \sum_{p_\ell > \sqrt{R}} \frac{\log p_\ell}{p_\ell^{1+s_{J_\ell}}} \right) \times \left(\prod_{j=1}^{2k} \hat{w}_R(s_j) \right) ds_1 \cdots ds_{2k} + O\left(\frac{1}{\log R}\right),$$

with the notational convention that $s_J := \sum_{j \in J} s_j$. By possibly re-indexing the variables $s_1, \dots, s_{2k}$, we may assume that $\mathcal{L} = \{1, \dots, L\}$, where $L = |\mathcal{L}|$, and that $\max J_\ell = 2k - L + \ell$ for all $\ell \in \{1, \dots, L\}$. We want to move the variables $s_{2k-L+1}, \dots, s_{2k}$ to the left. First, we need some bounds on the sum over p_ℓ . We note that

$$\sum_{p > \sqrt{R}} \frac{\log p}{p^{1+s}} = -\frac{\zeta'}{\zeta}(1+s) + O(1) - \sum_{p \leq R^{1/2}} \frac{\log p}{p^{1+s}}$$

for $\text{Re}(s) \geq -1/3$. Using standard bounds on the Riemann zeta function (see, for example, Titchmarsh [23, Theorem 3.11]), we find that

$$\begin{aligned} \sum_{p > \sqrt{R}} \frac{\log p}{p^{1+s}} &\ll \frac{1}{|s|} + \log(2 + |t|) + R^{\max\{0, -\sigma\}/2} \sum_{p \leq \sqrt{R}} \frac{\log p}{p} \\ &\ll R^{\max\{0, -\sigma\}/2} \log(R + |t|), \end{aligned}$$

where $s = \sigma + it$, as usual. Moreover, note that if $\sigma_j \geq -1/\log y$ for all $j \in \{1, \dots, 2k\}$, then

$$\begin{aligned} \left| \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \right| &\leq \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu^2(d_j) d_j^{1/\log y}}{[d_1, \dots, d_{2k}]} \\ &\leq \prod_{p \leq y} \left(1 + \frac{p^{1/\log y} (4^k - 1)}{p} \right) \ll (\log y)^{4^k - 1}, \end{aligned}$$

using the estimate $p^{1/\log y} = 1 + O(\log p / \log y)$ for $p \leq y$.

We are now ready to move the variables $s_{2k-L+1}, \dots, s_{2k}$ to the left. First, we move the variable s_{2k} to the line $\operatorname{Re}(s_{2k}) = -1/\log y$. (Here we can use (9.9) to justify that the contribution of the vertical lines at infinity is 0.) We pick up a simple pole from the sum over p_{2k} when $s_{J_L} = 0$. The integrand when $\operatorname{Re}(s_{2k}) = -1/\log y$ is

$$\ll (\log y)^{4^k - 1} (\log R)^{L-1} \log(R+|t|) R^{1/(2 \log y)} \prod_{j=1}^{2k} |\widehat{w}_R(s_j)| \ll \frac{(\log R)^{O(1)} \log(2+|t|) R^{-1/(2 \log y)}}{(|s_1|^2 + 1) \cdots (|s_{2k}|^2 + 1)}$$

by (9.9). So we find that

$$\begin{aligned} W(\mathbf{J}) &= \frac{2^{L-k} (\log R)^{-L}}{(2\pi i)^{2k-1}} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=1/\log R \\ 1 \leq j \leq 2k-1 \\ s_{J_L}=0}} \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \left(\prod_{\ell=1}^{L-1} \sum_{p_\ell > R^{1/2}} \frac{\log p_\ell}{p_\ell^{1+s_{J_\ell}}} \right) \\ &\quad \times \left(\prod_{j=1}^{2k} \widehat{w}_R(s_j) \right) ds_1 \cdots ds_{2k-1} + O\left(\frac{1}{(\log R)^{100}} \right). \end{aligned}$$

Note that there are no more poles left for the variables $s_j \in J_L \setminus \{2k\}$, so we move all of them to the line $\operatorname{Re}(s_j) = 0$. Then we repeat the same argument by moving s_{2k-1} to the left, then s_{2k-2} , and so on and so forth, until all the sums over primes have been removed and replaced by poles. Writing $s_j = it_j$, we conclude that

$$\begin{aligned} W(\mathbf{J}) &= \frac{2^{L-k} (\log R)^{-L}}{(2\pi)^{2k-L}} \int \cdots \int_{\substack{t_1, \dots, t_{2k} \in \mathbb{R} \\ t_{J_\ell}=0 \ (1 \leq \ell \leq L)}} \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-it_j}}{[d_1, \dots, d_{2k}]} \left(\prod_{j=1}^{2k} \widehat{w}_R(it_j) \right) dt_1 \cdots dt_{2k-L} \\ &\quad + O\left(\frac{1}{(\log R)^{100}} \right). \end{aligned} \tag{9.11}$$

9.3. An auxiliary result. Before we continue with the estimation of $W(\mathbf{J})$, we establish a preliminary result.

Lemma 9.1. *For $z \geq y \geq 3$ and $t \in \mathbb{R}$, we have that*

$$\sum_{y < p \leq z} \frac{1}{p^{1+it}} = \begin{cases} \log \left(\frac{\log z}{\log y} \right) + O(1) & \text{if } |t| \leq 1/\log z, \\ \log \left(\frac{1}{|t| \log y} \right) + O(1) & \text{if } 1/\log z < |t| \leq 1/\log y, \\ O(1) & \text{if } y \geq |t| \geq 1/\log y. \end{cases}$$

Finally, if $|t| \geq y$, then

$$\left| \sum_{y < p \leq z} \frac{1}{p^{1+it}} \right| \leq \log \left(\frac{\log(\min\{|t|, z\})}{\log y} \right) + O(1).$$

Proof. If $|t| \leq 1/\log z$, then we note that $p^{-it} = 1 + O(\log p / \log z)$ for $p \leq z$, so that

$$\sum_{y < p \leq z} \frac{1}{p^{1+it}} = \sum_{y < p \leq z} \frac{1}{p} + O(1) = \log \left(\frac{\log z}{\log y} \right) + O(1),$$

as claimed. For $|t| \geq 1/\log y$ and $y \geq |t|$, then we note that

$$\sum_{y < p \leq z} \frac{1}{p^{1+it}} \ll 1$$

by relation (4.4) in [14]. Next, if $1/\log z \leq |t| \leq 1/\log y$, then we apply the results we just proved to deduce that

$$\sum_{y < p \leq e^{1/|t|}} \frac{1}{p^{1+it}} = \log \left(\frac{\log e^{1/|t|}}{\log y} \right) + O(1) = \log \left(\frac{1}{|t| \log y} \right) + O(1)$$

and that

$$\sum_{e^{1/|t|} < p \leq z} \frac{1}{p^{1+it}} \ll 1.$$

Finally, if $|t| \geq y$, then we note that

$$\sum_{|t| < p \leq z} \frac{1}{p^{1+it}} \ll 1,$$

so that

$$\begin{aligned} \left| \sum_{y < p \leq z} \frac{1}{p^{1+it}} \right| &= \left| \sum_{y < p \leq \min\{|t|, z\}} \frac{1}{p^{1+it}} \right| + O(1) \leq \sum_{y < p \leq \min\{|t|, z\}} \frac{1}{p} + O(1) \\ &\leq \log \left(\frac{\log(\min\{|t|, z\})}{\log y} \right) + O(1). \end{aligned}$$

□

9.4. Lower bound for the main term. We now return to the study of the quantity $W(\mathbf{J})$, defined by (9.11). First, we show that the term when $|J_\ell| = 2$ for all ℓ contributes what we claim to be our main term. Here $L = |\mathcal{L}| = k$ and, by possibly permuting the t_j 's, we may assume that $J_\ell = \{\ell, k + \ell\}$ for all $\ell \in \{1, \dots, k\}$. Then $t_{k+\ell} = -t_\ell$.

We want to study the integrand in (9.11) and show that it is non-negative for all choices of $t_1, \dots, t_k$. We have that

$$\prod_{j=1}^{2k} \widehat{w}_R(it_j) = \prod_{j=1}^k \widehat{w}_R(it_j) \widehat{w}_R(-it_j) = \prod_{j=1}^k |\widehat{w}_R(it_j)|^2 = \prod_{j=1}^k |t_j^{A+1} \widehat{g}_R(it_j)|^2 \prod_{a=1}^{A+1} \frac{|1 - q_a^{-it_j}|^2}{t_j^2},$$

which is clearly non-negative for all $t_1, \dots, t_k$. Moreover, if $t_j \ll 1$ for all j , then relations (9.7) and (6.6) imply that

$$\prod_{j=1}^{2k} \widehat{w}_R(it_j) \geq \frac{c_2}{(\log R)^{2kA}}$$

for some $c_2 > 0$. Furthermore,

$$\sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-it_j}}{[d_1, \dots, d_{2k}]} = \prod_{q_{A+1} < p \leq y} \left(1 + \frac{\lambda_p(\mathbf{t})}{p} \right),$$

where

$$(9.12) \quad \lambda_p(\mathbf{t}) := \sum_{I \in \mathcal{S}^*(2k)} (-1)^{|I|} p^{-it_I} = -1 + \prod_{j=1}^{2k} (1 - p^{-it_j}) = -1 + \prod_{j=1}^k |1 - p^{-it_j}|^2 \geq -1$$

for such $\mathbf{t}$. In particular, we see that

$$\sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-it_j}}{[d_1, \dots, d_{2k}]} \geq 0$$

for $\mathbf{t}$ as above.

Since the integrand in (9.11) is non-negative, we may restrict the range of integration to any region we wish. We assume that $t_1 \in [1, 2]$ and that $t_j \in [t_1, t_1 + 1/\log y]$ for $2 \leq j \leq k$. The volume is $\asymp 1/(\log y)^{k-1}$. Moreover, under this assumption, we find that

$$\lambda_p(\mathbf{t}) = -1 + |1 - p^{it_1}|^{2k} + O\left(\frac{\log p}{\log y}\right).$$

Therefore

$$\sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-it_j}}{[d_1, \dots, d_{2k}]} \gg \exp \left\{ \sum_{p \leq y} \frac{-1 + |1 - p^{it_1}|^{2k}}{p} \right\} \asymp (\log y)^{\binom{2k}{k} - 1},$$

since

$$|1 - e^{i\theta}|^{2k} = \sum_{j+j'=2k} \binom{2k}{k} (-1)^{j'} e^{i(j-j')\theta}$$

and the terms with $j \neq j'$ can be handled by Lemma 9.1. Hence

$$W(\mathbf{J}) \gg \frac{(\log y)^{\binom{2k}{k} - k}}{(\log R)^{k+2kA}}$$

in this case. So the total contribution to the right hand side of (9.10) from such $\mathbf{J}$ is $\geq c_2(\log y)^{\binom{2k}{k}-k+1}/(\log R)^{(2A+1)k}$ for some $c_2 > 0$, which is greater than the claimed main term in (9.6) if ϵ' is small enough in terms of ϵ and k .

9.5. Upper bound for the error term. It remains to show that the contribution of the $\mathbf{J}$'s for which at least one of the $|J_a|$'s is odd is smaller than what we have above. Before we get started, we note that

$$\lambda_p(\mathbf{t}) = -1 + \prod_{j=1}^{2k} (1 - p^{-it_j}) = -1 + \prod_{j=1}^{2k} (p^{it_j/2} - p^{-it_j/2}) = -1 + (-4)^k \prod_{j=1}^k \sin\left(\frac{t_j \log p}{2}\right)$$

whenever $t_1 + \dots + t_{2k} = 0$, which is the case here. In particular, $-4^k \leq \lambda_p(\mathbf{t}) + 1 \leq 4^k$, whence

$$\left| \sum_{d_1, \dots, d_{2k} \in \mathcal{N}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-it_j}}{[d_1, \dots, d_{2k}]} \right| = \left| \prod_{q_{A+1} < p \leq y} \left(1 + \frac{\lambda_p(\mathbf{t})}{p}\right) \right| \ll \prod_{4^k < p \leq y} \left(1 + \frac{\lambda_p(\mathbf{t})}{p}\right).$$

Next, we split the region of integration in (9.11) into various subsets. First, we note that, by a dyadic decomposition argument and (9.9), we have that

$$W(\mathbf{J}) \ll \frac{(\log T)^{2k}}{(\log R)^{L+2kA}} \max_{1 \leq T_1, \dots, T_{2k} \leq T} \frac{\Lambda(\mathbf{T})}{T_1 \cdots T_{2k-L}} + \frac{(\log R)^{O(1)}}{T},$$

where

$$\Lambda(\mathbf{T}) := \int \cdots \int \prod_{\substack{T_j \leq |t_j| + 1 \leq 2T_j \\ 1 \leq j \leq 2k-L \\ t_{J_\ell} = 0 \ (\ell \in \mathcal{L})}} \left(1 + \frac{\lambda_p(\mathbf{t})}{p}\right) dt_1 \cdots dt_{2k-L}.$$

We take

$$T = \exp\{(\log \log R)^2\},$$

and fix a choice of $T_1, \dots, T_{2k-L}$ as above. We will further break the region of integration according to which sums $t_J = \sum_{j \in J} t_j$ are small. Indeed, by Lemma 9.1, only if there are such configurations, the product $\prod_{q_{A+1} < p \leq y} (1 + \lambda_p(\mathbf{t})/p)$ can become large. Note that if t_{J_1} and t_{J_2} are both small, so is any linear combination of t_{J_1} and t_{J_2} . Thus, we are naturally led to the following definition: given free variables $x_1, \dots, x_{2k}$ and $J \subset [2k]$, we define the linear form

$$L_J(x_1, \dots, x_{2k}) := \sum_{j \in J} x_j,$$

and, given $\mathcal{I} \subset \mathcal{S}(2k)$, we set

$$V_B(\mathcal{I}) := \left\{ \sum_{I \in \mathcal{I}} \frac{a_I}{q_I} \cdot L_I : a_I, q_I \in \mathbb{Z} \cap [-B, B] \right\}.$$

For the simplicity of notation, we also set

$$V(\mathcal{I}) := V_\infty(\mathcal{I}) = \text{Span}_{\mathbb{Q}}(\{L_I : I \in \mathcal{I}\}).$$

By Gaussian elimination, and since there are only finitely many linear forms L_I , $I \subset [2k]$, there is some finite $B = B(k)$ such that, for any $\mathcal{I}$, if $L_J \in V(\mathcal{I})$, then $L_J \in V_B(\mathcal{I})$. We fix B to be of this size from now on.

We set $m = \lfloor \log \log y \rfloor$ and, to each $\mathbf{t}$, we associate the sets

$$\mathcal{I}_j = \mathcal{I}_j(\mathbf{t}) := \{I \subset [2k] : e^m |t_I| + 1 \leq e^{j+1}\} \quad (0 \leq j \leq m).$$

Note that if $L_I \notin V_B(\mathcal{I}_m)$, then $|t_I| > 1$. Therefore, Lemma 9.1 implies that

$$\prod_{4^k < p \leq y} \left(1 + \frac{\lambda_p(\mathbf{t})}{p}\right) \ll (\log \log R)^{O(1)} \prod_{\substack{I \in \mathcal{S}^*(2k) \\ L_I \in V_B(\mathcal{I}_m)}} \min \left\{ \log y, \frac{1}{|t_I|} \right\}^{(-1)^{|I|}}.$$

If $L_I \in V_B(\mathcal{I}_j) \setminus V_B(\mathcal{I}_{j-1})$, where $\mathcal{I}_{-1} = \{\emptyset\}$ so that $V_B(\mathcal{I}_{-1}) = \{0\}$, then $I \notin \mathcal{I}_{j-1}$, which means that $e^m |t_I| + 1 > e^j$. On the other hand, since $L_I \in V_B(\mathcal{I}_j)$, then we find that $e^m |t_I| \leq B \cdot |\mathcal{I}_j| \cdot e^{j-m}$. We thus conclude that

$$\min \left\{ \log y, \frac{1}{|t_I|} \right\} \asymp e^{m-j} \quad \text{for } L_I \in V_B(\mathcal{I}_j) \setminus V_B(\mathcal{I}_{j-1}), \quad 0 \leq j \leq m.$$

Therefore

$$\prod_{4^k < p \leq y} \left(1 + \frac{\lambda_p(\mathbf{t})}{p}\right) \ll (\log \log R)^{4^k} \prod_{j=0}^m e^{(m-j)F_j},$$

where

$$F_j := \sum_{\substack{I \subset [2k] \\ L_I \in V_B(\mathcal{I}_j) \setminus V_B(\mathcal{I}_{j-1})}} (-1)^{|I|}.$$

(Here we use the fact that there are only finitely many indices j for which $\mathcal{I}_j \neq \mathcal{I}_{j-1}$.) Summing over the $\ll (\log \log R)^{O(1)}$ choices for $\mathcal{I}_0, \mathcal{I}_1, \dots, \mathcal{I}_m$, we conclude that

$$\Lambda(\mathbf{T}) \ll (\log \log R)^{O(1)} \max_{\mathcal{I}_m \supset \dots \supset \mathcal{I}_0 \supset \{J_1, \dots, J_L\}} \left(\nu(\mathbf{T}, \mathcal{I}) \cdot \prod_{j=0}^m e^{(m-j)F_j} \right),$$

where $\nu(\mathbf{T}, \mathcal{I})$ denotes the $(2k - L)$ -dimensional Lebesgue measure of $(t_1, \dots, t_{2k-L}) \in \mathbb{R}$ such that $T_j \leq |t_j| + 1 \leq 2T_j$ for $j \leq 2k - L$ and

$$\{I \subset [2k] : e^m |t_I| + 1 < e^{j+1}\} = \mathcal{I}_j \quad (0 \leq j \leq m),$$

where the variables $t_{2k-L+1}, \dots, t_{2k}$ are defined via the equations $t_{J_\ell} = 0$ for $\ell \in \{1, \dots, 2k\}$. (In particular, $\mathcal{I}_0 \supset \{J_1, \dots, J_L\}$.)

Next, we use linear algebra to understand $\nu(\mathbf{T}, \mathcal{I})$. If

$$D_j = \dim V(\mathcal{I}_j),$$

then we may find sets $I_1, \dots, I_{D_m}$ such that, for each $j \leq m$, $\{L_{I_1}, \dots, L_{I_{D_j}}\}$ is a basis of $V(\mathcal{I}_j)$. We may also assume that $\{J_\ell : 1 \leq \ell \leq L\}$ is contained in $\{I_1, \dots, I_{D_1}\}$. We apply Gaussian elimination to the asymptotic formulas

$$t_I = O(e^{j-m}) \quad (I \in \{I_1, \dots, I_{D_j}\})$$

to find that D_j of the variables $t_1, \dots, t_{2k}$, say the variables $\{t_i : i \in \mathcal{D}_j\}$ with $|\mathcal{D}_j| = D_j$, can be put within $O(e^{j-m})$ of a linear combination of the other variables. We can also arrange

the sets $\mathcal{D}_0, \dots, \mathcal{D}_L$ to satisfy $\mathcal{D}_0 \subset \dots \subset \mathcal{D}_L$. (Recall that $\mathcal{I}_j \neq \mathcal{I}_{j-1}$ for only finitely many j 's.) We may therefore prove that

$$\nu(\mathbf{T}, \mathbf{I}) \ll (\log y)^L \left(\prod_{j=0}^m e^{(D_j - D_{j-1})(j-m)} \right) \left(\prod_{\substack{1 \leq j \leq 2k \\ j \notin \mathcal{D}_0 \cup \dots \cup \mathcal{D}_L}} T_j \right),$$

where the extra factor $(\log y)^L$ is included because we are not integrating over the variables $t_{2k-L+1}, \dots, t_{2k}$, which are fixed via the conditions $t_{J_\ell} = 0$, $1 \leq \ell \leq L$. By the above discussion, we find that

$$\Lambda(\mathbf{T}) \ll T_1 \cdots T_{2k-L} (\log y)^L (\log \log R)^{O(1)} \max_{\mathcal{I}_m \supset \dots \supset \mathcal{I}_0 \supset \{J_1, \dots, J_L\}} \prod_{j=0}^m e^{(m-j)(F_j - (D_j - D_{j-1}))}.$$

We note that

$$\begin{aligned} \sum_{j=0}^m (m-j)(F_j - (D_j - D_{j-1})) &= \sum_{j=0}^{m-1} \sum_{i=1}^{m-j} (F_j - (D_j - D_{j-1})) \\ &= \sum_{i=1}^m \sum_{j=0}^{m-i} (F_j - (D_j - D_{j-1})) \\ &= \sum_{i=1}^m (\mathcal{A}(V(\mathcal{I}_{m-i})) - \dim(V(\mathcal{I}_{m-i}))) \end{aligned}$$

in the notation of Section 7, provided that B is large enough. Since $\exp$ is a convex function, we have that

$$e^{\sum_{j=0}^m (m-j)(F_j - (D_j - D_{j-1}))} \leq \frac{1}{m} \sum_{j=0}^{m-1} e^{m(\mathcal{A}(V(\mathcal{I}_j)) - \dim(V(\mathcal{I}_j)))} \asymp \frac{1}{m} \sum_{j=0}^{m-1} (\log y)^{\mathcal{A}(V(\mathcal{I}_j)) - \dim(V(\mathcal{I}_j))}$$

We then conclude that

$$W(\mathbf{J}) \ll \frac{(\log \log R)^{O(1)} (\log y)^L}{(\log R)^{L+2kA}} \max_{\mathcal{I} \supset \{J_1, \dots, J_L\}} (\log y)^{\mathcal{A}(V(\mathcal{I})) - \dim(V(\mathcal{I}))},$$

where $F(\mathcal{I})$ is defined as above.

The above discussion reduces (9.6) to proving that

$$(9.13) \quad \mathcal{A}(V(\mathcal{I})) - \dim(V(\mathcal{I})) \leq \binom{2k}{k} - 2k - 1$$

whenever $\mathcal{I}$ contains the sets $J_1, \dots, J_L$ and at least one of the J_i 's has an odd number of elements. This follows directly by Proposition 7.1, thus completing the proof of (9.1) and, hence, of Theorem 1.3.

10. ON THE ANATOMY OF INTEGERS CONTRIBUTING TO $\mathcal{M}_{f,2k}(R)$

This section is devoted to establishing Theorems 1.6 and 1.4. Throughout this section, given $n \in \mathbb{N}$ and $R \geq 1$, we adopt the notations

$$\Omega(n; R) := \sum_{p^\alpha \parallel n, p \leq R} \alpha \quad \text{and} \quad \Omega(n; r, R) := \sum_{p^\alpha \parallel n, r < p \leq R} \alpha.$$

10.1. An estimate for the logarithmic means. We start by proving a preliminary result, where the integer n is weighted by $1/n$. The transition to the uniform weights will be accomplished in the subsequent section.

Theorem 10.1. *Let $R \geq 2$, $k \in \mathbb{N}$ and $A \in \mathbb{Z}_{\geq 0}$.*

(a) *If $A > \frac{1}{2k} \binom{2k}{k} - 1$ and $\eta \in [\log 2 / \log R, 1]$, then*

$$\prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{\Omega(n; R^\eta) M_{f_A}(n; R)^{2k}}{n} \ll_{k,A} \frac{\eta^{2k}}{\log R} + (\log R)^{\binom{2k}{k} - 2k(A+1)} \log \log R.$$

(b) *If $A \leq \frac{1}{2k} \binom{2k}{k} - 1$ and $1 - 1/\binom{2k}{k} \leq v \leq 2 - \epsilon$, then*

$$\prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{v^{\Omega(n)} M_{f_A}(n; R)^{2k}}{n} \ll_{\epsilon,k,A} (\log R)^{v \binom{2k}{k} - 2k(A+1)} (\log \log R)^{O_k(1)}.$$

Proof. All implied constants might depend on k, A and ϵ .

(a) First of all, we claim we may restrict our attention to small enough η . To see this, it suffices to show that

$$(10.1) \quad \sum_{P^+(n) \leq R} \frac{\Omega(n; R^\delta, R) M_{f_A}(n; R)^{2k}}{n} \ll_\delta 1,$$

for any fixed $\delta > 0$. To prove this, we first note that if $(a, b) = 1$, then

$$(10.2) \quad \begin{aligned} M_{f_A}(ab; R) &= \sum_{d|a} \sum_{d'|b} \mu(d) \mu(d') f_A \left(\frac{\log d}{\log R} + \frac{\log d'}{\log R} \right) \\ &= \sum_{\substack{d'|b \\ d' \leq R}} \mu(d') \left(\frac{\log(R/d')}{\log R} \right)^A \sum_{d|a} \mu(d) f_A \left(\frac{\log d}{\log(R/d')} \right) \end{aligned}$$

by the definition of f_A . Hölder's inequality then yields

$$(10.3) \quad M_{f_A}(ab; R)^{2k} \leq \tau(b)^{2k-1} \sum_{\substack{d'|b \\ d' \leq R}} \mu^2(d') \left(\frac{\log(R/d')}{\log R} \right)^{2kA} M_{f_A}(a; R/d')^{2k}.$$

Therefore, writing $n = ab$, where $b = \prod_{p^v \parallel n, R^\delta < p \leq R} p^v$, we find that

$$\begin{aligned} &\sum_{P^+(n) \leq R} \frac{\Omega(n; R^\delta, R) M_{f_A}(n; R)^{2k}}{n} \\ &\leq \sum_{p|b \Rightarrow R^\delta < p \leq R} \frac{\tau(b)^{2k-1} \Omega(b)}{b} \sum_{\substack{d'|b \\ d' \leq R}} \mu^2(d') \left(\frac{\log(R/d')}{\log R} \right)^{2kA} \sum_{P^+(a) \leq R} \frac{M_{f_A}(a; R/d')^{2k}}{a} \\ &\ll \sum_{p|b \Rightarrow R^\delta < p \leq R} \frac{\tau(b)^{2k-1} \Omega(b)}{b} \sum_{\substack{d'|b \\ d' \leq R}} \mu^2(d') \left(\frac{\log(R/d')}{\log R} \right)^{2kA} \frac{\log R}{\log(R/d')} \ll_\delta 1 \end{aligned}$$

by Theorem 1.3, since $A \geq 1$ here. This proves (10.1), so for the rest of the proof, we may assume that η is sufficiently small.

Expanding $\Omega(n, R^\eta)$, we have that

$$\begin{aligned} S &:= \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{\Omega(n; R^\eta) M_{f_A}(n; R)^{2k}}{n} \\ &= \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{\substack{q \leq R^\eta \\ j \geq 1}} \frac{j}{q^j} \sum_{\substack{P^+(m) \leq R \\ q \nmid m}} \frac{M_{f_A}(mq^j; R)^{2k}}{m} \\ &\leq \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{q \leq R^\eta} \frac{q}{(q-1)^2} \sum_{P^+(m) \leq R} \frac{\left(\sum_{d|m} \mu(d) \left\{f_A\left(\frac{\log d}{\log R}\right) - f_A\left(\frac{\log(qd)}{\log R}\right)\right\}\right)^{2k}}{m}. \end{aligned}$$

where we used (10.2) with $a = q^j$ and $b = m$. Expanding the $2k^{th}$ power, we find that

$$S \leq \sum_{q \leq R^\eta} \frac{q}{(q-1)^2} \sum_{\substack{P^+(d_j) \leq R \\ 1 \leq j \leq 2k}} \frac{\prod_{j=1}^{2k} \mu(d_j) \left\{f_A\left(\frac{\log d_j}{\log R}\right) - f_A\left(\frac{\log(qd_j)}{\log R}\right)\right\}}{[d_1, \dots, d_{2k}]},$$

Here $A \geq 1$, so that $\widehat{(f_A)}(s) = A!R^s/(s^{A+1}(\log R)^A)$ decays fast, and Perron inversion implies that

(10.4)

$$S \leq \sum_{q \leq R^\eta} \frac{q}{(q-1)^2} \int \cdots \int_{\substack{\text{Re}(s_j)=1/\log R \\ 1 \leq j \leq 2k}} \sum_{d_1, \dots, d_{2k} \geq 1} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \prod_{j=1}^{2k} \frac{A!R^{s_j}(1 - q^{-s_j})}{(\log R)^A s_j^{A+1}} ds_1 \cdots ds_{2k}.$$

The above integral is amenable to the methods of Section 8. Precisely, we note that the integrand is of the form

$$A!^{2k} \left(\frac{\log q}{\log R}\right)^{2k} \frac{P(\mathbf{s}) R^{s_{[2k]}}}{(\log R)^{2k(A-1)}} \prod_{I \in \mathcal{S}^+(2k)} \zeta^{e_I}(1 + s_I) \prod_{j=1}^{2k} \frac{(1 - q^{-s_j})/(s_j \log q)}{(s_j \zeta(1 + s_j))^A},$$

where $P(\mathbf{s})$ is as in Section 8, $e_I = +1$ for $I \in \mathcal{S}^+(2k)$, $e_I = -1$ for $I \in \mathcal{S}^-(2k)$ with $|I| > 1$, and $e_I = A - 1 \geq 0$ for $|I| = 1$. If $q \leq R^\delta$ with δ small enough, then the argument leading to (8.1) yields that

$$\begin{aligned} (10.5) \quad &\int \cdots \int_{\substack{\text{Re}(s_j)=1/\log R \\ 1 \leq j \leq 2k}} \sum_{d_1, \dots, d_{2k} \geq 1} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \prod_{j=1}^{2k} \frac{A!R^{s_j}(1 - q^{-s_j})}{(\log R)^A s_j^{A+1}} ds_1 \cdots ds_{2k} \\ &\ll \frac{(\log q)^{2k}}{(\log R)^{2k+1}} + (\log R)^{\binom{2k}{k} - 2k(A+1)}, \end{aligned}$$

with the first term coming from Case 1a and the second one from Case 2. Inserting the above inequality into (10.4) completes the proof of part (a).

(b) We will use a variation of the argument of Section 9. The fact that Proposition 7.1 requires $s_{[2k]} = 0$ complicates the proof; otherwise, a direct application of the method of Section 9.5 would be possible.

Call S' the sum in question. As usual, we may replace f_0 by a sufficiently smooth function h . So write $g = h$ if $A = 0$, and $g = f_A$ otherwise. Note that, since $M_g(n; R)$ depends only on the square-free kernel of n , we have that

$$S' = \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{\mu^2(n) v^{\omega(n)}}{\varphi_v(n)} M_{f_A}(n; R)^{2k}$$

with $\varphi_v(n) = \prod_{p|n} (p-v)$. Set $y = R^{c/\log \log R}$ for a small enough but fixed c . Given an integer n , we decompose it as $n = ab$ with $P^+(a) \leq y < P^-(b)$. If $\omega(a) \leq A+1$, then $M_{f_A}(n; R) \ll 4^{k\omega(b)}$, and we immediately find that such n 's contribute at most $\ll (\log \log R)^{O(1)} / \log R$. Otherwise, we sum over all possible choices $a = qa'$ with $\omega(q) = A+1$ to deduce that

$$(10.6) \quad S' \leq v^{A+1} \sum_{\substack{P^+(q) \leq y \\ \omega(q) = A+1}} \frac{\mu^2(q)}{\varphi_v(q)} S'(q) + O\left(\frac{(\log \log R)^{O(1)}}{\log R}\right),$$

where

$$S'(q) := \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{\substack{P^+(a') \leq y < P^-(b) \leq R \\ (a', q) = 1}} \frac{\mu^2(a') v^{\omega(a')}}{\varphi_v(a')} M_{f_A}(qa'b; R)^{2k}.$$

Next, we apply (10.3) with $a = a'q$ to find that

$$S'(q) \leq \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{\substack{P^+(a') \leq y \\ y < P^-(b) \leq R \\ (a', q) = 1}} \frac{\mu^2(a') v^{\omega(a')} \tau(b)^{2k-1}}{\varphi_v(a'b)} \sum_{\substack{d|b \\ d \leq R}} \left(\frac{\log(R/d)}{\log R}\right)^{2kA} M_{f_A}(a'q; R/d)^{2k}.$$

We write $b = cd$ and note that the sum over c is $\leq (\log \log R)^{O(1)}$ by our choice of y . Moreover, $\varphi_v(n) \gg n/(\log \log n)^v$. Thus,

$$S'(q) \ll (\log \log R)^{O(1)} \sum_{\substack{d \leq R \\ P^-(d) > y}} \frac{4^{k\Omega(d; y, R)}}{d} S''(d, q) \leq (\log \log R)^{O(1)} \sum_{\substack{d \leq R \\ P^-(d) > y}} \frac{S''(d, q)}{d^{1-2k/\log y}},$$

where

$$\begin{aligned} S''(d, q) &:= \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{\substack{P^+(a') \leq y \\ (a', q) = 1}} \frac{\mu^2(a') v^{\omega(a')}}{\varphi_v(a')} \left(\frac{\log(R/d)}{\log R}\right)^{2kA} M_{f_A}(a'q; R/d)^{2k} \\ &\leq \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{\substack{P^+(a) \leq y \\ (a, q) = 1}} \frac{v^{\Omega(a)}}{a} \left(\frac{\log(R/d)}{\log R}\right)^{2kA} M_{f_A}(aq; R/d)^{2k}. \end{aligned}$$

Before we proceed to the estimation of $S''(d, q)$, we note that

$$(10.7) \quad S'(q) \ll (\log \log R)^{O(1)} \sum_{d \leq R} \frac{(\lambda^+ * 1)(d)}{d^{1-2k/\log y}} S''(d, q),$$

where $(\lambda^+(m))_{m \leq R^\delta}$ is an upper bound sieve with δ small enough, constructed using the fundamental lemma of sieve methods, taking $\kappa = 1$ in [12, Lemma 6.3, p. 159]. Then the

Dirichlet series $\sum_d (1 * \lambda^+)(d)/d^s$ has a simple pole at $s = 1$ of residue $\sum_m \lambda^+(m)/m \asymp (\log \log R)^{O(1)} / \log R$.

Next, if $q = p_1^{r_1} \cdots p_{A+1}^{r_{A+1}}$ is the prime factorisation of q , then (10.2) implies that

$$\left(\frac{\log(R/d)}{\log R} \right)^A M_{f_A}(aq; R/d) = \sum_{d'|aq} \mu(d') f_A \left(\frac{\log(dd')}{\log R} \right) = \sum_{d''|a} \mu(d'') w_q \left(\frac{\log(dd'')}{\log R} \right)$$

with

$$\text{with } w_q(x) := \sum_{J \subset [A+1]} (-1)^{|J|} f_A \left(x + \frac{\sum_{j \in J} \log p_j}{\log R} \right).$$

As usual, if $A = 0$, we may replace f_0 by a sufficiently smooth function h at the cost of a small error. Letting $g = h$ when $A = 0$ and $g = f_A$ otherwise, and letting W_q have the same definition as w_q with g in place of f_A , we find that

$$\begin{aligned} S''(d, q) &\leq \prod_{p \leq R} \left(1 - \frac{1}{p} \right) \sum_{\substack{P^+(a) \leq y \\ (a, q) = 1}} \frac{v^{\Omega(a)}}{a} \left(\sum_{f|a} \mu(f) W_q \left(\frac{\log(df)}{\log R} \right) \right)^{2k} + O \left(\frac{1}{\log R} \right) \\ &= \frac{\prod_{p \leq R} (1 - 1/p)}{\prod_{p \leq y, p \nmid q} (1 - v/p)} \sum_{\substack{(f_j, q) = 1 \\ 1 \leq j \leq 2k}} \frac{v^{\Omega([f_1, \dots, f_{2k}])}}{[f_1, \dots, f_{2k}]} \prod_{j=1}^{2k} \mu(f_j) W_q \left(\frac{\log(df_j)}{\log R} \right) + O \left(\frac{1}{\log R} \right). \end{aligned}$$

We apply Mellin inversion $2k$ times to find that

$$\begin{aligned} S''(d, q) &\leq \frac{\prod_{p \leq R} (1 - 1/p)}{(2\pi i)^{2k} \prod_{p \leq y, p \nmid q} (1 - v/p)} \int \cdots \int \sum_{\substack{\text{Re}(s_j) = 4k/\log y \\ |\text{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k}} \frac{v^{\Omega([f_1, \dots, f_{2k}])} \prod_{j=1}^{2k} \mu(f_j) f_j^{-s_j}}{[f_1, \dots, f_{2k}]} \\ &\quad \times \prod_{j=1}^{2k} \frac{\widehat{g}_R(s_j)}{d^{s_j}} \prod_{a=1}^{A+1} (1 - p_a^{-s_j}) ds_j + O \left(\frac{1}{\log R} \right). \end{aligned}$$

Together with (10.7), this implies that

$$\begin{aligned} S'(q) &\leq \frac{(\log \log R)^{O(1)} \prod_{p \leq R} (1 - 1/p)}{(2\pi i)^{2k} \prod_{p \leq y, p \nmid q} (1 - v/p)} \int \cdots \int \sum_{\substack{\text{Re}(s_j) = 4k/\log y \\ |\text{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k}} \frac{v^{\Omega([f_1, \dots, f_{2k}])} \prod_{j=1}^{2k} \mu(f_j) f_j^{-s_j}}{[f_1, \dots, f_{2k}]} \\ &\quad \times P(1 + s_{[2k]} - 2k/\log y) \prod_{j=1}^{2k} \widehat{g}_R(s_j) \prod_{a=1}^{A+1} (1 - p_a^{-s_j}) ds_j + O \left(\frac{1}{\log R} \right), \end{aligned}$$

where

$$P(s) := \sum_{d=1}^{\infty} \frac{(\lambda^+ * 1)(d)}{d^s} = \zeta(s) \sum_{m \leq R^\delta} \frac{\lambda^+(m)}{m^s}.$$

We fix $s_1, \dots, s_{2k-1}$ and move s_{2k} to the line $\text{Re}(s_{2k}) = -8k/\log y$ to pick up the pole at $s_{[2k]} = 2k/\log y$. If c is small enough in the definition of y , and the same is true for δ , the

complementary contours contribute $\ll 1/\log R$. There are no other poles, since the factors $1 - p_a^{-s_j}$ are annihilating the poles of $\widehat{g}_R(s_j)$. We conclude that

$$S'(q) \ll \frac{(\log \log R)^{O(1)}}{(\log R)^{2kA+2-v}} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=4k/\log y \\ |\operatorname{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k \\ s_{[2k]}=2k/\log y}} \left| \sum_{\substack{P^+(f_j) \leq y \\ 1 \leq j \leq 2k}} \frac{v^{\Omega([f_1, \dots, f_{2k}])} \prod_{j=1}^{2k} \mu(f_j) f_j^{-s_j}}{[f_1, \dots, f_{2k}]} \right| \\ \left(\prod_{j=1}^{2k} \frac{1}{|s_j|^{A+1}} \right) |ds_1 \cdots ds_{2k-1}| + \frac{1}{\log R}.$$

Recall the notation $\lambda_p(\mathbf{t})$ defined in (9.12), and combine the above with (10.6) to find that

$$S' \ll \frac{(\log \log R)^{O(1)}}{(\log R)^{2kA+2-v}} \int \cdots \int_{\substack{-T \leq t_j \leq T \ (1 \leq j < 2k) \\ t_{[2k]}=0}} \left(1 + \frac{v \cdot \lambda_p(\mathbf{t}) + (A+1) \sum_{j=1}^{2k} \cos(t_j \log p)}{p} \right) \\ \times \left(\prod_{j=1}^{2k} \frac{(\log(2 + |t_j|))^{2k(A+1)}}{(1 + |t_j|)^{A+1}} \right) dt_1 \cdots dt_{2k-1} + \frac{(\log \log R)^{O(1)}}{\log R},$$

where we used the fact that $s\zeta(1+s) \gg 1/\log(2+|t|)$ for $\operatorname{Re}(s) > 1$. Then, following the arguments of Section 9.5 (with $L = 1$), and recalling the notations $V(\mathcal{I})$ and $\mathcal{A}(V(\mathcal{I}))$, we find that

$$S' \ll \max_{\mathcal{I} \subset \mathcal{I}(2k)} (\log R)^{v(1+\mathcal{A}(V(\mathcal{I}))) - \dim(V(\mathcal{I})) + (A+1)U(\mathcal{I}) - 2kA - 1} (\log \log R)^{O(1)},$$

where

$$U(\mathcal{I}) := \#\{1 \leq j \leq 2k : L_{\{j\}} \in V(\mathcal{I})\}.$$

If $U(\mathcal{I}) \geq 1$, then Proposition 7.1(a) implies that $\mathcal{A}(V(\mathcal{I})) = -1$, and the exponent of $\log R$ is then

$$(A+1) \cdot U(\mathcal{I}) - \dim(V(\mathcal{I})) - 2kA - 1 \leq A \cdot U(\mathcal{I}) - 2kA - 1 \leq -1,$$

since we clearly have that $2k \geq \dim(V(\mathcal{I})) \geq U(\mathcal{I})$. Assume now that $U(\mathcal{I}) = 0$. If $\mathcal{A}(V(\mathcal{I})) = \binom{2k}{k} - 1$ and $\dim(V(\mathcal{I})) = 2k - 1$, then the exponent of $\log R$ is $v\binom{2k}{k} - 2k(A+1)$. Finally, if this is not the case, then Proposition 7.1 (together with the argument in the end of Section 9.5) implies that

$$\begin{aligned} v(1 + \mathcal{A}(V(\mathcal{I}))) - \dim(V(\mathcal{I})) - 1 &\leq \max\{0, v-1\} \binom{2k}{k} + \mathcal{A}(V(\mathcal{I})) - \dim(V(\mathcal{I})) \\ &\leq \max\{0, v-1\} \binom{2k}{k} + \binom{2k}{k} - 2k - 1 \\ &\leq v \binom{2k}{k} - 2k, \end{aligned}$$

by our assumption that $v \geq 1 - 1/\binom{2k}{k}$. This completes the proof of the theorem. $\square$

10.2. From logarithmic weights to uniform weights. In this section, we show how to go from Theorem 10.1 to the analogous result for the regular mean value and then prove Theorem 1.6.

Theorem 10.2. *Let $x \geq R \geq 2$, $k \in \mathbb{N}$, $A \in \mathbb{Z}_{\geq 0}$ and $\epsilon \in (0, 1/2)$.*

(a) *Assume that $A > \frac{1}{2k} \binom{2k}{k} - 1$. Then uniformly for $\eta \in [\log 2 / \log R, 1]$, we have*

$$\frac{1}{x} \sum_{n \leq x} \Omega(n; R^\eta) M_{f_A}(n; R)^{2k} \ll_{k,A} \frac{\eta}{\log R}.$$

(b) *If $A \leq \frac{1}{2k} \binom{2k}{k} - 1$ and $1 - 1/\binom{2k}{k} + \epsilon \cdot \mathbf{1}_{k=1} \leq v \leq 2 - \epsilon$, then*

$$\frac{1}{x} \sum_{n \leq x} v^{\Omega(n; R)} M_{f_A}(n; R)^{2k} \ll_{k,A,\epsilon} (\log R)^{v \binom{2k}{k} - 2k(A+1)} (\log \log R)^{O(1)}.$$

Proof. We start by proving a preparatory estimate. Our goal is to show that there is some constant $C = C(k)$ such that

$$(10.8) \quad \sum_{n \leq x} M_{f_A}(n; R)^{2k} \leq \frac{Cx}{\log R} \quad (x \geq R > 1).$$

When $x \leq 2^H$ for some fixed $H \in \mathbb{N}$ that will be taken large enough in terms of k and A , relation (10.8) trivially holds by taking C to be large enough in terms of J . Assume now that (10.8) holds for all $x \leq 2^h$, where $h \geq H$. We wish to prove that (10.8) is also true when $x \in (2^h, 2^{h+1}]$.

We follow a variation of the argument leading to Theorem III.3.5 in [22, p.308]: note that

$$(10.9) \quad \sum_{n \leq x} M_{f_A}(n; R)^{2k} = \sum_{n \leq x} M_{f_A}(n; R)^{2k} \frac{\log(x/n)}{\log x} + \sum_{n \leq x} M_{f_A}(n; R)^{2k} \frac{\log n}{\log x}.$$

For the first sum, we bound $\log(x/n)$ by x/n to give

$$(10.10) \quad \begin{aligned} \sum_{n \leq x} M_{f_A}(n; R)^{2k} \frac{\log(x/n)}{\log x} &\leq \sum_{n \leq x} M_{f_A}(n; R)^{2k} \frac{x}{n \log x} \\ &\ll \frac{x}{\log R} \sum_{P^+(n) \leq R} \frac{M_{f_A}(n; R)^{2k}}{n} \\ &\ll \frac{x}{\log R}, \end{aligned}$$

by Theorem 1.3. In the second sum, we write $\log n = \sum_{p^j \parallel n} j \log p$ to find that

$$\sum_{n \leq x} M_{f_A}(n; R)^{2k} \frac{\log n}{\log x} = \sum_{p^j \leq x} \frac{j \log p}{\log x} \sum_{\substack{m \leq x/p^j \\ p \nmid m}} M_{f_A}(mp^j; R)^{2k}.$$

Since

$$M_{f_A}(mp^j; R) = M_{f_A}(m; R) - \mathbf{1}_{p < R} \left(\frac{\log(R/p)}{\log R} \right)^A M_{f_A}(m; R/p)$$

by (10.2), Minkowski's inequality implies that

$$\sum_{n \leq x} M_{f_A}(n; R)^{2k} \frac{\log n}{\log x} \leq \left(S_1^{\frac{1}{2k}} + S_2^{\frac{1}{2k}} \right)^{2k},$$

where

$$S_1 := \sum_{p^j \leq x} \frac{j \log p}{\log x} \sum_{m \leq x/p^j} M_{f_A}(m; R)^{2k}.$$

and

$$S_2 := \sum_{\substack{p^j \leq x \\ p < R}} \frac{j \log p}{\log x} \left(\frac{\log(R/p)}{\log R} \right)^{2kA} \sum_{m \leq x/p^j} M_{f_A}(m; R/p)^{2k}.$$

For S_1 , we note that

$$S_1 = \sum_{m \leq x} M_{f_A}(m; R)^{2k} \sum_{p^j \leq x/m} \frac{j \log p}{\log x} \ll \frac{x}{\log x} \sum_{m \leq x} \frac{M_{f_A}(m; R)^{2k}}{m} \ll \frac{x}{\log R},$$

by (10.10). Finally, we need to bound S_2 . First, we bound its subsum with $j \geq 2$. We have that

$$\begin{aligned} & \sum_{\substack{p^j \leq x \\ j \geq 2 \\ p < R}} \frac{j \log p}{\log x} \left(\frac{\log(R/p)}{\log R} \right)^{2kA} \sum_{m \leq x/p^j} M_{f_A}(m; R/p)^{2k} \\ & \leq \sum_{\substack{p^j \leq x \\ j \geq 2 \\ p < R}} \frac{j \log p}{\log x} \left(\frac{\log(R/p)}{\log R} \right)^{2kA} \sum_{m \leq x/p^j} M_{f_A}(m; R/p)^{2k} \frac{x}{p^j m} \\ & \ll x \sum_{\substack{p^j \leq x \\ j \geq 2 \\ p < R}} \frac{j \log p}{p^j \log x} \left(\frac{\log(R/p)}{\log R} \right)^{2kA} \cdot \frac{\log x}{\log(R/p)}, \end{aligned}$$

by (10.10) with R replaced by R/p . Since $A \geq 1$ here, we find that the above is

$$\ll \sum_{\substack{p^j \leq x \\ j \geq 2}} \frac{j \log p}{p^j} \cdot \frac{x}{\log R} \ll \frac{x}{\log R},$$

where the implied constant is independent of C .

Finally, we need to bound the part of S_2 with $j = 1$. We note that $x/p \leq 2^h$ and that $R/p \leq x/p$, so we may apply the induction hypothesis. This gives a bound

$$\begin{aligned} & \leq \sum_{p < R} \frac{\log p}{\log x} \left(\frac{\log(R/p)}{\log R} \right)^{2kA} \cdot \frac{Cx}{p \log(R/p)} \leq \frac{Cx}{(\log x)(\log R)^2} \sum_{p < R} \frac{(\log p)(\log(R/p))}{p} \\ & \leq \frac{2C}{3} \cdot \frac{x}{\log R}, \end{aligned}$$

provided that J (and hence x) is big enough, where we used our assumptions that $A \geq 1$ and $x \geq R$. Combining the above, and assuming that C is big enough in terms of k and A completes the inductive step and hence the proof of (10.8).

(a) We now turn to proving part (a), that is bounding the sum

$$S(x, R, Q) := \sum_{n \leq x} \Omega(n; Q) M_{f_A}(n; R)^{2k}.$$

The proof is similar to the proof of (10.8), so we only give the main technical twists: we use induction on the dyadic interval in which x lies to prove that there is some constant $C' = C'(k, \epsilon)$ such that

$$(10.11) \quad S(x, R, Q) \leq \frac{C' x \log Q}{(\log R)^2} \quad (x \geq R \geq Q \geq 2).$$

When $x \leq 2^H$, for some fixed $H \in \mathbb{N}$ that will be taken large enough in terms of k , A and ϵ , this trivially holds by taking C' to be large enough in terms of H . Assume now that (10.8) holds for all $x \leq 2^h$, where $h \geq H$. We will prove that it also holds for $x \in (2^h, 2^{h+1}]$. Note that the analogues of (10.9) and (10.10) hold here as well, so let us focus on understanding the sum

$$T := \sum_{n \leq x} \Omega(n; Q) M_{f_A}(n; R)^{2k} \frac{\log n}{\log x}.$$

Fix a large integer N and call T_1 the portion of this sum with $\Omega(n; Q) > 2N$ and T_2 the remaining sum. Writing $\log n = \sum_{p^j \parallel n} j \log p$, we find that

$$T_1 = \sum_{p^j \leq x} \frac{j \log p}{\log x} \sum_{\substack{m \leq x/p^j, p \nmid m \\ \Omega(mp^j; Q) > 2N}} \Omega(mp^j; Q) M_{f_A}(mp^j; R)^{2k}.$$

If T'_1 is the part with $\Omega(mp^j; Q) \leq 2j$ and T''_1 is the rest, then

$$\begin{aligned} T'_1 &\leq \sum_{\substack{p^j \leq x \\ j > N}} \frac{2j^2 \log p}{\log x} \sum_{\substack{m \leq x/p^j \\ p \nmid m}} M_{f_A}(mp^j; R)^{2k} \frac{x}{p^j m} \\ &\ll x \sum_{\substack{p^j \leq x, p \leq e^{\sqrt{\log R}} \\ j > N}} \frac{j^2 \log p}{p^j \log R} \sum_{\substack{P^+(m) \leq R \\ p \nmid m}} \frac{M_{f_A}(mp; R)^{2k}}{m} + \frac{x(\log R)^{O(1)}}{e^{\sqrt{\log R}}} \\ &\ll_N x \sum_{\substack{p^j \leq x, p \leq e^{\sqrt{\log R}} \\ j > N}} \frac{j^2 \log p}{p^j \log R} \cdot \left\{ \left(\frac{\log p}{\log R} \right)^{2k} + (\log R)^{1 + \binom{2k}{k} - 2k(A+1)} \right\} \\ &\ll \frac{x}{(\log R)^{2k+1}} + x(\log R)^{\binom{2k}{k} - 2k(A+1)}, \end{aligned}$$

where the second to last bound follows from (10.5). Finally, in the range of T''_1 , we note that $\Omega(m; Q) \geq (\Omega(m; Q) + j)/2 > N$, so that $\Omega(mp^j; Q) \leq j(1 + \Omega(m; Q)) \leq (1 + 1/N) \cdot j \cdot \Omega(m; Q)$.

Therefore,

$$(10.12) \quad T_1 \leq \frac{N+1}{N} \sum_{p^j \leq x} \frac{j^2 \log p}{\log x} \sum_{\substack{m \leq x/p^j, p \nmid m \\ \Omega(mp^j; Q) > 2N}} \Omega(m; Q) M_{f_A}(mp^j; R)^{2k} \\ + O\left(\frac{x}{(\log R)^{2k+1}} + x(\log R)^{\binom{2k}{k} - 2k(A+1)}\right).$$

Next, we need to bound

$$T_2 = \sum_{\substack{n \leq x \\ \Omega(n; Q) \leq 2N}} \Omega(n; Q) M_{f_A}(n; R)^{2k} \frac{\log n}{\log x}.$$

If $Q > R^{1/2N^2}$, then we simply note that

$$T_2 \leq 2N \sum_{n \leq x} M_{f_A}(n; R)^{2k} \ll_N \frac{x}{\log R} \ll_N \frac{x \log Q}{(\log R)^2}.$$

by (10.8). Here the implied constant depends on N but does not depend on C' .

On the other hand, if $Q \leq R^{1/2N^2}$, then we have that $\sum_{p^j \parallel n, p \leq Q} j \log p \leq (\log x)/N$, so that

$$T_2 \leq \frac{S(x, R, Q)}{N} + \sum_{\substack{n \leq x \\ \Omega(n; Q) \leq 2N}} \Omega(n; Q) M_{f_A}(n; R)^{2k} \sum_{p^j \parallel n, p > Q} \frac{j \log p}{\log x} \\ \leq \frac{S(x, R, Q)}{N} + \sum_{\substack{p^j \leq x \\ p > Q}} \frac{j \log p}{\log x} \sum_{\substack{m \leq x/p^j, p \nmid m \\ \Omega(mp^j; Q) \leq 2N}} \Omega(m; Q) M_{f_A}(mp^j; R)^{2k}.$$

Combining the above inequality and (10.12), we deduce that

$$T \leq \frac{S(x, R, Q)}{N} + \frac{N+1}{N} \sum_{p^j \leq x} \frac{j^2 \log p}{\log x} \sum_{m \leq x/p^j, p \nmid m} \Omega(m; Q) M_{f_A}(mp^j; R)^{2k} + O_N\left(\frac{x \log Q}{(\log R)^2}\right).$$

We thus conclude that

$$S(x, R, Q) \leq \frac{N+1}{N-1} \sum_{p^j \leq x} \frac{j^2 \log p}{\log x} \sum_{m \leq x/p^j, p \nmid m} \Omega(m; Q) M_{f_A}(mp^j; R)^{2k} + O_N\left(\frac{x \log Q}{(\log R)^2}\right).$$

We can bound the sum on the right hand side in an entirely analogous way to the proof of (10.8). Choosing N sufficiently large, and then C' large enough in terms of N completes the inductive step and thus the proof of (10.11). This proves part (a) of the theorem.

(b) The proof of part (b) is, for the most part, similar to the proof of (10.8). An important detail is that, after applying the induction hypothesis, we use the fact that

$$(10.13) \quad \sum_{p < R} \frac{v \log p}{\log x} \left(\frac{\log(R/p)}{\log R}\right)^{2kA} \cdot \frac{(\log(R/p))^{v\binom{2k}{k} - 2k(A+1)} (\log \log(R/p))^D}{p} \\ \sim \frac{v}{v\binom{2k}{k} - 2k + 1} \cdot \frac{\log R}{\log x} \cdot (\log R)^{\binom{2k}{k} - 2k(A+1)} (\log \log R)^D$$

for any fixed $D \geq 0$, as $R \rightarrow \infty$. This is sufficient when $k \geq 2$, because

$$\frac{v}{v \binom{2k}{k} - 2k + 1} \leq \frac{1 - 1/\binom{2k}{k}}{\binom{2k}{k} - 2k} \leq \frac{1}{2}$$

in this case.

However, when $k = 1$, the situation is more tricky. First of all, we note that the above argument allows to establish the theorem for $x \geq R^{2v/(2v-1)}$. (Note that if $p < R$ and $x \geq R^{2v/(2v-1)}$, then we also have that $x/p \geq (R/p)^{2v/(2v-1)}$, so that the inductive hypothesis can be applied.) Finally, it remains to treat the case when $x \leq R^{2v/(2v-1)}$. We then observe that $\Omega(n; R^\delta, R) \ll_{\epsilon, \delta} 1$, for any fixed δ . It would thus suffice to prove that

$$(10.14) \quad \sum_{n \leq x} v^{\Omega(n; R^\delta)} \delta^{\Omega(n; R^\delta, R)} M_{f_0}(n; R)^2 \leq C'' x (\log R)^{2v-2} (\log \log R)^D \quad (x \geq R \geq 2),$$

for some appropriate constants $C'', D > 0$. Then, in place of (10.13), we note that

$$\begin{aligned} & \sum_{p < R} \frac{(v \cdot \mathbf{1}_{p \leq R^\delta} + \delta \cdot \mathbf{1}_{R^\delta < p \leq R}) \log p}{\log x} \left(\frac{\log(R/p)}{\log R} \right)^{2kA} \cdot \frac{(\log(R/p))^{v \binom{2k}{k} - 2k(A+1)} (\log \log(R/p))^D}{p} \\ & < \frac{1}{2} \cdot (\log R)^{\binom{2k}{k} - 2k(A+1)} (\log \log R)^D \end{aligned}$$

as $R \rightarrow \infty$, as long as δ is small enough. This allows us to complete the inductive step and establish (10.14), thus completing the proof of the theorem. $\square$

Given the above result, proving Theorem 1.6 is quite easy:

Proof of Theorem 1.6. (a) This is an immediate consequence of Theorem 10.2(a).

(b) We use Rankin's trick: Given a small $\alpha > 0$ and $1 \leq v \leq 3/2$, we have that

$$\sum_{\substack{n \leq x \\ \Omega(n; R) > \binom{2k}{k} (1+\alpha) \log \log R}} M_{f_A}(n; R)^{2k} \leq \sum_{n \leq x} v^{\Omega(n; R) - \binom{2k}{k} (1+\alpha) \log \log R} M_{f_A}(n; R)^{2k}.$$

Theorem 10.2(b) then implies that

$$\sum_{\substack{n \leq x \\ \Omega(n; R) > \binom{2k}{k} (1+\alpha) \log \log R}} M_f(n; R)^{2k} \ll (\log R)^{\binom{2k}{k} - 2k(A+1) + \binom{2k}{k} (v-1 - (1+\alpha) \log v)},$$

provided that v is close enough to 1. We optimize this by choosing $v = 1 + \alpha$, so that $1 - v + (1 + \alpha) \log v = \int_1^{1+\alpha} (\log t) dt > 0$.

Similarly, we have that

$$\sum_{\substack{n \leq x \\ \Omega(n; R) < \binom{2k}{k} (1-\alpha) \log \log R}} M_{f_A}(n; R)^{2k} \leq \sum_{n \leq x} v^{\Omega(n; R) - \binom{2k}{k} (1-\alpha) \log \log R} M_{f_A}(n; R)^{2k},$$

for any $1 - 1/\binom{2k}{k} + \epsilon \cdot \mathbf{1}_{k=1} v \leq 1$. Applying Theorem 10.2(b) and choosing $v = 1 - \alpha$ for small enough α completes the proof of the theorem. $\square$

10.3. Estimates for general weight functions. It is not so hard to go from estimates for the moments of $M_{f_A}(n; R)$ to the moments of $M_f(n; R)$ for a general weight function f . The following lemma provides the key link.

Lemma 10.3. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be supported in $(-\infty, 1]$. Assume further that $f \in C^A(\mathbb{R})$ and that all functions $f, f', \dots, f^{(A)}$ are uniformly bounded for some integer $A \geq 1$. Then*

$$M_f(n; R)^{2k} \ll_{A,k,f} \int_{\frac{\log 2}{\log R}}^1 u^{2k(A-1)} M_{f_{A-1}}(n; R^u)^{2k} du + \frac{1}{(\log R)^{2kA}}.$$

Proof. Since $f(x) = 0$ for $x > 1$ and $f \in C^A(\mathbb{R})$, we must have that $f^{(j)}(1) = 0$ for $j \leq A$. Taylor's theorem with the integral form of the remainder term implies that

$$f(x) = \int_1^x \frac{f^{(A)}(u)}{(A-1)!} (x-u)^{A-1} du = \frac{(-1)^A}{(A-1)!} \int_{[x,1]} f^{(A)}(u) (u-x)^{A-1} du,$$

for all x , since both sides vanish if $x > 1$. Therefore,

$$\begin{aligned} M_f(n; R) &= \frac{(-1)^A}{(A-1)!} \sum_{d|n} \mu(d) \int_{\frac{\log d}{\log R} \leq u \leq 1} f^{(A)}(u) \left(u - \frac{\log d}{\log R}\right)^{A-1} du \\ &= \frac{(-1)^A}{(A-1)!} \int_0^1 f^{(A)}(u) \sum_{\substack{d|n \\ d \leq R^u}} \mu(d) \left(u - \frac{\log d}{\log R}\right)^{A-1} du \\ &= \frac{(-1)^A}{(A-1)!} \int_{\frac{\log 2}{\log R}}^1 f^{(A)}(u) u^{A-1} M_{f_{A-1}}(n; R^u) du + O\left(\frac{1}{(\log R)^A}\right), \end{aligned}$$

by noting that $d = 1$ if $d \leq R^u < 2$. Hölder's inequality then completes the proof. $\square$

We use the above lemma to show the analogue of Theorem 10.2 for general weight functions f .

Theorem 10.4. *Let $k \in \mathbb{N}$, $x \geq R \geq 2$ and $f : \mathbb{R} \rightarrow \mathbb{R}$ be supported in $(-\infty, 1]$. Assume further that $f \in C^A(\mathbb{R})$ and that all functions $f, f', \dots, f^{(A)}$ are uniformly bounded for some integer $A \geq 1$, and fix some $\epsilon \in (0, 1)$.*

(a) *Let $A > \frac{1}{2k} \binom{2k}{k}$. Uniformly for $\eta \in [\log 2 / \log R, 1]$, we have*

$$\frac{1}{x} \sum_{n \leq x} \Omega(n; R^\eta) M_f(n; R)^{2k} \ll \frac{\eta}{\log R}.$$

(b) *If $A \leq \frac{1}{2k} \binom{2k}{k}$ and $1 - 1/\binom{2k}{k} + \epsilon \cdot \mathbf{1}_{k=1} \leq v \leq 2 - \epsilon$, then*

$$\frac{1}{x} \sum_{n \leq x} v^{\Omega(n; R)} M_f(n; R)^{2k} \ll (\log R)^{v \binom{2k}{k} - 2kA} (\log \log R)^{O(1)}.$$

All implied constants depend at most on k , f and ϵ .

Proof. (a) Lemma 10.3 implies that

$$\begin{aligned} \sum_{n \leq x} \Omega(n; R^\eta) M_f(n; R)^{2k} &\ll \int_{\frac{\log 2}{\log R}}^1 u^{2k(A-1)} \sum_{n \leq x} \Omega(n; R^\eta) M_{f_{A-1}}(n; R^u)^{2k} du \\ &\quad + O\left(\frac{x \log \log R}{(\log R)^{2kA}}\right), \end{aligned} \tag{10.15}$$

When $u \geq \eta$, Theorem 10.2(a) implies that

$$(10.16) \quad \int_{\eta}^1 u^{2k(A-1)} \sum_{n \leq x} \Omega(n; R^{\eta}) M_{f_{A-1}}(n; R^u)^{2k} du \ll \frac{\eta x}{\log R}.$$

Finally, we consider the integral over $u \in [\log 2 / \log R, \eta]$. Observe that

$$\sum_{n \leq x} \Omega(n; R^u) M_{f_{A-1}}(n; R^u)^{2k} \ll \frac{x}{u \log R},$$

by Theorem 10.2(a). If $x \leq R^{100}$, then we also have that $\Omega(n; R^u, R^{\eta}) \leq 100/u$ for each $n \leq x$, so that

$$\sum_{n \leq x} \Omega(n; R^u, R^{\eta}) M_{f_{A-1}}(n; R^u)^{2k} \ll \frac{x}{u^2 \log R}$$

by Theorem 10.2. We thus find that

$$\int_{\frac{\log 2}{\log R}}^{\eta} u^{2k(A-1)} \sum_{n \leq x} \Omega(n; R^{\eta}) M_{f_{A-1}}(n; R^u)^{2k} du \ll \int_{\frac{\log 2}{\log R}}^{\eta} \frac{u^{2k(A-1)-2} x}{\log R} du \ll \frac{\eta x}{\log R}.$$

Together with (10.15) and (10.16), this proves part (a) in the case when $x \leq R^{100}$.

Finally, let us consider the case when $x > R^{100}$. Then

$$\sum_{n \leq x} \Omega(n; R^u, R^{\eta}) M_{f_{A-1}}(n; R^u)^{2k} \leq \sum_{\substack{p^j \leq x \\ R^u < p \leq R^{\eta}}} j \sum_{m \leq x/p^j} M_{f_{A-1}}(m; R^u)^{2k}.$$

When $p^{j-1} \geq \sqrt{x}$, then we bound the inner sum trivially by $\ll (x/p^j)(\log R)^{O(1)}$, so that the total contribution of such summands is $\ll \sqrt{x}(\log R)^{O(1)}$. Finally, when $p^{j-1} \leq \sqrt{x}$, then we see that $x/p^j \geq \sqrt{x}/R \geq x^{0.499} \geq R^u$, so that the sum over m is $\ll (x/p^j)/\log(R^u)$, by Theorem 10.2(a). We thus conclude that

$$\sum_{n \leq x} \Omega(n; R^u, R^{\eta}) M_{f_{A-1}}(n; R^u)^{2k} \ll \frac{x \log(\eta/u)}{u \log R} + \frac{x}{(u \log R)^2} \ll \frac{x}{u \log R}.$$

Therefore,

$$\int_{\frac{\log 2}{\log R}}^{\eta} u^{2k(A-1)} \sum_{n \leq x} \Omega(n; R^{\eta}) M_{f_{A-1}}(n; R^u)^{2k} du \ll \frac{\eta x}{\log R}.$$

in this case as well, thus completing the proof of part (a) of the theorem.

(b) This proof of this part is similar. We start again with Lemma 10.3 to find that Lemma 10.3 implies that

$$(10.17) \quad \sum_{n \leq x} v^{\Omega(n; R)} M_f(n; R)^{2k} \ll \int_{\frac{\log 2}{\log R}}^1 u^{2k(A-1)} \sum_{n \leq x} v^{\Omega(n; R)} M_{f_{A-1}}(n; R^u)^{2k} du + O(x(\log R)^{v-1-2kA}),$$

where we also used Theorem III.3.5 in [22, p.308]. Next, if $w = \max\{v, 1\}$ and $\log 2 / \log R \leq u \leq 1$, then note that

$$\sum_{n \leq x} v^{\Omega(n; R)} M_{f_{A-1}}(n; R^u)^{2k} \leq \sum_{\substack{a \leq x \\ P^+(a) \leq R^u}} v^{\Omega(a; R^u)} M_{f_{A-1}}(a; R^u)^{2k} \sum_{\substack{b \leq x/a \\ P^-(b) > R^u}} w^{\Omega(b; R^u, R)}.$$

When $x/a \geq R^u$, the sum over b is $\ll u^{1-w}x/(a \log(R^u))$ by Theorem III.3.5 in [22, p.308]. Hence,

$$\begin{aligned} \sum_{n \leq x} v^{\Omega(n;R)} M_{f_{A-1}}(n; R^u)^{2k} &\ll \frac{u^{1-w}x}{\log(R^u)} \sum_{\substack{a \leq x \\ P^+(a) \leq R^u}} \frac{v^{\Omega(a;R^u)} M_{f_{A-1}}(a; R^u)^{2k}}{a} \\ &+ \sum_{\substack{a \leq x \\ P^+(b) \leq R^u}} v^{\Omega(a;R^u)} M_{f_{A-1}}(a; R^u)^{2k}. \end{aligned}$$

We bound the first sum by Theorem 10.1(b) and the second one by Theorem 10.2(b) to find that

$$\sum_{n \leq x} v^{\Omega(n;R)} M_{f_{A-1}}(n; R^u)^{2k} \ll xu^{1-w+v\binom{2k}{k}-2kA} (\log R)^{v\binom{2k}{k}-2kA}.$$

Inserting the above bound into (10.17) completes the proof of the theorem. $\square$

We conclude this section with the proof of Theorem 1.4. We need a preliminary lemma.

Lemma 10.5. *If $m \in \mathbb{N}$, $g \in C^1(\mathbb{R}^m)$ and $z \geq y \geq 3$, then there is a positive constant $c > 0$ such that*

$$\begin{aligned} \sum_{y < p_1 < \dots < p_m \leq z} \frac{g(\log p_1, \dots, \log p_m)}{(p_1 - 1) \dots (p_m - 1)} &= \frac{1}{m!} \int_{[\log y, \log z]^m} \frac{g(t_1, \dots, t_m)}{t_1 \dots t_m} dt_1 \dots dt_m \\ &+ O\left(\frac{\|g\|_\infty + \|\nabla g\|_\infty}{e^{c\sqrt{\log y}}} \cdot \frac{(\sum_{y < p \leq z} 1/p)^{m-1} + (\int_y^z dt/t \log t)^{m-1}}{m!/m^2} \right). \end{aligned}$$

Proof. First of all, note that

$$\begin{aligned} \sum_{y < p_1 < \dots < p_m \leq z} \frac{g(\log p_1, \dots, \log p_m)}{(p_1 - 1) \dots (p_m - 1)} &= \frac{1}{m!} \sum_{\substack{y < p_1, \dots, p_m \leq z \\ \text{distinct}}} \frac{g(\log p_1, \dots, \log p_m)}{(p_1 - 1) \dots (p_m - 1)} \\ &= \frac{1}{m!} \sum_{y < p_1, \dots, p_m \leq z} \frac{g(\log p_1, \dots, \log p_m)}{p_1 \dots p_m} \\ &+ O\left(\frac{m^2 \|g\|_\infty (\sum_{y < p \leq z} 1/p)^{m-1}}{m!y} \right). \end{aligned}$$

So, if we can show that

$$\begin{aligned} \sum_{y < p_1, \dots, p_m \leq z} \frac{g(\log p_1, \dots, \log p_m)}{p_1 \dots p_m} &= \int_{[\log y, \log z]^m} \frac{g(t_1, \dots, t_m)}{t_1 \dots t_m} dt_1 \dots dt_m \\ &+ O\left(\sum_{j=1}^m \frac{\|g\|_\infty + \|\partial g / \partial x_j\|_\infty}{e^{c\sqrt{\log y}}} \left(\int_y^z \frac{dt}{t \log t} \right)^{j-1} \left(\sum_{y < p \leq z} \frac{1}{p} \right)^{m-j} \right), \end{aligned}$$

then the lemma will follow. But this estimate can be easily proved by induction on m and the Prime Number Theorem, and the proof is completed. $\square$

Let us now see how we can deduce Theorem 1.4 from the above results:

Proof of Theorem 1.4. The first estimate of part (a) follows by 10.4(a) and part (b) follows by Theorem 10.2(b). It remains to prove the second estimate of part (a). Note that it suffices to prove that

$$\prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{M_f(n; R)^{2k}}{n} = \frac{c_{k,f}}{\log R} + O\left(\frac{1}{(\log R)^{2-\epsilon}}\right),$$

since $x \geq R^{2k} \log^2 R$ here. Fix $\eta \in [\log 2 / \log R, 1]$ to be chosen later. Then Theorem 10.4(a) implies that

$$\sum_{\substack{P^+(n) \leq R \\ P^-(n) \leq R^\eta}} \frac{M_f(n; R)^{2k}}{n} \leq \sum_{P^+(n) \leq R} \frac{\Omega(n; R^\eta) M_f(n; R)^{2k}}{n} \ll \eta.$$

Moreover, for each positive integer m , we have that

$$\sum_{\substack{p|n \Rightarrow R^\eta < p \leq R \\ \omega(n)=m}} \frac{M_f(n; R)^{2k}}{n} = \sum_{R^\eta < p_1 < \dots < p_m \leq R} \frac{M_f(p_1 \cdots p_m; R)}{(p_1 - 1) \cdots (p_m - 1)}$$

We note that

$$M_f(p_1 \cdots p_m; R) = g_m \left(\frac{\log p_1}{\log R}, \dots, \frac{\log p_m}{\log R} \right),$$

where

$$g_m(t_1, \dots, t_m) := \sum_{J \subset \{1, \dots, m\}} (-1)^{|J|} f \left(\sum_{j \in J} t_j \right)$$

which is a smooth function satisfying the estimates $\|g_m^{2k}\|_\infty \leq 4^{km} \|f\|_\infty^{2k}$ and $\|\nabla g_m^{2k}\|_\infty \leq 2k 4^{km} \|f'\|_\infty \|f\|_\infty^{2k-1}$. Therefore Lemma 10.5 implies that

$$\sum_{\substack{p|n \Rightarrow R^\eta < p \leq R \\ \omega(n)=m}} \frac{M_f(n; R)^{2k}}{n} = \frac{1}{m!} \int_{[\eta, 1]^m} \frac{g_m(t_1, \dots, t_m)^{2k}}{t_1 \cdots t_m} dt_1 \cdots dt_m + O\left(\frac{(4^k \log(1/\eta) + O(1))^m}{e^{c\sqrt{\log(R^\eta)}} m! / m^2}\right)$$

for all $m \in \mathbb{N}$. We thus conclude that

$$(10.18) \quad \sum_{P^+(n) \leq R} \frac{M(n; R)^{2k}}{n} = F(\eta) + O\left(\eta + \frac{\log^2(1/\eta)}{\eta^{4^k} e^{c\sqrt{\log(R^\eta)}}}\right) \quad (0 < \eta \leq 1, R^\eta \geq 2),$$

where

$$F(\eta) := 1 + \sum_{m=1}^{\infty} \frac{1}{m!} \int_{[\eta, 1]^m} \frac{g_m(t_1, \dots, t_m)^{2k}}{t_1 \cdots t_m} dt_1 \cdots dt_m.$$

Completing the proof is now an exercise in real analysis. We start by proving that $\lim_{\eta \rightarrow 0^+} F(\eta)$ exists. Indeed, applying (10.18) twice, we deduce that

$$F(\eta_1) - F(\eta_2) \ll \max_{j \in \{1, 2\}} \left\{ \eta_j + \frac{\log^2(1/\eta_j)}{\eta_j^{4^k} e^{c\sqrt{\log(R^{\eta_j})}}} \right\}$$

whenever $0 < \eta_1 \leq \eta_2 \leq 1$ and $R^{\eta_1} \geq 2$. Letting $R \rightarrow \infty$, we find that

$$(10.19) \quad F(\eta_1) - F(\eta_2) \ll \eta_2 \quad (0 < \eta_1 \leq \eta_2 \leq 1).$$

In particular, Cauchy's convergence criterion implies that $\lim_{\eta \rightarrow 0^+} F(\eta)$ exists. Call F this limit, which clearly equals $F(0)$, and note that letting $\eta_1 \rightarrow 0^+$ in (10.19) implies that

$$F(\eta) = F + O(\eta) \quad (0 < \eta \leq 1).$$

Together with (10.18), this yields the estimate

$$\sum_{P^+(n) \leq R} \frac{M(n; R)^{2k}}{n} = F + O\left(\eta + \frac{\log^2(1/\eta)}{\eta^{4k} e^{c\sqrt{\log(R^\eta)}}}\right) \quad (0 < \eta \leq 1, R^\eta \geq 2).$$

Selecting η such that

$$R^\eta = e^{(\log \log R)^3}$$

completes the proof of Theorem 1.4. $\square$

We conclude this section with the proof of Theorem 1.7.

Proof of Theorem 1.7. The first estimate of part (a) can be proven following *mutatis mutandis* the proof of Theorem 1.6(a) above, using Theorem 10.4 in place of Theorem 10.2. Similarly, part (b) follows from the proof of Theorem 1.6(b). $\square$

11. THE ANALOGY FOR NON-EXCEPTIONAL DIRICHLET CHARACTERS

In the section, we study the sum

$$\mathcal{X}_{2k}(R) = \prod_{p \leq R} \left(1 - \frac{1}{p}\right) \sum_{P^+(n) \leq R} \frac{1}{n} \left(\sum_{\substack{d|n \\ R/2 < d \leq R}} \chi(d) \right)^{2k}$$

when $k \geq 2$ and $L(1, \chi)$ is not very small and prove Theorem 1.5(b).

We have that

$$\mathcal{X}_{2k}(R) = \sum_{R/2 < d_1, \dots, d_{2k} \leq R} \frac{\chi(d_1)\chi(d_2) \cdots \chi(d_{2k})}{[d_1, \dots, d_{2k}]}.$$

We want to introduce new variables D_I , $I \in \mathcal{S}^*(2k)$, as in Section 2, but first we perform a technical manoeuvre to simplify the situation. We write $d_i = d'_i d''_i$, where $P^+(d'_i) \leq y < P^-(d''_i)$, where

$$y := (\log R)^{4k+1}.$$

The contribution to $\mathcal{X}_{2k}(R)$ of $\mathbf{d}$'s for which d''_i is not square-free for some i is $\ll (\log R)^{4k-1}/y$ by a crude upper bound, and so is the contribution of those $\mathbf{d}$'s with $\max_i d'_i > B$, where

$$B := e^{(\log \log R)^3},$$

by Rankin's trick. Then we let D_I , $I \in \mathcal{S}^*(2k)$, be the product of those primes that divide d''_i when $i \in I$, and are coprime to the other d''_i 's. The numbers D_I are pairwise coprime and square-free, and $d''_i = \prod_{I \in \mathcal{S}^*(2k), I \ni i} D_I$, so that

$$\frac{\chi(d''_1)\chi(d''_2) \cdots \chi(d''_{2k})}{[d''_1, \dots, d''_{2k}]} = \frac{\prod_{I \in \mathcal{S}^+(2k)} \chi_0(D_I) \prod_{I \in \mathcal{S}^-(2k)} \chi(D_I)}{\prod_{I \in \mathcal{S}^*(2k)} D_I}.$$

We may now drop the condition that the D_I 's are square-free and coprime, since the contribution to $\mathcal{I}_{2k}(R)$ of the D_I 's not satisfying these conditions is $\ll (\log R)^{4k-1}/y$. Finally, we may drop the condition that $(D_I, q) = 1$ for $I \in \mathcal{S}^+(2k)$, encoded in the notation $\chi_0(D_I)$, since

the contribution of D_I 's with $P^-(D_I) > y$ and $(D_I, q) > 1$ is $\ll (\log R)^{4^k-1} \sum_{p|q, p>y} 1/p \ll (\log R)^{4^k}/y$. The above discussion implies that

$$\begin{aligned} \mathcal{X}_{2k}(R) = & \sum_{\substack{P^+(d'_i) \leq y, d'_i \leq B \\ 1 \leq i \leq 2k}} \frac{\chi(d'_1) \cdots \chi(d'_{2k})}{[d'_1, \dots, d'_{2k}]} \sum_{\substack{P^-(D_I) > y \ (I \in \mathcal{S}^*(2k)) \\ R/(2d'_i) < \prod_{I \in \mathcal{S}^*(2k), I \ni i} D_I \leq R/d'_i \\ 1 \leq i \leq 2k}} \frac{\prod_{I \in \mathcal{S}^-(2k)} \chi(D_I)}{\prod_{I \in \mathcal{S}^*(2k)} D_I} \\ & + O\left(\frac{1}{\log R}\right). \end{aligned}$$

Next, we note that

$$\sum_{\substack{n \leq x \\ P^-(n) > y}} \chi(n) \ll x^{1-1/(30 \log y)} \quad (x \geq \max\{q^4, y\}),$$

by Lemma 2.4 in [14]. Therefore,

$$(11.1) \quad \sum_{\substack{n > q^4 B \\ P^-(n) > y}} \frac{\chi(n)}{n} \ll \frac{1}{y}.$$

This implies that the contribution to $\mathcal{X}_{2k}(R)$ with $D_I > q^4 B$ for some $I \in \mathcal{S}^-(2k)$ is $\ll (\log R)^{4^k-1}/y$. To conclude, we have shown

$$\begin{aligned} \mathcal{X}_{2k}(R) = & \sum_{\substack{P^+(d'_i) \leq y, d'_i \leq B \\ 1 \leq i \leq 2k}} \sum_{\substack{D_I \leq q^4 B \\ P^-(D_I) > y \\ I \in \mathcal{S}^-(2k)}} \frac{\chi(d'_1) \cdots \chi(d'_{2k})}{[d'_1, \dots, d'_{2k}]} \cdot \frac{\prod_{I \in \mathcal{S}^-(2k)} \chi(D_I)}{\prod_{I \in \mathcal{S}^-(2k)} D_I} \cdot T(R_1, \dots, R_{2k}) \\ (11.2) \quad & + O\left(\frac{1}{\log R}\right). \end{aligned}$$

where $R_i = R/(d'_i \prod_{I \in \mathcal{S}^-(2k), I \ni i} D_I)$ and

$$T(\mathbf{R}) := \sum_{\substack{P^-(D_I) > y \ (I \in \mathcal{S}^+(2k)) \\ R_i/2 < \prod_{I \in \mathcal{S}^+(2k), I \ni i} D_I \leq R_i \\ 1 \leq i \leq 2k}} \frac{1}{\prod_{I \in \mathcal{S}^*(2k)} D_I}.$$

Our task now becomes estimating $T(\mathbf{R})$. Let $d = 2^{2k-1} - 2k$ and recall the definition of $V_d(\cdot)$ from the statement of Theorem 1.5(b). The proof of Theorem 1.1 shows that $V_k(m) \asymp m^d$. We claim that

$$(11.3) \quad T(\mathbf{R}) = V_k(\log R) \left(1 + O\left(\frac{\log B}{\log R}\right)\right) \prod_{p \leq y} \left(1 - \frac{1}{p}\right)^{2^{2k-1}-1}$$

whenever $R/B \leq R_i \leq R$ for all i , as is the case here. Proving (11.3) can be accomplished easily using a lattice point count and the fundamental lemma of sieve methods. First of all, note that the part of $T(\mathbf{R})$ where $D_I \leq B$ for some $I \in \mathcal{S}^+(2k)$ is trivially $\ll (\varphi(P)/P)^{2^{2k-1}-1} (\log R)^{d-1} \log B$ by an upper bound sieve, where we have set $P := \prod_{p \leq y} p$ for simplicity. In the rest of the range, we set $\rho = 1 + 1/\log R$ and divide the variables D_I into boxes of the form $(\rho^{m_I}, \rho^{m_I+1}]$, $m_I \geq 0$. Replacing D_I by ρ^{m_I} in the conditions

$R_i/2 < \prod_{J \in \mathcal{S}^+(2k), J \ni i} D_J \leq R_i$ creates a total error of size $\ll (\varphi(P)/P)^{2^{2k-1}-1} (\log R)^{d-1}$. In addition, if $\rho^{m_I} \geq B = e^{(\log \log R)^3}$, then we have that

$$\sum_{\substack{\rho^{m_I} < D_I \leq \rho^{m_I+1} \\ P^-(D_I) > y}} \frac{1}{D_I} = (\log \rho) \frac{\varphi(P)}{P} + O_C \left(\frac{1}{(\log R)^C} \right)$$

for any fixed C , by the fundamental lemma of sieve methods (see, for example, [22, Theorem I.4.3]). We thus conclude that

$$\begin{aligned} T(\mathbf{R}) &= ((\log \rho) \varphi(P)/P)^{2^{2k-1}-1} \sum_{\substack{m_I \geq \log B / \log \rho \quad (I \in \mathcal{S}^+(2k)) \\ \frac{\log(R_i/2)}{\log \rho} < \sum_{I \in \mathcal{S}^+(2k), I \ni i} m_I \leq \frac{\log R_i}{\log \rho}}} 1 \\ &\quad + O \left((\varphi(P)/P)^{2^{2k-1}-1} (\log R)^{d-1} \log B \right). \end{aligned}$$

A straightforward lattice point counting argument implies that the sum on the right hand side equals

$$\frac{W_k(\log R_1, \dots, \log R_{2k})}{(\log \rho)^{2^{2k-1}-1}} \left(1 + O \left(\frac{\log B}{\log R} \right) \right),$$

where $W_k(\mathbf{m})$ is the volume of the polytope $\{(x_I)_{I \in \mathcal{S}^+(2k)} : x_I \geq 0 \forall I, m_i - \log 2 \leq \sum_{I \ni i} x_I \leq m_i \forall i\}$. Since $m_i = \log R_i = \log R + O(\log B)$ here, we may show using the Mean Value Theorem that $W_k(\mathbf{m}) = V_k(m) + O((\log R)^{d-1} \log B)$. Relation (11.3) then follows.

We are now ready to complete the proof of Theorem 1.5(a): inserting the estimate (11.3) into (11.2), we conclude that

$$\begin{aligned} \mathcal{X}_{2k}(R) &= \frac{V_k(\log R)}{(P/\varphi(P))^{2^{2k-1}-1}} \sum_{\substack{P^+(d'_i) \leq y, d_i \leq B \\ 1 \leq i \leq 2k}} \sum_{\substack{D_I \leq q^4 B \\ P^-(D_I) > y \\ I \in \mathcal{S}^-(2k)}} \frac{\chi(d'_1) \cdots \chi(d'_{2k})}{[d'_1, \dots, d'_{2k}]} \cdot \frac{\prod_{I \in \mathcal{S}^-(2k)} \chi(D_I)}{\prod_{I \in \mathcal{S}^-(2k)} D_I} \\ &\quad + O((\log R)^{d-1} (\log(q \log R))^{O(1)}). \end{aligned}$$

We now remove the conditions $D_I \leq q^4 B$ and $d'_i \leq B$ via (11.1) and Rankin's trick, respectively. We conclude that

$$\begin{aligned} \mathcal{X}_{2k}(R) &= \frac{V_k(\log R)}{(P/\varphi(P))^{2^{2k-1}-1}} \sum_{\substack{P^+(d'_i) \leq y \\ 1 \leq i \leq 2k}} \sum_{\substack{P^-(D_I) > y \\ I \in \mathcal{S}^-(2k)}} \frac{\chi(d'_1) \cdots \chi(d'_{2k})}{[d'_1, \dots, d'_{2k}]} \cdot \frac{\prod_{I \in \mathcal{S}^-(2k)} \chi(D_I)}{\prod_{I \in \mathcal{S}^-(2k)} D_I} \\ &\quad + O((\log R)^{d-1} (\log(q \log R))^{O(1)}). \end{aligned}$$

Finally, we note that

$$\sum_{\substack{P^+(d'_i) \leq y \\ 1 \leq i \leq 2k}} \frac{\chi(d'_1) \cdots \chi(d'_{2k})}{[d'_1, \dots, d'_{2k}]} = \prod_{p \leq y} \left(1 + \sum_{j=1}^{\infty} \sum_{\substack{j_1, \dots, j_{2k} \geq 0 \\ \max\{j_1, \dots, j_{2k}\} = j}} \frac{\chi(p)^{j_1 + \dots + j_{2k}}}{p^j} \right).$$

The coefficient of $1/p$ is $(2^{2k-1} - 1)\chi_0(p) + 2^{2k-1}\chi(p)$. We thus conclude that

$$\begin{aligned}
& (\varphi(P)/P)^{2^{2k-1}-1} \sum_{\substack{P^+(d'_i) \leq y \\ 1 \leq i \leq 2k}} \sum_{\substack{P^-(D_I) > y \\ I \in S^-(2k)}} \frac{\chi(d'_1) \cdots \chi(d'_{2k})}{[d'_1, \dots, d'_{2k}]} \cdot \frac{\prod_{I \in S^-(2k)} \chi(D_I)}{\prod_{I \in S^-(2k)} D_I} \\
&= \prod_{p \leq y} \left(1 + \sum_{j=1}^{\infty} \sum_{\substack{j_1, \dots, j_{2k} \geq 0 \\ \max\{j_1, \dots, j_{2k}\} = j}} \frac{\chi(p)^{j_1 + \dots + j_{2k}}}{p^j} \right) \left(1 - \frac{1}{p} \right)^{2^{2k-1}-1} \prod_{p > y} \left(1 - \frac{\chi(p)}{p} \right)^{-2^{2k-1}} \\
&= \prod_p \left(1 + \sum_{j=1}^{\infty} \sum_{\substack{j_1, \dots, j_{2k} \geq 0 \\ \max\{j_1, \dots, j_{2k}\} = j}} \frac{\chi(p)^{j_1 + \dots + j_{2k}}}{p^j} \right) \left(1 - \frac{1}{p} \right)^{2^{2k-1}-1} + O\left(\frac{1}{y}\right).
\end{aligned}$$

An easy calculation then completes the proof of Theorem 1.5.

12. THE ANALOGY FOR EXCEPTIONAL DIRICHLET CHARACTERS

In this section, we consider the quantity $\mathcal{X}_{2k}(R)$ when $k = 1$, or when $\chi(p) = -1$ for most primes $p \leq R$. Our arguments here resemble closely the ones of Section 8.

12.1. Initial preparations. Arguing as in Section 8.7, we find that

$$\begin{aligned}
\mathcal{X}_{2k}(R) &= \frac{1}{(2i\pi)^{2k}} \int \cdots \int_{\substack{\operatorname{Re}(s_j) = \lambda^j / \log R \\ |\operatorname{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k}} \sum_{d_1, \dots, d_{2k} \geq 1} \frac{\prod_{j=1}^{2k} \chi(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \prod_{j=1}^{2k} \widehat{h}_R(s_j) (1 - 2^{-s_j}) ds_1 \cdots ds_{2k} \\
&\quad + O\left(\frac{1}{(\log R)^{100}}\right),
\end{aligned}$$

where h is a smooth function with $h(x) = 1$ for $x \leq 1 - 1/(\log R)^{(2k-1)2^{2k+1}+200k+2}$ and $h(x) = 0$ for $x > 1$, $T = \exp\{(\log \log R)^2\}$, and λ is some large parameter > 1 to be chosen later.

By expanding as an Euler product, we find that for $\operatorname{Re}(s_1), \dots, \operatorname{Re}(s_{2k}) \geq -1/4^k$ we have

$$\begin{aligned}
\sum_{d_1, \dots, d_{2k} \geq 1} \frac{\prod_{j=1}^{2k} \chi(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} &= \prod_p \left(1 + \sum_{I \in S^*(2k)} \frac{\chi(p)^{|I|}}{p^{1+s_I}} + O\left(\frac{1}{p^{2-2k/4^k}}\right) \right) \\
&= P(\mathbf{s}) \prod_{I \in S^*(2k)} L(1 + s_I, \chi^{|I|}),
\end{aligned}$$

where $P(\mathbf{s})$ is given by an Euler product which converges absolutely in the region $\operatorname{Re}(s_j) > -1/4^k$ for all j . Next, we set

$$F(\mathbf{s}) = P(\mathbf{s}) \prod_{j=1}^{2k} \frac{\widehat{h}_R(s_j)(1 - 2^{-s_j})}{R^{s_j}}$$

and

$$\zeta_q(s) := L(s, \chi_0) = \zeta(s) \prod_{p|q} \left(1 - \frac{1}{p^s}\right),$$

so that

$$(12.1) \quad \begin{aligned} \mathcal{X}_{2k}(R) &= \frac{1}{(2i\pi)^{2k}} \int \cdots \int_{\substack{\operatorname{Re}(s_j) = \lambda^j / \log R \\ |\operatorname{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k}} F(\mathbf{s}) R^{s_{[2k]}} \prod_{I \in \mathcal{S}^-(2k)} L(1 + s_I, \chi) \\ &\quad \times \prod_{I \in \mathcal{S}^+(2k)} \zeta_q(1 + s_I) ds_1 \cdots ds_{2k} + O\left(\frac{1}{(\log R)^{100}}\right). \end{aligned}$$

Note that F is analytic when $\operatorname{Re}(s_j) \geq -1/2^{2k+2}$ for all j , and that its derivatives satisfy

$$(12.2) \quad \frac{\partial^{j_1+\cdots+j_{2k}} F}{\partial s_1^{j_1} \cdots \partial s_{2k}^{j_{2k}}}(\mathbf{s}) \ll_{j_1, \dots, j_{2k}} \frac{1}{(|s_1| + 1) \cdots (|s_{2k}| + 1)}$$

in this region by (6.5). In particular, F is in the class of functions $\mathcal{C}_{2k}$, as defined in Section 8 after relation (8.7). Moreover, we know that there is an absolute constant $c_1 > 0$ such that

$$(12.3) \quad L^{(j)}(s, \psi) \ll_m (1 + (q + |t|)^{c_1(1-\sigma)}) \log^{j+1}(q + |t|) + \frac{\mathbf{1}_{\psi=\chi_0}}{|s-1|^{j+1}}$$

for $j \in \mathbb{Z}_{\geq 0}$, $\psi \in \{\chi, \chi_0\}$ and $j \in \{0, 1\}$, a standard consequence of bounds on the exponential sum $\sum_{n \leq N, n \equiv a \pmod{q}} n^{it}$ (see, for example, Lemma 4.1 in [14]). Throughout the proof, we assume that $R \geq q^{2c_1}$, the complimentary case being trivial. We follow closely the argument used in Section 8, so we only highlight the key points here.

12.2. The case $k = 1$. We first deal with the case $k = 1$ that is easy and will help us clarify some of the technical details of the argument. When $k = 1$, we move the variable s_2 to the line $\operatorname{Re}(s_2) = -\epsilon$, for a sufficiently small ϵ . The contribution of the vertical contours is $\ll (\log R)^{O(1)}/T$, and the contribution of the contour $\operatorname{Re}(s_2) = -\delta$ is $\ll R^{-\delta}$, for some positive $\delta = \delta(\epsilon)$ by relations (12.2) and (12.3), and our assumption that $R \geq q^{2c_1}$. In conclusion,

$$\mathcal{X}_2(R) = \frac{1}{2i\pi} \int_{\substack{\operatorname{Re}(s_1) = \lambda / \log R \\ |\operatorname{Im}(s_1)| \leq T \\ 1 \leq j \leq 2k}} F(s_1, -s_1) L(1 + s_1, \chi) L(1 - s_1, \chi) ds_1 + O\left(\frac{1}{(\log R)^{100}}\right).$$

Finally, we move s_1 to the line $\operatorname{Re}(s_1) = 0$. No poles are encountered, and the contribution of the vertical lines is easily seen to be $\ll (\log R)^{O(1)}/T$, so that

$$\begin{aligned} \mathcal{X}_2(R) &= \frac{1}{2\pi} \int_{-T}^T F(it, -it) |L(1 + it, \chi)|^2 dt + O\left(\frac{1}{(\log R)^{100}}\right) \\ &= \frac{1}{2\pi} \int_{-(\log R)^{102}}^{(\log R)^{102}} P(it, -it) \left| L(1 + it, \chi) \widehat{h}_R(it) (1 - 2^{it}) \right|^2 dt + O\left(\frac{1}{(\log R)^{100}}\right), \end{aligned}$$

since $\widehat{h}_R(it)(1 - 2^{it}) \ll 1/(1 + |t|)$. Finally, using 6.6 to replace $\widehat{h}_R(s)$ by R^s/s , choosing C to be large enough, and then extending the range of integration to $\mathbb{R}$ yields the estimate

$$\mathcal{X}_2(R) = \frac{1}{2\pi} \int_{-\infty}^{\infty} P(it, -it) \left| L(1 + it, \chi) \cdot \frac{\sin(t(\log 2)/2)}{t} \right|^2 dt + O\left(\frac{1}{(\log R)^{100}}\right),$$

which proves Theorem 1.5(a). (Obviously, we can obtain a much stronger error term, but we have chosen to content ourselves with a more qualitative result.)

12.3. Contour shifting. Next, we focus on the case $k \geq 2$. From now on, we will always be working under the assumptions and notations

$$L(\beta, \chi) = 0, \quad \beta > 1 - 1/(100 \log q), \quad Q = e^{1/(1-\beta)}.$$

Since $\sum_{q < p \leq Q} (1 + \chi(p))/p \ll 1$ and $\sum_{p > Q} \chi(p)/p \ll 1$, we note once and for all

$$(12.4) \quad L(1, \chi) \asymp \frac{1}{\log Q} \prod_{p \leq q} \left(1 + \frac{1 + \chi(p)}{p}\right).$$

As in Section 8, we shift the contours of the variables $s_1, \dots, s_{2k}$ in a certain order. Assume we have moved N variables, say $s_{j_1}, s_{j_2}, \dots, s_{j_N}$, and that we have picked up poles coming from the linear equations $s_{I_n} = 0$, $1 \leq n \leq N$ (which will be linearly independent by construction). We then set

$$V_N = \text{Span}_{\mathbb{Q}}(s_{I_1}, \dots, s_{I_N}) \quad \text{and} \quad \mathcal{I}_N = \{I \in \mathcal{S}(2k) : s_I \in V_N\}.$$

Each time we shift contours, the contribution of the new contours to $\mathcal{X}_k(R)$ can be shown to be $\ll 1/\log R$ by (12.2) and (12.3). So, we only need to understand the contribution of the poles after the contour shifting.

Let $L_{N,I}$ be the linear form $\sum_{i \in I} s_i$ in the variables $\{s_1, \dots, s_{2k}\} \setminus \{s_{j_1}, \dots, s_{j_N}\}$, after we have substituted $s_{j_1}, \dots, s_{j_N}$ using the relations $s_{I_1} = \dots = s_{I_N} = 0$. Then $I \in \mathcal{I}_N$ if, and only if, $L_{N,I} = 0$. Given the sets $\mathcal{I}_1, \dots, \mathcal{I}_N$, call a pair $(d, \mathbf{h})$, where $d \in \mathbb{Z}_{\geq 0}$ and $\mathbf{h} = \{h_{n,I} : 0 \leq n \leq N, I \in \mathcal{S}^*(2k)\}$ is a vector of non-negative integers, admissible if:

- $0 = h_{0,I} \leq h_{1,I} \leq \dots \leq h_{N,I}$ for $I \in \mathcal{S}^*(2k)$;
- If $I \in \mathcal{I}_n \setminus \mathcal{I}_{n-1}$ for some $n \in \{1, \dots, N\}$, then $h_{m,I} = h_{n,I}$ for all $m \geq n$;
- $X_N \geq N + d$, where

$$X_N := |\mathcal{I}_N \cap \mathcal{S}^+(2k)| - \sum_{I \in \mathcal{S}^-(2k) \cup (\mathcal{S}^+(2k) \setminus \mathcal{I}_N)} h_{N,I}.$$

We claim that the main contribution after N steps as the ones described above is a linear combination of terms of the form

$$\begin{aligned} J(I_1, \dots, I_N; \mathbf{h}, d) &:= \left(\frac{\varphi(q)}{q}\right)^{M_N} \frac{(\log R)^{X_N - N - d}}{(2i\pi)^{2k - N}} \\ &\times \int \dots \int_{\substack{\text{Re}(s_j) = \lambda_j / \log R \\ |\text{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k - N}} G_N(\mathbf{s}) R^{E_N(\mathbf{s})} \prod_{I \in \mathcal{S}^-(2k)} L^{(h_{N,I})}(1 + L_{N,I}(\mathbf{s}), \chi) \\ &\times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_N} \zeta_q^{(h_{N,I})}(1 + L_{N,I}(\mathbf{s})) ds_{2k-N} \dots ds_1 \end{aligned}$$

with $\lambda_j/\lambda_{j-1} \geq \lambda$,

$$M_N := \sum_{n=1}^N \sum_{I \in (\mathcal{I}_n \setminus \mathcal{I}_{n-1}) \cap \mathcal{S}^+(2k)} (h_{n-1,I} + 1),$$

$$E_N(\mathbf{s}) := L_{N,[2k]}(\mathbf{s}) \quad \text{and} \quad G_N(\mathbf{s}) := G(L_{N,\{1\}}(\mathbf{s}), \dots, L_{N,\{2k\}}(\mathbf{s})).$$

Note that the integral on the right hand side of (12.1) is the above integral when $N = 0$ and $h_{1,I} = 0$ for all I , with the conventions that $\mathcal{I}_0 = \{\emptyset\}$ and $L_{0,I}(s_1, \dots, s_{2k}) = s_I$.

Now, we need to understand what happens to J_N when we move some appropriate variable to the left. As in Section 8, we write

$$E_N(\mathbf{s}) = \gamma_1 s_1 + \dots + \gamma_{2k-N} s_{2k-N}$$

and let j_{N+1} denote the largest element of $\{1, \dots, 2k - N\}$, if it exists, for which $\gamma_{j_{N+1}} \neq 0$. We then separate four cases:

12.4. Case 1: $\text{Re}(E_N(\mathbf{s})) > 0$. In this case, $\gamma_{j_{N+1}} > 0$, by choosing a big enough λ . We make the change of variables

$$s'_j = s_j \quad (1 \leq j < j_{N+1}), \quad s'_j = s_{j+1} \quad (j_{N+1} \leq j \leq 2k - N), \quad s'_{2k-N} = s_{j_{N+1}},$$

and shift the s'_{2k-N} contour to the line $\text{Re}(s'_{2k-N}) = -\epsilon$ for a small enough ϵ . As we discussed, the contour contributions are negligible, and we are left with having to analyze the pole contributions. The poles occur when $L_{N,I_{N+1}}(\mathbf{s}') = 0$ for some $I_{N+1} \in \mathcal{S}^+(2k) \setminus \mathcal{I}_N$ such that the coefficient of s'_{2k-N} in $L_{N,I_{N+1}}$ is non-zero. Using the relation $L_{N,I_{N+1}}(\mathbf{s}') = 0$ to write s'_{2k-N} as a linear combination of $s'_1, \dots, s'_{2k-N-1}$, say $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$, we deduce that

$$L_{N,I}(s'_1, \dots, s'_{2k-N}) = L_{N,I}(s'_1, \dots, s'_{2k-N-1}, C(s'_1, \dots, s'_{2k-N-1})) =: L_{N+1,I}(s'_1, \dots, s'_{2k-N-1})$$

at this pole. We also set $E_{N+1} = L_{N+1,[2k]}$ and let $\mathcal{I}_{N+1}$ be the set of $I \subset [2k]$ such that $L_{N+1,I} = 0$. The generic order of the pole at $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$ is

$$(12.5) \quad m = \sum_{I \in (\mathcal{I}_{N+1} \setminus \mathcal{I}_N) \cap \mathcal{S}^+(2k)} (h_{N,I} + 1) - \nu,$$

where ν is the generic order of the zero of

$$G(\mathbf{s}) \prod_{I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N)} L^{(h_{N,I})}(1 + L_{N,I}(\mathbf{s}), \chi)$$

at the same point. In particular, $\nu = 0$ if $d = 0$ (so that $G(\mathbf{s}) = F(L_{N,\{1\}}(\mathbf{s}), \dots, L_{N,\{2k\}}(\mathbf{s}))$) and $h_{N,I} = 0$ for all $I \in \mathcal{S}^-(2k) \cap (\mathcal{I}_{N+1} \setminus \mathcal{I}_N)$. By a direct computation, we then find that

$$(12.6) \quad X_{2k-1} + m = |\mathcal{I}_{N+1} \cap \mathcal{S}^+(2k)| - \sum_{I \in \mathcal{S}^-(2k) \cup (\mathcal{S}^+(2k) \setminus \mathcal{I}_{N+1})} h_{N,I} - \nu.$$

We note that $m \geq 1$ for all $N \leq 2k - 1$ when $k \geq 2$ and $\nu = 0$; otherwise, we would have that $\mathcal{S}^+(2k) \subset \mathcal{I}_N$, which is impossible because the dimension of V_N is N , whereas the dimension of the span of the linear forms s_I , $I \in \mathcal{S}^+(2k)$ is $2k$.

We want to understand the pole contribution when $m \geq 1$. We separate two subcases:

Case 1a: $N = 2k - 1$. In this case, we have that $s'_j = L_{2k-1,\{j\}}(s'_1) = a_j s'_1$ for all j , where $a_j \in \mathbb{Q}$. Then the pole occurs necessarily when $s'_1 = 0$. Thus $\mathcal{I}_{2k} = \mathcal{S}(2k)$, and we obtain

an evaluation of $J_{2k-1}(\mathbf{I}; \mathbf{h})$ as finite linear combination of powers of $\log R$, the highest of which has exponent

$$X_{2k-1} + m - 2k = 2^{2k-1} - 2k - 1 - \sum_{I \in \mathcal{S}^-(2k)} h_{2k-1,I} - \nu,$$

up to an admissible error. The coefficients of the polynomial in $\log R$ are given in terms of the derivatives $L^{(j)}(1, \chi)$. Specifically, the coefficient of $(\log R)^{X_{2k-1}+m-2k-h}$, $0 \leq h \leq m-1$, is a linear combination of products of the form

$$\left(\frac{\varphi(q)}{q}\right)^{M_{2k}} \prod_{I \in \mathcal{S}^-(2k)} L^{(h_{2k,I})}(1, \chi),$$

with the coefficients of this linear combination being $\ll 1$, and with the parameters $h_{2k,I}$ satisfying $h_{2k,I} \geq h_{2k-1,I}$ with equality if $I \in \mathcal{I}_{2k-1} \setminus \{0\}$, and $\sum_{I \in \mathcal{S}^-(2k)} (h_{2k,I} - h_{2k-1,I}) \leq h$. Relations (12.3) and (12.4) and imply that

$$\left(\frac{\varphi(q)}{q}\right)^{M_{2k}} \prod_{I \in \mathcal{S}^-(2k)} L^{(h_{2k,I})}(1, \chi) \ll \frac{(\log q)^{O(1)}}{(\log Q)^{\#\{I \in \mathcal{S}^-(2k): h_{2k,I}=0\}}}.$$

Therefore, the contribution of this term of $J_{2k-1}(\mathbf{I}; \mathbf{h})$ is

$$\ll (\log R)^{X_{2k-1}+m-2k-h} \cdot \frac{(\log q)^{O(1)}}{(\log Q)^{\#\{I \in \mathcal{S}^-(2k): h_{2k,I}=0\}}} \leq (\log q)^{O(1)} \frac{(\log R)^{2^{2k-1}-2k-1-\sum_{I \in \mathcal{S}^-(2k)} h_{2k,I}}}{(\log Q)^{\#\{I \in \mathcal{S}^-(2k): h_{2k,I}=0\}}},$$

where we used the fact that $h \geq \sum_{I \in \mathcal{S}^-(2k)} (h_{2k,I} - h_{2k-1,I})$. Since $Q \geq R$, the above is maximized when $h_{2k,I} = 1$ for all $I \in \mathcal{S}^-(2k)$, whence

$$J_{2k-1}(\mathbf{I}; \mathbf{h}) \ll \frac{(\log q)^{O(1)}}{(\log R)^{2k+1}},$$

which is smaller than the claimed error term in Theorem 1.5(c).

Case 1b: $N \leq 2k-2$. Similarly to the Case 1b of Section 8.2, the contribution of the pole $s'_{2k-N} = C(s'_1, \dots, s'_{2k-N-1})$ to $J_N(\mathbf{I}, \mathbf{h})$ can be seen to be a linear combination of terms of the form

$$\begin{aligned} & \left(\frac{\varphi(q)}{q}\right)^{M_{N+1}} \frac{(\log R)^{X_N+m-h-N-1-d}}{(2i\pi)^{2k-N-1}} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=\lambda'_j/\log R, |\operatorname{Im}(s_j)| \leq T \\ 1 \leq j \leq 2k-N-1}} \tilde{G}(\mathbf{s}) R^{E_{N+1}(\mathbf{s})} \\ & \times \prod_{I \in \mathcal{S}^-(2k)} L^{(h_{N+1,I})}(1 + L_{N+1,I}(\mathbf{s}), \chi) \\ & \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{N+1}} \zeta_q^{(h_{N+1,I})}(1 + L_{N+1,I}(\mathbf{s})) ds_{2k-N-1} \cdots ds_1 \end{aligned}$$

plus an error term of size $O(1/\log R)$, where $h \in \{0, \dots, m-1\}$, $h_{N+1,I} \geq h_{N,I}$ with equality if $I \in \mathcal{I}_{N+1} \setminus \{0\}$, and $\sum_{I \in \mathcal{S}^-(2k) \cup (\mathcal{S}^+(2k) \setminus \mathcal{I}_{N+1})} (h_{N+1,I} - h_{N,I}) \leq h$. Relation 12.6 then implies that the power of $\log R$ is then $X_{N+1} - N - 1 - d'$ with

$$d' = d + \nu + h - \sum_{I \in \mathcal{S}^-(2k) \cup (\mathcal{S}^+(2k) \setminus \mathcal{I}_{N+1})} (h_{N+1,I} - h_{N,I}) \geq 0.$$

Moreover, $X_{N+1} - d' = X_N + m - h \geq N + 1$.

This completes the inductive step, which we keep iterating until until we arrive to one of Cases 2, 3 and 4 below, or to Case 1a above.

12.5. Case 2: $E_N(\mathbf{s}) = 0$ and $N = 2k - 1$. As in Case 1a, we have that $s_j = L_{2k-1, \{j\}}(s_1) = a_j s_1$ for all j , where $a_j \in \mathbb{Q}$. Then

$$\begin{aligned} J_{2k-1}(\mathbf{I}; \mathbf{h}, d) &= \left(\frac{\varphi(q)}{q} \right)^{M_{2k-1}} \frac{(\log R)^{X_{2k-1} - 2k + 1 - d}}{2\pi i} \prod_{I \in \mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1}} L^{(h_{2k-1, I})}(1, \chi) \\ &\quad \times \int_{\substack{\operatorname{Re}(s_1) = \lambda_1 / \log R \\ |\operatorname{Im}(s_1)| \leq T}} G_{2k-1}(s_1) \prod_{I \in \mathcal{S}^-(2k) \setminus \mathcal{I}_{2k-1}} L^{(h_{2k-1, I})}(1 + a_I s_1, \chi) \\ &\quad \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}} \zeta_q^{(h_{2k-1, I})}(1 + a_I s_1) ds_1, \end{aligned}$$

We first show a crude bound on $J_{2k-1}(\mathbf{I}; \mathbf{h}, d)$, that will allow us to focus on a more convenient subcase. We move the line of integration to $\operatorname{Re}(s_1) = \lambda / \log(qT)$ to find that

$$\begin{aligned} J_{2k-1}(\mathbf{I}; \mathbf{h}, d) &\ll (\log(q \log R))^{O(1)} \frac{(\log R)^{X_{2k-1} - 2k + 1 - d}}{(\log Q)^{\#\{I \in \mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1} : h_{2k-1, I} = 0\}}} \\ &= (\log(q \log R))^{O(1)} \frac{(\log R)^{|\mathcal{S}^+(2k) \cap \mathcal{I}_{2k-1}| - 2k + 1 - \sum_{I \in \mathcal{S}^-(2k) \cup (\mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1})} h_{2k-1, I} - d}}{(\log Q)^{\#\{I \in \mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1} : h_{2k-1, I} = 0\}}}, \end{aligned}$$

by relations (12.3) and (12.4). Since $Q \geq R$, if $d \geq 1$, if $h_{2k-1, I} \geq 2$ for some $I \in \mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1}$, or if $h_{2k-1, I} \geq 1$ for some $I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}$, then

$$\begin{aligned} J_{2k-1}(\mathbf{I}; \mathbf{h}, d) &\ll (\log(q \log R))^{O(1)} \cdot (\log R)^{|\mathcal{S}^+(2k) \cap \mathcal{I}_{2k-1}| - |\mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1}| - 2k} \\ &= (\log(q \log R))^{O(1)} \cdot (\log R)^{\sum_{I \in \mathcal{S}(2k)} (-1)^{|I|} - 2k - 1} \\ &\leq (\log(q \log R))^{O(1)} \cdot (\log R)^{\binom{2k}{k} - 2k - 1}, \end{aligned}$$

where we used Proposition 7.1. Therefore, these cases contribute to the claimed error term in Theorem 1.5(c).

Finally, assume that $d = 0$, $h_{2k-1, I} \in \{0, 1\}$ for all $I \in \mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1}$, and that $h_{2k-1, I} = 0$ for all $I \in \mathcal{S}(2k) \setminus \mathcal{I}_{2k-1}$. In this case,

$$\begin{aligned} J_{2k-1}(\mathbf{I}; \mathbf{h}, 0) &= \left(\frac{\varphi(q)}{q} \right)^{M_{2k-1}} \frac{(\log R)^{|\mathcal{S}^+(2k) \cap \mathcal{I}_{2k-1}| - 2k + 1}}{2\pi i} \prod_{I \in \mathcal{S}^-(2k) \cap \mathcal{I}_{2k-1}} \frac{L^{(h_{2k-1, I})}(1, \chi)}{(\log R)^{h_{2k-1, I}}} \\ &\quad \times \int_{\substack{\operatorname{Re}(s_1) = \lambda / \log R \\ |\operatorname{Im}(s_1)| \leq T}} G(s_1) \prod_{I \in \mathcal{S}^-(2k) \setminus \mathcal{I}_{2k-1}} L(1 + a_I s_1, \chi) \\ &\quad \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}} \zeta_q(1 + a_I s_1) ds_1. \end{aligned}$$

As before, we have the crude bound

$$J_{2k-1}(\mathbf{I}; \mathbf{h}, 0) \ll (\log(q \log R))^{O(1)} \cdot (\log R)^{\sum_{I \in \mathcal{I}_{2k-1}} (-1)^{|I|} - 2k}.$$

So, unless half of the a_j 's are $+1$ and the other half -1 , Proposition 7.1 implies that $J_{2k-1}(\mathbf{I}, \mathbf{h})$ contributes to the claimed error term in Theorem 1.5(c). So, assume that exactly half of the a_j 's are $+1$, and the other half are -1 . In particular, we find that $\mathcal{I}_{2k-1} \cap \mathcal{S}^-(2k) = \emptyset$ and that $|\mathcal{I}_{2k-1} \cap \mathcal{S}^+(2k)| = \binom{2k}{k} - 1$, so that

$$J_{2k-1}(\mathbf{I}; \mathbf{h}, 0) = \left(\frac{\varphi(q)}{q} \right)^{M_{2k-1}} \frac{(\log R)^{\binom{2k}{k}-2k}}{2\pi i} \int_{\substack{\operatorname{Re}(s_1)=\lambda/\log R \\ |\operatorname{Im}(s_1)| \leq T}} G(s_1) \prod_{I \in \mathcal{S}^-(2k)} L(1 + a_I s_1, \chi) \\ \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}} \zeta_q(1 + a_I s_1) ds_1.$$

Since $\nu = 0$ here, we saw before that the integrand has a genuine pole of order $m \geq 1$ at $s_1 = 0$ by a dimension argument. In fact, we have that $m \geq 2$: indeed, we know that $[2k] \in \mathcal{I}_{2k-1}$ by our assumption that $E_{2k-1} = 0$, so that $I \in \mathcal{I}_{2k-1}$ if and only if $[2k] \setminus I \in \mathcal{I}_{2k-1}$. In particular, since we know that there is at least one $I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}$, there must be at least two such I 's.

The presence of this pole make the estimation of $J_{2k-1}(\mathbf{I}, \mathbf{h})$ tricky. In particular, we cannot shift the contour to the line $\operatorname{Re}(s_1) = 0$ as in Case 2 of Section 8.2. Instead, we write

$$L(1 + a_I s_1, \chi) = L(\beta + a_I s_1, \chi) + R(a_I s_1),$$

where

$$R(s) := L(1 + s, \chi) - L(\beta + s, \chi).$$

We thus find that

$$R(s) = \int_{\beta}^1 L'(u + s, \chi) du \ll \frac{\log(q + |t|)}{\log Q} \quad (\sigma \geq -1/\log(q + |t|)),$$

by (12.3) and the assumption that $\beta > 1 - 1/(100 \log q)$. With this notation,

$$J_{2k-1}(\mathbf{I}; \mathbf{h}) = M + E,$$

where

$$M = \left(\frac{\varphi(q)}{q} \right)^{M_{2k-1}} \frac{(\log R)^{\binom{2k}{k}-2k}}{2\pi i} \int_{\substack{\operatorname{Re}(s_1)=\lambda/\log R \\ |\operatorname{Im}(s_1)| \leq T}} G_{2k-1}(s_1) \prod_{I \in \mathcal{S}^-(2k)} L(\beta + a_I s_1, \chi) \\ \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}} \zeta_q(1 + a_I s_1) ds_1 + E,$$

where E is a sum of similar expressions where at least one of the $L(\beta + a_I s_1, \chi)$ factors is replaced by $R(a_I s_1)$. Then moving s_1 to the line $\operatorname{Re}(s_1) = 1/\log(qT)$, we find that

$$E \ll \frac{(\log(q \log R))^{O(1)} (\log R)^{\binom{2k}{k}-2k}}{\log Q}.$$

Finally, we estimate M by moving the line of integration of s_1 to $\operatorname{Re}(s_1) = 0$, and use the fact that $G(s_1) \ll 1/(1 + |s_1|)^{2k}$ (note that the integrand is now analytic, since the pole of

the ζ_q 's is annihilated by the zeroes of the $L(\cdot, \chi)$'s to find that

$$J_{2k-1}(\mathbf{I}; \mathbf{h}) = \left(\frac{\varphi(q)}{q} \right)^{M_{2k-1}} \frac{(\log R)^{\binom{2k}{k}-2k}}{2\pi} \int_{-T}^T G(it) \prod_{I \in \mathcal{S}^-(2k)} L(\beta + ia_I t, \chi) \\ \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}} \zeta_q(1 + ia_I t) dt + O \left(\frac{(\log(q \log R))^{O(1)} (\log R)^{\binom{2k}{k}-2k}}{\log Q} \right).$$

Finally, we note that $G(s) = F(a_1 s, \dots, a_{2k} s)$ here, because $d = 0$. When $|t| \leq \log R$, we use (6.6) to replace $\hat{h}_R$ by R^s/s , and when $|t| > \log R$ we use the bound $\hat{h}_R(s) \ll R^\sigma/|s|$ by (6.5). Taking C to be large enough, we thus conclude that

$$\frac{J_{2k-1}(\mathbf{I}; \mathbf{h})}{(\log R)^{\binom{2k}{k}-2k}} = \frac{(\varphi(q)/q)^{M_{2k-1}}}{2\pi} \int_{-\infty}^{\infty} F(ia_1 t, \dots, ia_{2k} t) \prod_{I \in \mathcal{S}^-(2k)} L(\beta + ia_I t, \chi) \\ \times \prod_{I \in \mathcal{S}^+(2k) \setminus \mathcal{I}_{2k-1}} \zeta_q(1 + ia_I t) \prod_{j=1}^{2k} \frac{1 - 2^{-ia_j t}}{ia_j t} dt + O \left(\frac{(\log(q \log R))^{O(1)}}{\log R} \right).$$

Clearly, $J_{2k-1}(\mathbf{I}; \mathbf{h}, 0)$ for such $\mathbf{I}$ and $\mathbf{h}$ contributes towards the main term in Theorem 1.5(c).

12.6. Case 3: $\operatorname{Re}(E_N(s)) < 0$. This case is treated exactly like the corresponding case of Section 8, with the difference that the contours of $s_{2k-N}, s_{2k-N-1}, \dots, s_{j_{N+1}}$ are shifted to the lines $\operatorname{Re}(s_j) = \lambda^j / ((\log q) + (\log T)^{3/2})$, $j_{N+1} \leq j \leq 2k - N$. Keeping in mind that $q \leq e^{\sqrt{\log R}}$, a consequence of the hypotheses of Theorem 1.5(c), we find that

$$J_N(\mathbf{I}; \mathbf{h}) \ll e^{-(1+o(1))(\log \log R)^2}.$$

12.7. Case 4: $E(s) = 0$ and $N \leq 2k - 2$. As in the corresponding case of Section 8, we find that

$$J_N(\mathbf{I}; \mathbf{h}) \ll (\log(q \log R))^{O(1)} (\log R)^{X_N - N} \prod_{I \in \mathcal{S}^-(2k) \cap \mathcal{I}_N} L^{(h_{N,I})}(1, \chi) \\ \ll (\log(q \log R))^{O(1)} \frac{(\log R)^{|\mathcal{I}_N \cap \mathcal{S}^+(2k)| - N - \sum_{I \in \mathcal{S}^-(2k) \cup (\mathcal{S}^+(2k) \setminus \mathcal{I}_N)} h_{N,I}}}{(\log Q)^{\#\{I \in \mathcal{S}^-(2k) \cap \mathcal{I}_N : h_{N,I} = 0\}}},$$

by relations (12.3) and (12.4). Since $Q \geq R$, this is maximized when $h_{N,I} = 1$ for all $I \in \mathcal{S}^-(2k) \cap \mathcal{I}_N$ and $h_{N,I} = 0$ for all other I , so that

$$J_N(\mathbf{I}; \mathbf{h}) \ll (\log(q \log R))^{O(1)} (\log R)^{|\mathcal{I}_N \cap \mathcal{S}^+(2k)| - |\mathcal{I}_N \cap \mathcal{S}^-(2k)| - N}.$$

Then Proposition 7.1(c) implies that

$$J_N(\mathbf{I}; \mathbf{h}) \ll (\log(q \log R))^{O(1)} (\log R)^{\binom{2k}{k} - 2k - 2},$$

which contributes towards the claimed error term in Theorem 1.5(c).

12.8. Lower bounds. The argument presented above implies that, if $k \geq 2$ and $e^{(\log q)^2} \leq R \leq Q$, then

$$\mathcal{X}_{2k}(R) = c_k(\chi) \cdot (\log R)^{\binom{2k}{k} - 2k} + O\left((\log(q \log R))^{O(1)} (\log R)^{\binom{2k}{k} - 2k - 1}\right),$$

for some constant $c_k(\chi)$. Moreover, it is clear that $c_k(\chi) \ll (\log q)^{O(1)}$. In order to complete the proof of Theorem 1.5(c), we need to prove that $c_k(\chi) \gg (\log q)^{-O(1)}$ as well. In order to do so, we follow the argument of Section 9 and prove that, for any $\epsilon > 0$, there is a constant $c_k > 0$ such that

$$(12.7) \quad \mathcal{X}_{2k}(R) \geq c_k \frac{(\log R)^{\binom{2k}{k} - 2k - \epsilon}}{(\log q)^{O(1)}} - O_\epsilon\left((\log(q \log R))^{O(1)} (\log R)^{\binom{2k}{k} - 2k - 1}\right),$$

provided that $\sqrt{\log Q} \geq \log R \geq 2c_1 \log q$, where c_1 is the constant appearing in (12.3). (For this section, all constants will be independent of ϵ , unless specified by a subscript, as above.)

We set

$$y = \exp\{(\log R)^{1-\epsilon'}\} \quad \text{and} \quad Y = \exp\{(\log R)^{1-\epsilon'/2}\},$$

where ϵ' will be taken to be small enough in terms of ϵ , and focus our attention on integers of the form $n = ap_1 \cdots p_k$ with $P^+(a) \leq y$, $a \leq Y$, and $p_1, \dots, p_k$ are distinct primes such that $p_\ell > \sqrt{R}$ and $\chi(p_\ell) = -1$ for all $\ell \in \{1, \dots, k\}$. Then

$$\mathcal{X}_{2k}(R) \geq \frac{\prod_{p \leq R} (1 - 1/p)}{k!} \sum_{\substack{P^+(a) \leq y \\ a \leq Y}} \sum_{\substack{p_j > \sqrt{R} \\ \chi(p_j) = -1 \\ 1 \leq j \leq k}} \frac{\mu^2(p_1 \cdots p_k)}{ap_1 \cdots p_k} \left(\sum_{\ell=1}^k \sum_{\substack{R/(2p_\ell) < d \leq R/p_\ell \\ d|a}} \chi(d) \right)^{2k}.$$

The next step is to drop the condition that $a \leq Y$ by an application of Rankin's trick and to remove the condition that the p_j 's are distinct. We further replace the sharp cut-off $R/(2p_\ell) < d \leq R/p_\ell$ by the smooth cut-off $h(\frac{\log(dp_\ell)}{\log R}) - h(\frac{\log(2dp_\ell)}{\log R})$, where $h(x) = 1$ for $x \leq 1 - 1/(\log R)^B$ and $h(x) = 0$ for $x \geq 1$, with B is sufficiently large. To conclude, we have that

$$\begin{aligned} \mathcal{X}_{2k}(R) &\geq \frac{\prod_{p \leq R} (1 - 1/p)}{k! (\log R)^k} \sum_{P^+(a) \leq y} \sum_{\substack{\sqrt{R} < p_j \leq R \\ \chi(p_j) = -1 \\ 1 \leq j \leq k}} \frac{\prod_{j=1}^k \log p_j}{ap_1 \cdots p_k} \left(\sum_{\ell=1}^k \sum_{d|a} \chi(d) w\left(\frac{\log(dp_\ell)}{\log R}\right) \right)^{2k} \\ &\quad + O\left(\frac{1}{\log R}\right), \end{aligned}$$

where $w(x) = h(x) - h(x + \log 2 / \log R)$.

The rest of the proof follows the argument of Section 9, with a small twist, as we will explain in the end. We expand the $2k$ -th power and focus on a convenient subset of summands. We then conclude that

$$\mathcal{X}_{2k}(R) \geq \frac{\prod_{y < p \leq R} (1 - 1/p)}{k!} \sum_{\mathbf{J} \in \mathcal{J}} X(\mathbf{J}) + O\left(\frac{1}{\log R}\right)$$

with

$$X(\mathbf{J}) = \frac{1}{(\log R)^k} \sum_{\substack{P^+(d_j) \leq y \\ 1 \leq j \leq 2k}} \frac{\chi(d_1) \cdots \chi(d_{2k})}{[d_1, \dots, d_{2k}]} \prod_{\ell=1}^k \sum_{\sqrt{R} < p_\ell \leq R} \frac{(1 - \chi(p_\ell)) \log p_\ell}{2p_\ell} \prod_{j \in J_\ell} w\left(\frac{\log(p_\ell d_j)}{\log R}\right),$$

where $\mathbf{J}$ is as in Section 9. We set

$$S := \sum_{\sqrt{R} < p \leq R} \frac{(1 - \chi(p)) \log p}{2p} \asymp \log R$$

and write $\mathcal{L}$ for the set of $\ell \in \{1, \dots, k\}$ such that $J_\ell \neq \emptyset$. Then, using Perron's formula $2k$ times to write each appearance of w as an integral of $\widehat{w}_R$, we find that

$$\begin{aligned} X(\mathbf{J}) &= \frac{S^{k-|\mathcal{L}|} (\log R)^{-k}}{(2\pi i)^{2k}} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=1/\log R \\ 1 \leq j \leq 2k}} \sum_{\substack{P^+(d_j) \leq y \\ 1 \leq j \leq 2k}} \frac{\prod_{j=1}^{2k} \chi(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \left(\prod_{\ell \in \mathcal{L}} \sum_{p_\ell > \sqrt{R}} \frac{(1 - \chi(p_\ell)) \log p_\ell}{2p_\ell^{1+s_{J_\ell}}} \right) \\ &\quad \times \left(\prod_{j=1}^{2k} \widehat{w}_R(s_j) \right) ds_1 \cdots ds_{2k} + O\left(\frac{1}{\log R}\right), \end{aligned}$$

where $s_J = \sum_{j \in J} s_j$, as usual, and the condition that $p_\ell \leq R$ was dropped because it is encoded in the support of w . By possibly re-indexing the variables $s_1, \dots, s_{2k}$, we may assume that $\mathcal{L} = \{1, \dots, L\}$, where $L = |\mathcal{L}|$, and that $\max J_\ell = 2k - L + \ell$ for all $\ell \in \{1, \dots, L\}$ with $L = |\mathcal{L}|$. As in Section 9, we will move the variables $s_{2k-L+1}, \dots, s_{2k}$ to the left. We note that

$$\sum_{p > \sqrt{R}} \frac{(1 - \chi(p)) \log p}{p^{1+s}} = -\frac{\zeta'}{\zeta}(1+s) + \frac{L'}{L}(1+s, \chi) + O(1) - \sum_{p \leq R^{1/2}} \frac{\log p}{p^{1+s}}$$

for $\operatorname{Re}(s) \geq -1/3$. The above has simple poles at $s = 0$ and $s = \beta - 1$, each of residue 1. Therefore, using the argument leading to (9.11), we find that

$$\begin{aligned} (12.8) \quad X(\mathbf{J}) &= \sum_{\substack{\epsilon_\ell \in \{0, \beta-1\} \\ 1 \leq \ell \leq L}} \frac{S^{k-L} (\log R)^{-k}}{2^L (2\pi i)^{2k-L}} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=1/\log R \ (1 \leq j \leq 2k-L) \\ s_{J_\ell} = \epsilon_\ell \ (1 \leq \ell \leq L)}} \sum_{\substack{P^+(d_j) \leq y \\ 1 \leq j \leq 2k}} \frac{\prod_{j=1}^{2k} \mu(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]} \\ &\quad \times \left(\prod_{j=1}^{2k} \widehat{w}_R(s_j) \right) ds_1 \cdots ds_{2k-L} + O\left(\frac{1}{\log R}\right). \end{aligned}$$

The above expression is sufficient for handling the terms $\mathbf{J} \in \mathcal{J}$ with at least one J_ℓ of odd cardinality: following the argument of Section 9.5 with the obvious modifications implies that

$$(12.9) \quad X(\mathbf{J}) \ll (\log(q \log R))^{O(1)} \frac{(\log y)^{\binom{2k}{k} - 2k - 1 + L}}{(\log R)^L},$$

where we used the fact that $\sup_{q < p \leq Q} (1 + \chi(p))/p \ll 1$.

However, we need to be more careful on our lower bound for the main term, that is to say for $X(\mathbf{J})$ with $|J_\ell| = 2$ for all ℓ . First of all, by relabelling our variables, we may assume that $J_\ell = \{\ell, \ell + k\}$ for all ℓ . In our expression (12.8) for $X(\mathbf{J})$, we see that $s_{J_\ell} = \epsilon_\ell \in \{0, \beta - 1\}$

implies that $s_{\ell+k} = -s_\ell + O(1/\log Q)$. We want to replace $s_{\ell+k}$ by $-s_\ell$. This introduces an error that we will control by an application of the mean value theorem. In particular, we need to understand the derivative of the integrand. If

$$G(\mathbf{s}) = \sum_{\substack{P^+(d_j) \leq y \\ 1 \leq j \leq k}} \frac{\prod_{j=1}^{2k} \chi(d_j) d_j^{-s_j}}{[d_1, \dots, d_{2k}]},$$

then $-G'(\mathbf{s})/G(\mathbf{s})$ equals $\sum_{I \in \mathcal{S}^*(2k)} \sum_{p \leq y} \chi^{|I|}(p) \log p / p^{1+s_I}$, plus lower order terms, so that $G'(\mathbf{s}) \ll (\log y)G(\mathbf{s})$ for the vectors $\mathbf{s}$ we are considering. Similarly, $\widehat{w}_R(s + \epsilon)/R^{s+\epsilon} = \widehat{w}_R(s)/R^s + O(\epsilon/(|s| + 1))$ by (6.5). Since we also have that

$$\int \cdots \int_{\substack{\operatorname{Re}(s_j)=1/\log R \\ |s_{j+k}+s_j| \leq 1-\beta \\ 1 \leq j \leq k}} |G(\mathbf{s})| \prod_{j=1}^k \frac{|ds_j|}{1 + |s_j|^2} \ll (\log(q \log R))^{O(1)} \frac{(\log y)^{\binom{2k}{k}-k}}{(\log R)^k},$$

by the argument leading to (12.9), we conclude that

$$\begin{aligned} X(\mathbf{J}) &= \frac{(1 + R^{\beta-1})^k}{(2\pi i \log R)^k} \int \cdots \int_{\substack{\operatorname{Re}(s_j)=1/\log R \\ s_{j+k}=-s_j \\ 1 \leq j \leq k}} G(\mathbf{s}) \left(\prod_{j=1}^{2k} \widehat{w}_R(s_j) \right) ds_1 \cdots ds_k \\ &\quad + O \left(\frac{\log y}{\log Q} \cdot \frac{(\log(q \log R))^{O(1)} (\log y)^{\binom{2k}{k}-k}}{(\log R)^k} \right). \end{aligned}$$

The main term can now be bounded from below as in Section 9.4. We thus arrive to the lower bound

$$X(\mathbf{J}) \geq c_k \frac{(\log y)^{\binom{2k}{k}-k} (\log R)^{-k}}{(\log q)^{O(1)}} - O \left(\frac{(\log(q \log R))^{O(1)} (\log y)^{\binom{2k}{k}-k+1}}{(\log R)^{k+2}} \right),$$

using that $\sum_{q < p \leq Q} (1 + \chi(p))/p \ll 1$ and $y \leq R \leq e^{\sqrt{\log Q}}$ here. This completes the proof of (12.7), and thus of Theorem 1.5(c).

REFERENCES

- [1] M. Balazard, M. Naimi and Y.-F. S. Pétermann, *Étude d'une somme arithmétique multiple liée à la fonction de Möbius*, Acta Arith. **132** (2008), 245–298.
- [2] R. de la Bretèche, *Estimation de sommes multiples de fonctions arithmétiques*, Compos. Math. **128** (2001), 261–298.
- [3] Harold Davenport. *Multiplicative number theory*, volume 74 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2000. Revised and with a preface by Hugh L. Montgomery.
- [4] F. Dress, H. Iwaniec and G. Tenenbaum, *Sur une somme liée à la fonction de Möbius*, J. Reine Angew. Math. **340** (1983), 53–58.
- [5] S. Eberhard, K. Ford and B. Green, *Permutations fixing a k -set*, IMRN, to appear. arXiv:1507.04465
- [6] P. Erdős, and R. R. Hall, *On the Möbius function*, J. Reine Angew. Math. **315** (1980), 121–126.
- [7] K. Ford, *The distribution of integers with a divisor in a given interval*, Ann. of Math. **168**, (2008), 367–433.
- [8] J. Friedlander and H. Iwaniec, *Opera de cribro*. American Mathematical Society Colloquium Publications, 57. American Mathematical Society, Providence, RI, 2010.

- [9] A. Granville, *Cycle lengths in a permutation are typically Poisson*. Electron. J. Combin. 13 (2006), no. 1, Research Paper 107, 23 pp.
- [10] R. R. Hall and G. Tenenbaum, *Divisors*. Cambridge Tracts in Mathematics, 90. Cambridge University Press, Cambridge, 1988.
- [11] A. Goldston, J. Pintz and C. Y. Yıldırım, *Primes in tuples. I*. Ann. of Math. (2) 170 (2009), no. 2, 819–862.
- [12] H. Iwaniec and E. Kowalski, *Analytic number theory*. American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.
- [13] D. Koukoulopoulos, *Localized factorizations of integers*. Proc. Lond. Math. Soc. (3) 101 (2010), no. 2, 392–426.
- [14] —, *Pretentious multiplicative functions and the prime number theorem for arithmetic progressions*. Compos. Math., 149 (2013), no. 7, 1129–1149.
- [15] —, *On multiplicative functions which are small on average*. Geom. Funct. Anal. 23 (2013), no. 5, 1569–1630.
- [16] J. Maynard, *Small gaps between primes*, Ann. of Math. 181 (2015), 383–413.
- [17] Y. Motohashi, , *A multiple sum involving the Möbius function*, Publ. Inst. Math. (Beograd) (N.S.) **76(90)** (2004), 31–39.
- [18] , M. Rosen, *Number theory in function fields*. Graduate Texts in Mathematics, 210. Springer-Verlag, New York, 2002.
- [19] A. Selberg, *On an elementary method in the theory of primes*, Norske Vid. Selsk. Forh., Trondhjem 19 (1947), 64–67.
- [20] A. Selberg, *Collected papers. II*, Springer Collected Works in Mathematics, Springer, Heidelberg, 2014.
- [21] T. Tao, *Polymath8b: Bounded intervals with many primes, after Maynard*, Blog note. <https://terrytao.wordpress.com/2013/11/19/polymath8b-bounded-intervals-with-many-primes-after-maynard/>
- [22] ———, *Introduction to analytic and probabilistic number theory*. Translated from the second French edition (1995) by C. B. Thomas. Cambridge Studies in Advanced Mathematics, 46. Cambridge University Press, Cambridge, 1995.
- [23] E. C. Titchmarsh, *The theory of the Riemann zeta-function*. Second edition. Edited and with a preface by D. R. Heath-Brown. The Clarendon Press, Oxford University Press, New York, 1986.
- [24] Y. Zhang, *Bounded gaps between primes*, Ann. of Math. (2) 179 (2014), no. 3, 1121–1174.

AG: DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP 6128 SUCC. CENTRE-VILLE, MONTRÉAL, QC H3C 3J7, CANADA; AND DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON WC1E 6BT, ENGLAND.

E-mail address: `andrew@dms.umontreal.ca`

DK: DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP 6128 SUCC. CENTRE-VILLE, MONTRÉAL, QC H3C 3J7, CANADA

E-mail address: `koukoulo@dms.umontreal.ca`

JM: MATHEMATICAL INSTITUTE, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD OX2 6GG, ENGLAND

E-mail address: `james.alexander.maynard@gmail.com`